

AD-A182 804

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

DTIC FILE COPY

②

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
AFOSR-TN 87-0813		2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
TITLE (and Subtitle) Infrared Emission and Atomic Transitions		5. TYPE OF REPORT & PERIOD COVERED Final Technical Report 1 August 1983-31 January 1987	
AUTHOR(s) Chun C. Lin		6. PERFORMING ORG. REPORT NUMBER	
PERFORMING ORGANIZATION NAME AND ADDRESS Department of Physics University of Wisconsin-Madison 1150 University Ave., Madison, WI 53706		8. CONTRACT OR GRANT NUMBER(s) AFOSR-83-0312	
CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research, NP Bolling Air Force Base, D.C. 20332		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 611025, 2301, A4	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) SCMTC CW 11		12. REPORT DATE March 27, 1987	
		13. NUMBER OF PAGES 4	
		15. SECURITY CLASS. (of this report) Unclassified	
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release, distribution unlimited			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Approved for public release, distribution unlimited			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) infrared emission; atoms in highly excited levels, number density of metastable atoms			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The general objective of this project is to study radiative transitions in atoms and molecules, particularly those relevant to infrared radiation. Our major efforts include (a) study of electron-impact dissociative excitation of oxygen and its mechanism, (b) production of excited nitrogen atoms and their infrared emission by electron impact on nitrogen molecules, (c) development of a method for determining number density of metastable atoms produced by electron beams, (d) study of collisional behaviors of 3s lone electrons.			

DTIC
ELECTE
JUL 06 1987
S E D

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

816

Final Scientific Report

Grant AFOSR-83-0312, 1 August 1983 - 31 January 1987

Principal Investigator: Chun C. Lin

The general objective of this project is to study radiative transitions in atoms and molecules, particularly those which are relevant to infrared radiation. Our major efforts were in the following areas:

(i) Passage of an electron beam to a chamber containing O_2 molecules produces excited O_2 molecules in bound electronic states as well as unbound states. In the latter case the excited molecule dissociates into two oxygen atoms with one or both in excited states. To study this process we measure the radiation from the excited oxygen atoms produced by electron impact on O_2 molecules. We have measured absolute optical emission cross sections for a large number of transitions originating from excited electron configurations $1s^2 2s^2 2p^3 n\ell$ ($n=3,4,5,6,7,8$) of the oxygen atom produced by incident electron energy from threshold to 500 eV. (Production of oxygen atoms in the highly excited levels is of special interest because such highly excited atoms are sources of infrared radiation.) Studies of detailed characteristics of the energy dependence of the cross sections (including the appearance potential) allow us to identify the key mechanisms for producing the excited oxygen atoms in the electron-impact dissociation experiment. The excitation functions show a broad peak at about 90 eV with a shoulder-like structure of varying degree near 35 eV. The observed threshold energy is very close to the energy defect of the dissociation process. Near the threshold, dissociative excitation through partly bound Rydberg states of the oxygen molecule is believed to be a major mechanism for producing the observed atomic-oxygen emission, whereas simultaneous ionization and excitation followed by dissociation becomes important at energies

above 50 eV. Absolute optical emission cross sections for a series of atomic-oxygen transitions in the long-wave infrared region have been obtained.

(ii) Production of excited nitrogen atoms $1s^2 2s^2 2p^2 n\ell$ in the low excited states ($n=3,4$) by electron-impact dissociation was previously reported in the literature. Nitrogen atoms in the highly excited states are of special importance because of the intrinsic interest in atomic physics and their potential roles as infrared emitters. Therefore we have conducted a study of production of excited nitrogen atoms in the highly excited states of $n=5,6,7,8,9$. Excitation cross sections and excitation functions for optical emission of some fifty transitions of the type $1s^2 2s^2 2p^2 n\ell \rightarrow 1s^2 2s^2 2p^2 n'\ell'$ have been measured. Like the case of oxygen, the cross section data, especially the variation of the cross section with incident electron energy, provide information about the key mechanisms for the production of the excited nitrogen atoms. Absolute optical cross sections for a series of atomic-nitrogen transitions in the long-wave infrared region have also been obtained.

(iii) A new method for determining the number density of metastable atoms produced by electron-beam excitation of ground-state atoms has been developed in our laboratory and applied successfully to neon. We generate metastable Ne atoms (in the $1s^2 2s^2 2p^5 3s$ electron configuration) by an electron beam through a container of Ne atoms. A pulsed laser, tuned to the absorption frequency of one of the $1s^2 2s^2 2p^5 3s \rightarrow 1s^2 2s^2 2p^5 3p$ transitions, pumps the metastable atoms to a $1s^2 2s^2 2p^5 3p$ level. With a high-power pulsed laser we saturate the transition, i.e., transfer the atoms from the $1s^2 2s^2 2p^5 3s$ metastable level to the higher $1s^2 2s^2 2p^5 3p$ level to equalize the population of these two levels. The number of photons emitted from this $1s^2 2s^2 2p^5 3p$ level after cessation of the laser pulse is measured and used to determine the number density of the metastable Ne atom number density. At a Ne gas pressure of 15 mTorr, an electron beam current

density of 0.016 A/cm^2 , and an electron beam energy of 100 eV, we find the number density of the metastable Ne atoms in the $1s^2 2s^2 2p^5 3s$, $J=2$ level to be $9 \times 10^8 \text{ cm}^{-3}$. Comparing this with the ground-state atom number density of $5 \times 10^{14} \text{ cm}^{-3}$, we find a metastable atom concentration of about one or two parts in 10^6 .

(iv) As a way to understand the collisional behaviors of the $\text{Ne}(1s^2 2s^2 2p^5 3s)$ metastable atoms, we try to have some indications as to the roles played by the 3s electron under various collision conditions. The 3s electron is a "lone" electron in an outer shell and relatively loosely bound, so it is in some way similar to the 3s electron in a ground-state Na atom ($1s^2 2s^2 2p^6 3s$). Because of its small binding energy, the 3s electron can be readily excited to the higher levels, ionized, or, upon colliding with another atom, may transfer to the collision partner. Thus we have conducted collision experiments of Na atoms with H^+ , H_2^+ , H_3^+ , H^- ions and neutral H atoms as projectiles, and measured the cross sections of $3s \rightarrow 3d$ excitation and of electron transfer, $\text{H} + \text{Na} \rightarrow \text{H}^- + \text{Na}^+$. At the same projectile velocity, the $\text{Na}(3s \rightarrow 3d)$ excitation cross sections are the same for H^+ , H_2^+ , H_3^+ , and the same for electron projectile when the projectile velocity is somewhat above the electron excitation threshold. However, the $\text{Na}(3s \rightarrow 3d)$ excitation cross sections due to H^- ion impact are about 45% lower than the corresponding cross sections due to H^+ , H_2^+ , H_3^+ , and electron impact at the same projectile velocity. These results provide very valuable information about the collisional properties of the various partners.



Accession For		☒ ☐ ☐	
NTIS GRA&I			
DTIC TAB			
Unannounced			
Justification			
By			
Distribution/			
Availability Codes			
Avail and/or			
Special			
Dist		A-1	

Publications

"Excitation of the Na(3p) Level by H Ions and H⁰ Atoms", Physical Review Letters 51, 2029 (1983).

"Charge Changing Cross Sections for 1-25 keV H(1s) Incident on a Na Vapor Target", Physical Review A 29, 1083 (1984).

"Electron-Impact Excitation of the $D^3\Sigma_u^+$ and $c_4^1\Sigma_u^+$ Rydberg States of N₂", Physical Review A 29, 1709 (1984).

"Emission from Oxygen Atoms Produced by Electron-Impact Dissociative Excitation of Oxygen Molecules", Physical Review A 32, 2100 (1985).

"Electron Excitation Cross Sections for the Metastable and Resonant Levels of Ne(2p⁵3s)", Physical Review A 32, 2117 (1985).

"Emission from Atomic Nitrogen Produced by Electron Impact on Nitrogen Molecules", submitted to the Journal of Chemical Physics.

Professional Personnel

Researchers associated with this project include the following:

(a) Senior researchers: C. C. Lin (Principal Investigator), L. W. Anderson, F. A. Sharpton.

(b) Graduate Students: J. S. Allen, A. R. Filippelli, J. E. Gastineau, A. M. Howald, M. B. Schulman, D.L. A. Rall.

Interactions

The principal investigator has regular scientific contact with Drs. Edward T. P. Lee and Randall Murphy in the Radiation Effect Branch, Optical Physics Division of the Air Force Geophysics Laboratory (Bedford, Massachusetts).

Approved for public release;
distribution unlimited.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSO)
NOTICE OF TRANSMITTAL TO DTIC

This technical report has been reviewed and is
approved for public release IAW AFR 190-12.
Distribution is unlimited.

MATTHEW J. KERPER
Chief, Technical Information Division