

Unclassified

DTIC FILE COPY

②

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS	
AD-A182 809		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited	
		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 87-0909	
6a. NAME OF PERFORMING ORGANIZATION Universities Space Research Association	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION AFOSR/NM	
6c. ADDRESS (City, State and ZIP Code) NASA Langley Research Center Hampton, Va 23665-5225		7b. ADDRESS (City, State and ZIP Code) Bldg 410 Bolling AFB DC 20332-6448	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR	8b. OFFICE SYMBOL (If applicable) NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-83-0089	
8c. ADDRESS (City, State and ZIP Code) Bldg 410 Bolling AFB DC 20332-6448		10. SOURCE OF FUNDING NO.	
		PROGRAM ELEMENT NO. 61102F	PROJECT NO. 2304
		TASK NO. A3	WORK UNIT NO.
11. TITLE (Include Security Classification) Spectral methods for time dependent partial differential equations			
12. PERSONAL AUTHOR(S) David Gottlieb			
13a. TYPE OF REPORT Final	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Yr., Mo., Day) 86 11 20	15. PAGE COUNT 5
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB. GR.	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)			
The investigator proved research on the use of spectral methods in computational fluid dynamics. In particular, he looked at their implementation for the solution of time dependent partial differential equations. Other topics pursued included the adaption spectral methods for compressible flow problems involving shocks, and the exploration of information content in spectral calculations.			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT. ☐ DTIC USERS ☐		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL John Thomas, Maj, USAF	22b. TELEPHONE NUMBER (Include Area Code) (202) 767-5027	22c. OFFICE SYMBOL NM	

DTIC
ELECTE
S JUL 30 1987 **D**
E

87 7 28 399

FINAL SCIENTIFIC REPORT

GRANT 83-0089

PRINCIPAL INVESTIGATOR

DAVID GOTTLIEB

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

→ This document discusses briefly
The following ^{research} topics were researched:

- 1) Spectral methods for time dependent partial differential equations.

The theory of spectral methods for time dependent partial equations is reviewed. When the domain is periodic, Fourier methods are presented while for nonperiodic problems both Chebyshev and Legendre methods are discussed. The theory is presented for both hyperbolic and parabolic systems using both Galerkin and collocation procedures. While most of the review considers problems with constant coefficients, the extension to nonlinear problems is also discussed. Some results for problems with shocks are presented. (See Appendix 1)

ASPECTED

- 2) Spectral methods for compressible flow problems.

In this article we review recent results concerning numerical simulation of shock waves using spectral methods. We discuss shock fitting techniques as well as shock capturing techniques with finite difference artificial viscosity. We also discuss the notion of the information contained in the numerical results obtained by spectral methods and show how this information can be recovered. (See Appendix 2)

P 2

3) Recovering pointwise values of discontinuous data within spectral accuracy.

We show how pointwise values of a function, $f(x)$, can be accurately recovered either from its spectral or pseudospectral approximations, so that the accuracy solely depends on the local smoothness of f in the neighborhood of the point x . Most notably, given the equidistant function grid values, its intermediate point values are recovered within spectral accuracy, despite the possible presence of discontinuities scattered in the domain. (Recall that the usual spectral convergence rate decelerates otherwise to first order, throughout.)

To this end we employ a highly oscillatory smoothing kernel in contrast to the more standard positive unit-mass mollifiers.

In particular, post-processing of a stable Fourier method applied to hyperbolic equations with discontinuous data recovers the exact solution modulo a spectrally small error. Numerical examples are presented. (See Appendix 3)

4) Information content in spectral calculations.

Spectral solutions of hyperbolic partial differential equations induce a Gibbs phenomenon near local discontinuities or strong gradients.

A procedure is presented for extracting the piecewise smooth behavior of the solution out of the oscillatory numerical solution data. The procedure is developed from the theory of linear partial differential equations. Its application to a non-linear system (the two-dimensional Euler equations of gas dynamics) is shown to be efficacious for the particular situation. (See Appendix 4)

5) Spectral methods for discontinuous problems.

We show that spectral methods yield high-order accuracy even when applied to problems with discontinuities, though not in the sense of pointwise accuracy. Two different procedures are presented which recover pointwise accurate approximations from the spectral calculations. (See Appendix 5)

6) Stability and Lyapunov stability of dynamical systems: a differential approach and a numerical method. *and* *→* *See p. 4*

A set of differential equations for the eigenvalues and eigenvectors of the stability matrix of a dynamical system, as well as for the Lyapunov exponents and the corresponding eigenvectors is derived. The rate of convergence of the Lyapunov eigenvectors is shown to be exponential. The eigenvectors of the stability matrix can be grouped into sets, each spanning a subspace which converges at an exponential rate. It is demonstrated that, generically, the real parts of the eigenvalues of the stability matrix equal the corresponding Lyapunov exponents. This statement has been tested numerically. The values of the Lyapunov exponents, μ_1 , are shown to be related to the corresponding finite time values of the Lyapunov exponents (e.g. those deduced from a finite time numerical simulation), $\mu_1(t)$, by: $\mu_1(t) = \mu_1 + \frac{b_1 + \xi_1(t)}{t}$. The b_1 's are constants and $\xi_1(t)$ are "noise" terms of zero mean. This observation leads to a method of extrapolation, which has been used to predict Lyapunov exponents from a finite amount of data. It is shown that the use of the standard (numerical) methods to compute Lyapunov exponents introduces an error $\frac{a_1}{t}$ in the value of $\mu_1(t)$, where the a_1 's are

constants. Thus the standard method has a rate of convergence which is the same as that of the exact $\mu_1(t)$'s. Finally, we have shown how one can compute the eigenvectors associated with each of the eigenvalues of the stability matrix as well as the Lyapunov eigenvectors. (See Appendix 6)

✓ 7) Boundary conditions for incompressible flows. ①

In this paper, we have analyzed the effect of boundary conditions on incompressible flows. We have explained and analyzed a numerical boundary layer of thickness $\sqrt{\nu \Delta t}$ that appears in many formulations of incompressible flow problems and we have explained techniques for the development of high-order methods.

For first- or second-order methods, we recommend the use of splitting methods in the form (4.1) - (4.5) or (7.1) - (7.5) with the tangential-derivative boundary conditions (7.16) or the modified velocity boundary conditions (7.28). To achieve higher-order accuracy, we may employ either the extrapolation methods outlined in Sec. 6 or, perhaps even better, we may use schemes of the form (1.11) with high-order pressure boundary conditions of the form (7.23) or iterative conditions of the form (7.25) [with only a few iterations per time step].

In this paper, we have shown that we can characterize methods for the solution of incompressible flow problems as belonging to either parabolic or elliptic type with regard to the determination of the pressure field. The elliptic schemes typically have smaller errors in the divergence field, with the errors decaying exponentially away from the boundaries of

the computational domain. On the other hand, the parabolic schemes have smooth solutions, without numerical boundary layers, but care should be exercised with respect to the boundary conditions in order the initial divergence errors be eliminated. This analysis explains why 'elliptic' schemes, like that introduced by Harlow & Welch (1965) have been found to be more accurate than parabolic schemes.

We have also shown, using Weyl's lemma for the decomposition of a flow into its solenoidal and irrotational components, that it is possible to derive accurate boundary conditions for the pressure that involve only the tangential derivative of the boundary vorticity. This form of the boundary condition tends to minimize the effects of numerical boundary layers induced by splitting methods. (See Appendix 7)

The following papers were prepared under Grant 83-0089 and appeared in the following journals:

Gottlieb, D. and E. Turkel: Spectral methods for time dependent partial differential equations. Proc. of 3rd 1983 C.I.M.E. Series Conference on Numerical Methods in Fluid Dynamics, Como, Italy, July 7-12, 1983. Lecture Notes in Math., Vol. 1127, Springer-Verlag, 1985, pp. 115-155.

Gottlieb, D.: Spectral methods for compressible flow problems. Lecture Notes in Physics, No. 218, (Soubbaramayer and J. P. Boujot, eds.), Springer-Verlag, 1984, Proc. of 9th Intl. Conference on Numerical Methods in Fluid Dynamics, Saclay, France, pp. 48-61.

Gottlieb, D. and E. Tadmor: Recovering pointwise values of discontinuous data within spectral accuracy. Progress and Supercomputing in Computational Fluid Dynamics, (E. M. Murman and S. S. Abarbanel, eds.), Birkhauser Boston, Inc., 1985, pp. 357-375.

Abarbanel, S. and D. Gottlieb: Information content in spectral calculations. Progress and Supercomputing in Computational Fluid Dynamics, (E. M. Murman and S. S. Abarbanel, eds.), Birkhauser Boston, Inc. 1985, pp. 345-356.

Abarbanel, S., D. Gottlieb, and E. Tadmor: Spectral methods for discontinuous problems. Proc. of Conference on Numerical Methods for Fluid Dynamics, Reading, England.

Goldhirsch, I., P. Sulem, S. Orszag: Stability and Lyapunov stability of dynamical systems: a differential approach and a numerical method. J. Sci. Comp., to appear.

Orszag, S. A., M. Israeli, and M. Deville: Boundary conditions for incompressible flows. J. Sci. Comp., to appear.