

AD-A183 878

SPECIES PROFILES LIFE HISTORIES AND ENVIRONMENTAL
REQUIREMENTS OF COASTAL (U) GULF COAST RESEARCH LAB
OCEAN SPRINGS MS F C SUTTER ET AL. MAR 87 FHS-82/11.71

1/1

UNCLASSIFIED

F/G 8/1

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A


FWS
Biological Report 82/11.71
March 1987

WES/TR/EL-82-4.71

**Species Profiles: Life Histories and
Environmental Requirements of Coastal Fishes
and Invertebrates (Gulf of Mexico)**

JUL 16 1987

PIGFISH



**Fish and Wildlife Service
U.S. Department of the Interior**

**Coastal Ecology Group
Waterways Experiment Station
U.S. Army Corps of Engineers**

Biological Report 82(11.71)
TR EL-82-4
March 1987

**Species Profiles: Life Histories and Environmental Requirements
of Coastal Fishes and Invertebrates (Gulf of Mexico)**

PIGFISH

by

Frederick C. Sutter and Thomas D. McIlwain
Gulf Coast Research Laboratory
Fisheries Research and Development
East Beach Boulevard
Ocean Springs, MS 39564

Project Manager
Carroll Cordes

Project Officer
David Moran
National Wetlands Research Center
U.S. Fish and Wildlife Service
1010 Gause Boulevard
Slidell, LA 70458

Performed for
Coastal Ecology Group
Waterways Experiment Station
U.S. Army Corps of Engineers
Vicksburg, MS 39180

and

National Wetlands Research Center
Research and Development
Fish and Wildlife Service
U.S. Department of Interior
Washington, DC 20240



A1

This series may be referenced as follows:

U.S. Fish and Wildlife Service. 1983-19 . Species profiles: life histories and environmental requirements of coastal fishes and invertebrates. U.S. Fish Wildl. Serv. Biol. Rep. 82(11). U.S. Army Corps of Engineers, TR EL-82-4.

This profile may be cited as follows:

Sutter, F. C., and T. D. McIlwain. 1987. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (Gulf of Mexico)--pigfish. U.S. Fish Wildl. Serv. Biol. Rep. 82(11.71). U.S. Army Corps of Engineers, TR EL-82-4. 11 pp.

PREFACE

This species profile is one of a series on coastal aquatic organisms, principally fish, of sport, commercial, or ecological importance. The profiles are designed to provide coastal managers, engineers, and biologists with a brief comprehensive sketch of the biological characteristics and environmental requirements of the species and to describe how populations of the species may be expected to react to environmental changes caused by coastal development. Each profile has sections on taxonomy, life history, ecological role, environmental requirements, and economic importance, if applicable. A three-ring binder is used for this series so that new profiles can be added as they are prepared. This project is jointly planned and financed by the U.S. Army Corps of Engineers and the U.S. Fish and Wildlife Service.

Suggestions or questions regarding this report should be directed to one of the following addresses.

Information Transfer Specialist
National Wetlands Research Center
U.S. Fish and Wildlife Service
NASA-Slidell Computer Complex
1010 Gause Boulevard
Slidell, LA 70458

or

U.S. Army Engineer Waterways Experiment Station
Attention: WESER-C
Post Office Box 631
Vicksburg, MS 39180

CONVERSION TABLE

Metric to U.S. Customary

<u>Multiply</u>	<u>By</u>	<u>To Obtain</u>
millimeters (mm)	0.03937	inches
centimeters (cm)	0.3937	inches
meters (m)	3.281	feet
meters (m)	0.5468	fathoms
kilometers (km)	0.6214	statute miles
kilometers (km)	0.5396	nautical miles
square meters (m ²)	10.76	square feet
square kilometers (km ²)	0.3861	square miles
hectares (ha)	2.471	acres
liters (l)	0.2642	gallons
cubic meters (m ³)	35.31	cubic feet
cubic meters (m ³)	0.0008110	acre-feet
milligrams (mg)	0.00003527	ounces
grams (g)	0.03527	ounces
kilograms (kg)	2.205	pounds
metric tons (t)	2205.0	pounds
metric tons (t)	1.102	short tons
kilocalories (kcal)	3.968	British thermal units
Celsius degrees (°C)	1.8(°C) + 32	Fahrenheit degrees

U.S. Customary to Metric

inches	25.40	millimeters
inches	2.54	centimeters
feet (ft)	0.3048	meters
fathoms	1.829	meters
statute miles (mi)	1.609	kilometers
nautical miles (nmi)	1.852	kilometers
square feet (ft ²)	0.0929	square meters
square miles (mi ²)	2.590	square kilometers
acres	0.4047	hectares
gallons (gal)	3.785	liters
cubic feet (ft ³)	0.02831	cubic meters
acre-feet	1233.0	cubic meters
ounces (oz)	283.5	milligrams
ounces (oz)	28.35	grams
pounds (lb)	0.4536	kilograms
pounds (lb)	.00045	metric tons
short tons (ton)	0.9072	metric tons
British thermal units (Btu)	0.2520	kilocalories
Fahrenheit degrees (°F)	0.5556(°F - 32)	Celsius degrees

CONTENTS

	<u>Page</u>
PREFACE	iii
CONVERSION TABLE	iv
ACKNOWLEDGMENTS	vi
 NOMENCLATURE/TAXONOMY/RANGE	 1
MORPHOLOGY/IDENTIFICATION AIDS	1
REASON FOR INCLUSION IN THE SERIES	3
LIFE HISTORY	3
Spawning	3
Eggs	3
Larvae	3
Juveniles	4
Adults	5
GROWTH CHARACTERISTICS	5
THE FISHERY	5
ECOLOGICAL ROLE	5
Food Habits	5
Predators	6
Parasites and Diseases	6
ENVIRONMENTAL REQUIREMENTS	6
Temperature and Salinity	6
Dissolved Oxygen	8
Substrate	8
 LITERATURE CITED	 9

ACKNOWLEDGMENTS

We gratefully acknowledge peer reviews by R.L. Shipp, University of South Alabama, Mobile, and H.D. Hoese, University of Southwestern Louisiana, Lafayette.




Figure 1. Pigfish (adult).

PIGFISH

NOMENCLATURE/TAXONOMY/RANGE

Scientific name Orthopristis chrysoptera (Linnaeus)

Preferred common name Pigfish (Figure 1)

Class Osteichthyes

Order Perciformes

Family Haemulidae

Geographical range: Pigfish are distributed along the Atlantic coast from Massachusetts (Hoes and Moore 1977) to the tip of Florida, but are uncommon north of Virginia (Breder 1948). They are found throughout the Gulf of Mexico (Figure 2; Briggs 1958) and in coastal waters around Bermuda.

ovate-elliptical, considerably compressed, its depth contained 2.6 to 3.0 times in standard length (SL). Posterior edge of upper jaw not reaching to below eye; two pores and a median groove on chin; jaws with a narrow band of slender teeth; preopercular margin very slightly serrate; gill rakers short and slender, about 12 on lower limb of first arch. Dorsal fin with 12 or 13 spines and 15 or 16 soft rays; anal fin with 3 spines and 12 or 13 soft rays; dorsal and anal fin spines enclosed in a deep scaly sheath, the soft rays naked. Scales ctenoid; pored lateral-line scales 55 to 58; 10 longitudinal rows of scales above the lateral line and 19 rows below.

MORPHOLOGY/IDENTIFICATION AIDS

The following descriptive characteristics for pigfish were taken from Courtenay and Sahlman (1978): Body

Color of body: light blue-gray above, shading gradually into silver below; each scale of body with a blue center, the edge with a bronze spot; these spots form distinct orange-brown stripes extending obliquely upward




Figure 2. Distribution of the pigfish.

and backward on back and sides, those below being nearly horizontal; head with bronze spots; fins yellow-bronze with dusky margins.

REASON FOR INCLUSION IN THE SERIES

Pigfish are common inhabitants of warm gulf waters (Reid 1954). They are frequently taken by sport anglers, especially in Florida waters, and are considered to be a good quality food fish (Darcy 1983); however, they have only limited economic importance (Joseph and Yerger 1956). Pigfish are often trapped and used for live bait (Carr 1976). They are also used as a source of food by other predatory species (Smith 1907).

LIFE HISTORY

Spawning

Pigfish mature by their second year of life (Taylor 1916; Hildebrand and Cable 1930). Reid (1954) concluded that spawning in gulf waters near Cedar Key, Florida, was probably in spring, since small young-of-the-year (23 to 28 mm SL) first appeared in May. He also noted a bimodal length-frequency curve for that month, suggesting that pigfish in this area of the gulf may have two breeding peaks or "growth spurts." Adult females taken during July from the Cedar Key area showed some signs of maturing gonads; however, they were far from ripe. After comparing average monthly lengths and seasonality of fish in other areas, Grimes and Mountain (1971) concluded that pigfish apparently spawn in about March in gulf waters near Crystal River, Florida. Gunter (1945) found ripe males in Texas gulf waters in March and April and suggested that the fish probably spawn there before June.

Other studies from the Gulf of Mexico also indicate that spawning

occurs in the late winter or spring: Tampa Bay, Florida, March-May (Springer and Woodburn 1960); Alligator Harbor, Florida, March (Joseph and Yerger 1956); eastern Gulf of Mexico, January-May (Darcy 1983); and Horn Island, Mississippi, March-April (Franks 1970). Hoese (1965) reported that pigfish larvae were found off Port Aransas from late February through June. Hastings (1972) noted that spawning may occur in open water prior to inshore migrations during March-April in St. Andrew Bay, Florida.

Pigfish spawn during March to June along the inside shores of Bogue and Shackleford Banks, North Carolina, and within the harbor and estuaries on the outer shores of these banks (Hildebrand and Cable 1930). Spawning apparently takes place during the early evening hours (Towers 1928; Hildebrand and Cable 1930).

Eggs

Johnson (1978) reported that pigfish eggs are buoyant, highly transparent, and spherical (0.7 to 0.8 mm in diameter). The eggs have a single oil globule (rarely two or three) that averages 0.16 mm in diameter. Pigfish eggs are easily confused with those of silver perch (*Bairdiella chrysoura*). This problem is magnified by the fact that the two species spawn at nearly the same time and in similar areas.

Larvae

Hildebrand and Cable (1930) provided descriptions of larval pigfish collected from North Carolina waters. Watson (1983) updated their descriptions using a series of larval pigfish specimens from lower Cape Fear Estuary, North Carolina, and from the gulf waters off Texas. Pigfish larvae can be separated from those of other haemulids (grunts) by the presence of 11 soft rays in the anal fin (Watson 1983). Darcy (1983)

cited a study by Houde et al. (1979) describing the distribution and abundance of pigfish larvae in the eastern gulf. They found one peak of abundance in late winter and spring, mainly inside a depth of 50 m.

Juveniles

Juvenile pigfish are fully scaled by 25 mm SL and achieve adult form by 70 mm SL (Figure 3; Hildebrand and Cable 1930). At 25 mm, they have a prominent dark midlateral band with an additional dark band from the nape to the base of the second dorsal fin. The mid-lateral bands often disappear at a length of approximately 40 mm SL, but the anterior position of the lower band may remain longer (Johnson 1978). Juvenile pigfish have yellow and green horizontal lines along their sides that are most prominent on the cheeks and opercles.

Grimes and Mountain (1971), working in an area of thermal effluent near Crystal River, Florida, first noted young-of-the-year pigfish in trawl samples taken during June. Juvenile pigfish (smallest, 12.5 mm SL) were first taken in April from Tampa Bay (Springer and Woodburn 1960) and St. Andrew Bay, Florida (Hastings 1972). Juveniles were most abundant during May in Cedar Key, Florida (Reid 1954). In Alligator Harbor, Florida, Joseph and Yerger (1956) reported that juveniles (41 mm SL) were present by June. Gunter (1945) and Hildebrand (1954) collected smallest pigfish in shallow bays along the Texas gulf coast.

Juvenile pigfish may associate with other species of fish. Darcy (1983) referenced a study by Wang and Raney (1971) from Charlotte Harbor, Florida, where young pigfish (16-25 mm SL) were found in mixed schools with small pinfish.




Figure 3. Juvenile pigfish, 38 mm TL (from Hildebrand and Cable 1930, Figure 37).

Adults

Adult pigfish are common in the northern and more saline coastal areas of the Gulf of Mexico (Springer and Woodburn 1960; Moe and Martin 1965; Hoese and Moore 1977). Reid (1954) reported that adult pigfish were found in Cedar Key, Florida, throughout the year (except January); however, pigfish were most abundant during summer months. Grimes and Mountain (1971) also noted that pigfish were more commonly taken during later summer and fall months in Crystal River, Florida. Along the coastal areas of north Florida, pigfish have been reported to be among the most abundant finfish collected (Joseph and Yerger 1956; Hastings et al. 1976; Ogren and Brusher 1977; Pristas and Trent 1978; Darcy 1983).

Adult pigfish also occur in the offshore and open-shelf areas of the Gulf of Mexico. Darcy (1983) noted a study by Cody et al. (1978) stating that pigfish occurred in 40%-43% of all trawl catches on the white shrimp grounds off the Texas gulf coast. Pigfish have also been taken from offshore reefs and platforms along the northern gulf (Hastings 1972; Hastings et al. 1976; Darcy 1983). Moe and Martin (1965) reported that pigfish may be more abundant in offshore waters than inshore in the southern regions of the Gulf of Mexico. Hildebrand (1954) noted that pigfish were abundant on the shell banks off Campeche, Mexico.

GROWTH CHARACTERISTICS

Pigfish reach a maximum length of 46 cm SL (Courtenay and Sahlman 1978) and a weight of 0.9 kg (2 lb) (Hildebrand and Cable 1930; Konchina 1977; Darcy 1983). Few pigfish older than 3 years old and very few age 4 fish have been taken along the Atlantic coast (Taylor 1916; Hildebrand and Cable 1930).

Information on growth of pigfish in gulf waters is limited to age 0 and age 1 fish (Table 1). Growth rates based on these data range from 7 mm SL/mo (Reid 1954) to 9.3 mm SL/mo from June to October. Growth during October to April slows to 3.1 mm SL/mo (Grimes and Mountain 1971) to 5.5 mm SL/mo (Reid 1954).

THE FISHERY

Commercial statistics are not specifically reported for pigfish, but have been combined with those for the other grunts by the Bureau of Commercial Fisheries and National Marine Fisheries Service. Pigfish are taken in seines, traps, trawls, and by handlines (Courtenay and Sahlman 1978). Most are marketed as live bait (especially in Texas); they are also considered a good quality food fish (Darcy 1983) but have little economic value. Marine recreational landing statistics for pigfish in the Gulf of Mexico for 1979 and 1981-85 are summarized in Table 2; most are taken along Florida (gulf and Atlantic) and the Texas gulf coast waters.

ECOLOGICAL ROLE

Food Habits

Feeding habits of pigfish vary with growth stage; they are primarily benthic carnivores as adults, possibly feeding nocturnally (Hastings et al. 1976), while young fish are planktivorous (Carr and Adams 1973; Darcy 1983). Reid (1954) found a shift in diet relative to increasing length for pigfish taken from Crystal River, Florida (Table 3). Small fish (25-50 mm TL) ate mostly copepods, while larger fish shifted to a diet of amphipods, shrimps, and other benthic organisms. Carr and Adams (1973), who also studied juvenile pigfish (16-80 mm SL) in the Crystal River, found two distinct feeding phases. Smaller pigfish (16-30 mm) were planktivorous (eating copepods, mysids, and

Table 1. Comparative monthly mean standard lengths (mm) for age 0 pigfish (from Grimes and Mountain 1971).

Study	Year collected	June	Aug.	Oct.	Dec.	Feb.	April
Springer and Woodburn 1960 (Tampa Bay)	1957	40.0	62.1				17.5
Reid 1954 (Cedar Keys) ^a	1951	49.0	66.0	77.0		84.0	110.0
Grimes 1971 ^b	1969						
Crystal River-Affected		44.0	66.0	70.0	79.0		98.0
Crystal River-Nonaffected		51.0	75.0				111.0
Grimes and Mountain 1971 ^b	1970						
Crystal River-Affected			78.9	94.0	88.2		102.9
Crystal River-Nonaffected		50.0	68.9	87.2	90.8		105.8

^aLengths estimated from graph.

^bStatistical comparison of annual growth (monthly mean standard length vs. time) of fish from thermally affected (warmed by the effluent from a steam electric station) vs. nonaffected areas revealed no significant difference.

postlarval shrimp), followed by a two-phase carnivorous stage in which benthic invertebrates were the major food items. The transition from planktivore to carnivore was gradual, beginning at about 26 mm SL, and was complete by 41-45 mm SL. Polychaetes were important in the diet of pigfish longer than 30 mm, but as fish grew larger than 55 mm, caridean and penaeid shrimp were consumed more frequently (Darcy 1983).

Predators

Pigfish are prey of Atlantic sharp-nose sharks, spotted seatrout, and weakfish (Radcliffe 1916; Hastings 1972; Darcy 1983). Other large piscivores, such as snapper and grouper, probably also prey on pigfish (Darcy 1983).

Parasites and Diseases

Pigfish have been reported to be parasitized by a monogenetic trematode that infests the gill filaments (Suydam 1971). Springer and Woodburn (1960) report that pigfish were killed by red tides in Tampa Bay, Florida.

ENVIRONMENTAL REQUIREMENTS

Temperature and Salinity

Reid (1954) found mean water temperature and salinity values of 25.2 °C and 25.1 ppt in Crystal River, Florida, when pigfish were most abundant. During December to April, when pigfish were taken less frequently, values of 15.9 °C and

Table 2. Summary of recreational fishing statistics for pigfish in the Gulf of Mexico.

Time period	Total U.S. catch (thousands of fish)	Percent of catch taken in Gulf of Mexico	Catch by Gulf States (thousands of fish)				
			FL	AL	MS	LA	TX
Jan-Dec ^a 1979	1,992	76.4	700	--	34	--	770
Mar-Dec ^b 1981	2,281	67.0	1,071	--	--	--	428
Jan-Dec ^b 1982	2,643	76.2	1,769	--	--	72	169
Jan-Dec ^c 1983	2,386	49.2	1,001	39	*	--	115
Jan-Dec ^c 1984	1,731	66.8	741	--	--	*	413
Jan-Dec ^d 1985	2,770	47.4	587	*	*	*	726

^aU.S. National Marine Fisheries Service (1980).

^bU.S. National Marine Fisheries Service (1985a).

^cU.S. National Marine Fisheries Service (1985b).

^dU.S. National Marine Fisheries Service (1986).

*means none reported.

--means less than 30,000 reported; however, the figure is included in removed column totals.

26.5 ppt were recorded. Roessler (1970) collected pigfish from Florida waters at temperatures of 19.5 to 30.6 °C and salinities of 17.2 to 44.1 ppt. A temperature range of 13.7 to 36 °C and a salinity range of 0 to 38 ppt was also provided by Roessler (1970) on the basis of published values for pigfish throughout the gulf. Springer and Woodburn (1960) reported a tolerance range of 19.1 to 35 ppt (mean = 28.9 ppt) and 17.5 to 32.5 °C for pigfish collected in Tampa Bay, Florida. In Barataria Bay, Louisiana, pigfish were taken between 6.2 and 24.3 ppt and between 17.3 and 30.0 °C (Dunham 1972).

Pigfish apparently avoid low-temperature water, migrating to deeper water during the winter (Hildebrand and Cable 1930; Gunter 1945; Reid 1954; Wang and Raney 1971; Grimes 1971; Hastings 1972; Ogren and Brusher 1977; Naughton and Saloman 1978; Darcy 1983). Hastings (1972) noted that pigfish were absent in collections made at water temperatures of 12 to 14 °C; however adults were abundant when waters of St. Andrew Bay, Florida, warmed to 16.5 to 31.0 °C. Moe and Martin (1965) also noted that pigfish were not collected when water temperatures in Pinellas County, Florida, dropped

Table 3. Food of pigfish (percentage frequency of occurrence) from Crystal River, Florida (from Reid 1954).

Food item	Size of fish (mm)		
	25-50	51-150	151-170
Copepods	83	38	--
Ostracods	50	--	--
Amphipods	--	54	10
Shrimps	17	56	40
Crabs	--	5	20
Mollusks	--	--	20
Polychaetes	17	8	60
Fishes	--	5	--
Insects	--	--	10

below 12.5 °C. Moore (1976) reported that pigfish were killed during a cold wave that caused water temperatures to drop to 4.5 °C (Darcy 1983).

Low salinity areas also appear to be avoided by pigfish. In Charlotte Harbor, Florida, Wang and Raney (1971) reported that pigfish were not taken in waters with salinity values less than 15 ppt. Gunter (1945) found all pigfish less than 50 mm TL and greater than 200 mm TL in Texas gulf waters in salinities greater than 25 ppt; no specimens were taken in less than 10 ppt. Reid (1954) noted a mass mortality of pigfish, as well as other species, after a hurricane caused salinities to drop from 23.5 ppt to 9.7 ppt over a 4-day period in Goose Cove, Florida. However, it was probably the rapid drop rather than the low salinity that caused the mortality (Darcy 1983).

Dissolved Oxygen

Schwartz et al. (1982) collected pigfish from waters with dissolved oxygen concentrations of 2.1 to 11.8 ppm in Cape Fear Estuary, North Carolina.

Substrate

Juvenile pigfish in Crystal River were found on shallow flats with considerable plant growth during spring and early summer (Reid 1954). As the summer and fall progressed, juvenile pigfish moved to deep flats and the edges of channels. Adults were taken from deeper flats and channels with sparse vegetation. Adult pigfish occurred most frequently over mud bottoms and occasionally over sandy, vegetated areas (Hildebrand and Schroeder 1928; Tabb and Manning 1961), hard substrates such as reefs and jetties (Hastings 1972), and offshore platforms (Hastings et al. 1976).

LITERATURE CITED

- Breder, C. M. Jr. 1948. Field book of marine fishes of the Atlantic coast from Labrador to Texas. G. P. Putnam's Sons, New York. 332 pp.
- Briggs, J. C. 1958. A list of Florida fishes and their distribution. Bull. Fla. State Mus. Biol. Ser. 2(8):223-318.
- Carr, W. E. S. 1976. Chemoreception and feeding behavior in the pigfish, Orthopristis chrysopterus: characterization and identification of stimulatory substances in a shrimp extract. Comp. Biochem. Physiol. 55(A):153-157.
- Carr, W. E. S., and C. A. Adams. 1973. Food habits of juvenile marine fishes occupying seagrass beds in the estuarine zone near Crystal River, Florida. Trans. Am. Fish. Soc. 102:511-540.
- Cody, T. J., K. W. Rice, and C. E. Bryan. 1978. Commercial fish and penaeid shrimp studies northwestern Gulf of Mexico. Pt. 5. Abundance and distribution of fauna on the white shrimp, Penaeus setiferus (Linnaeus), grounds off the central Texas coast. Coastal Fish. Branch, Tex. Parks Wildl. Dep., P.L.88-309, Proj. 2-276-R. 39 pp.
- Courtenay, W. R., Jr., and H. F. Sahlman. 1978. Pomadasyidae. Vol. 4, Unpaginated in W. Fischer, ed. FAO species identification sheets for fishery purposes, western central Atlantic (Fishing area 31). Food and Agriculture Organization of the United Nations, Rome.
- Darcy, G. H. 1983. Synopsis of biological data on the pigfish, Orthopristis chrysoptera. (Pisces: Haemulidae). FAO Fish. Synop. No. 134. 23 pp.
- Dunham, F. 1972. A study of commercially important estuarine-dependent industrial fishes. La. Wildl. Fish. Comm., Tech. Bull 4. 63 pp.
- Franks, J. S. 1970. An investigation of the fish population within the inland waters of Horn Island, Mississippi, a barrier island in the northern Gulf of Mexico. Gulf Res. Rep. 3:3-104.
- Grimes, C. B. 1971. Thermal addition studies of the Crystal River steam electric station. Fla. Dep. Nat. Resour. Mar. Res. Lab. Prof. Pap. Ser. 11. 53 pp.
- Grimes, C. B., and J. A. Mountain. 1971. Effects of thermal effluent upon marine fishes near the Crystal River steam electric station. Fla. Dep. Nat. Resour. Mar. Res. Lab. Prof. Pap. Ser. 17. 64 pp.
- Gunter, G. 1945. Studies on marine fishes of Texas. Publ. Inst. Mar. Sci. Univ. Tex. 1:1-90.
- Hastings, R. W. 1972. The origin and seasonality of the fish fauna on a new jetty in the northeastern Gulf of Mexico. Ph.D. Thesis. Florida State University, Tallahassee. 555 pp.
- Hastings, R. W., L. H. Ogren, and M. T. Mabry. 1976. Observations

- on the fish fauna associated with offshore platforms in the north-eastern Gulf of Mexico. U.S. Natl. Mar. Fish. Serv. Fish. Bull. 74: 387-402.
- Hildebrand, H. H. 1954. A study of the fauna of the brown shrimp (*Penaeus aztecus* Ives) grounds in the western Gulf. Publ. Inst. Mar. Sci. Univ. Tex. 3:233-366.
- Hildebrand, S. F., and L. E. Cable. 1930. Development and life history of fourteen teleostean fishes at Beaufort, N.C. U.S. Bur. Fish. Bull. 46:383-488.
- Hildebrand, S. F., and W. C. Schroeder. 1928. Fishes of Chesapeake Bay. U.S. Bur. Fish. Bull. 43, Pt. 1:1-388.
- Hoese, H. D. 1965. Spawning of marine fishes in Port Aransas, Texas, area as determined by the distribution of young and larvae. Ph.D. Dissertation, University of Texas, Austin. 144 pp.
- Hoese, H. D. and R. H. Moore. 1977. Fishes of the Gulf of Mexico, Texas, Louisiana, and adjacent waters. Texas A&M University Press, Austin. 327 pp.
- Houde, E. D., J. C. Leak, C. E. Dowd, S. A. Berkeley, and W. J. Richards. 1979. Ichthyoplankton abundance and diversity in the eastern Gulf of Mexico. Report to the Bureau of Land Management, under Contract No. AA550-CT7-28, 546 pp.
- Joseph, E. B., and R. W. Yerger. 1956. The fishes of Alligator Harbor, Florida, with notes on their natural history. Pap. Oceanogr. Inst. Fla. State Univ. Stud. 22:111-156.
- Johnson, G. D. 1978. Development of fishes of the mid-Atlantic bight; an atlas of eggs, larval and juvenile stages. Vol. IV. U.S. Fish Wildl. Serv. Biol. Serv. Program FWS/OBS-78/12.
- Konchina, Y. V. 1977. Some data on the biology of grunts (Family Pomadasyidae). [In Russ.] Vopr. Ikhtiol. 17:621-633. (Transl. J. Ichthyol. 17:548-558).
- Moe, M. A., Jr., and G. T. Martin. 1965. Fishes taken in monthly trawl samples offshore of Pinellas County, Florida, with new additions to the fish fauna of the Tampa Bay area. Tulane Stud. Zool. 12:129-151.
- Moore, R. H. 1976. Observations on fishes killed by cold at Port Aransas, Texas, 11-12 January 1973. Southwest. Nat. 20:461-466.
- Naughton, S. P., and C. H. Saloman. 1978. Fishes of the nearshore zone of St. Andrew Bay, Florida, and adjacent coast. Northeast Gulf Sci. 2:43-55.
- Ogren, L.H., and H. A. Brusher. 1977. The distribution and abundance of fishes caught with a trawl in the St. Andrew Bay System, Florida. Northeast Gulf Sci. 1:83-105.
- Pristas, P. J., and L. Trent. 1978. Seasonal abundance, size and sex ratio of fishes caught with gill nets in St. Andrew Bay, Florida. Bull. Mar. Sci. 28:581-589.
- Radcliffe, L. 1916. The sharks and rays of Beaufort, North Carolina. U.S. Bur. Fish. Bull. 34:239-284.
- Reid, G. K. 1954. An ecological study of the Gulf of Mexico fishes in the vicinity of Cedar Key, Fla. Bull. Mar. Sci. Gulf Caribb. 4(1): 1-94.
- Roessler, M. A. 1970. Checklist of fishes in Buttonwood Canal, Everglades National Park, Florida, and observations on the seasonal

- occurrence and life histories of selected species. *Bull. Mar. Sci.* 20(4):860-893.
- Schwartz, F. J., W. T. Hogarth, and M. P. Weinstein. 1982. Marine and freshwater fishes of the Cape Fear Estuary, North Carolina, and their distribution in relation to environmental factors. *Brimleyana* 7:17-37.
- Smith, H. M. 1907. The fishes of North Carolina. *N.C. Geol. Econ. Surv.* 2:1-458.
- Springer, V. G., and K. D. Woodburn. 1960. An ecological study of the fishes of the Tampa Bay area. *Fla. Board Conserv. Mar. Lab., Prof. Pap. Ser. 1.* 104 pp.
- Suydam, E. L. 1971. The microecology of three species of monogenetic trematodes of fishes from the Beaufort-Cape Hatteras area. *Proc. Helminthol. Soc. Wash.* 38:240-246.
- Tabb, D. C., and R. B. Manning. 1961. A checklist of the flora and fauna of northern Florida Bay and adjacent brackish waters of the Florida mainland collected during the period July, 1957 through September, 1960. *Bull. Mar. Sci. Gulf Caribb.* 11: 552-649.
- Taylor, H. F. 1916. The structure and growth of the scales of the squeteague and the pigfish as indicative of life history. *U.S. Bur. Fish. Bull.* 34:285-330.
- Towers, I. L. 1928. Embryology of the pigfish. Pages 622-624 in E. J. G. Rehn, ed. *Progress in biological sciences.* U.S. Fish. Comm. Rep. 1927:App. VII. *U.S. Bur. Fish. Doc.* 1029:517-588.
- U.S. National Marine Fisheries Service. 1980. Marine recreational statistics survey, Atlantic and gulf coasts, 1979. *U.S. Natl. Mar. Fish. Serv. Curr. Fish. Stat.* 8063. 139 pp.
- U.S. National Marine Fisheries Service. 1985a. Marine recreational fishery statistics survey, Atlantic and gulf coasts, 1981-1982. *U.S. Natl. Mar. Fish. Serv. Curr. Fish. Stat.* 8324. 215 pp.
- U.S. National Marine Fisheries Service. 1985b. Marine recreational fishery statistics survey, Atlantic and gulf coasts, 1983-1984. *U.S. Natl. Mar. Fish. Serv. Curr. Fish. Stat.* 8326. 222 pp.
- U.S. National Marine Fisheries Service. 1986. Marine recreational fishery statistics survey, Atlantic and gulf coasts, 1985. *U.S. Natl. Mar. Fish. Serv. Curr. Fish. Stat.* 8327. 130 pp.
- Wang, J. C. S., and E. C. Raney. 1971. Distribution and fluctuations in the fish fauna of the Charlotte Harbor Estuary, Florida. *Charlotte Harbor Estuarine Study, Mote Marine Laboratory, Sarasota, Florida* 56 pp.
- Watson, W. 1983. Redescription of larvae of the pigfish, *Orthopristis chrysoptera* Linnaeus (Pisces, Haemulidae). *U.S. Fish Wildl. Serv. Fish. Bull.* 81:847-854.

AD-A183070

REPORT DOCUMENTATION PAGE		1. REPORT NO. Biological Report 82(11.71)*		2.		3. Recipient's Accession No.	
4. Title and Subtitle Species Profiles: Life Histories and Environmental Requirements of Coastal Fishes and Invertebrates (Gulf of Mexico)--Pigfish						5. Report Date March 1987	
7. Author(s) F.C. Sutter and T.D. McIlwain						8. Performing Organization Report No.	
9. Performing Organization Name and Address						10. Project/Task/Work Unit No.	
12. Sponsoring Organization Name and Address National Wetlands Research Center U.S. Army Corps of Engineers Fish and Wildlife Service Waterways Experiment Station U.S. Dept. of the Interior P.O. Box 631 Washington, DC 20240 Vicksburg, MS 39180						11. Contract(G) or Grant(G) No. (C) (G)	
						13. Type of Report & Period Covered	
15. Supplementary Notes *U.S. Army Corps of Engineers Report No. TR EL-82-4						14.	
16. Abstract (Limit: 200 words) Species profiles are literature summaries of the taxonomy, morphology, range, life history, and environmental requirements of coastal aquatic species. They are designed to assist in environmental impact assessment. Pigfish are common inhabitants of warm gulf waters. Pigfish spawn during late winter and spring in open gulf waters. Young fish first appear in nearshore or estuarine environments in April to June. Juvenile pigfish may school with other species of fish (e.g. pinfish). Adult pigfish are found in higher salinity, coastal areas; they are also found in offshore, open shelf waters, as well as offshore reefs and platforms. Recreational fishermen frequently catch pigfish, especially in Florida gulf waters. Pigfish have little economic value; however, they are used for live bait and are considered to be a good quality food fish. Young pigfish are planktivorous, becoming carnivorous as they grow in length (first polychaetes, then shrimps). Pigfish have been taken in water temperatures ranging from 13.7 to 36°C, and in salinities of 0 to 38 ppt; however, they have been found in warmer water with salinities greater than about 15 ppt. Pigfish are found over vegetated sandy substrates, muddy bottoms, or hard substrates, such as reefs, jetties, and offshore platforms. Pigfish are prey of spotted seatrout and weakfish.							
17. Document Analysis a. Descriptors Estuaries Temperature Nearshore gulf water Salinity Growth Dissolved oxygen Life cycles Fisheries Food habits Fishes b. Identifiers/Open-Ended Terms Pigfish <u>Orthopristis chrysoptera</u> c. CDSAT Field/Group							
18. Availability Statement Unlimited availability				19. Security Class (This Report) Unclassified		21. No. of Pages vi + 11	
				20. Security Class (This Page) Unclassified		22. Price	

END

9-87

Dtic