

AD-A183 492

THE APPLICATION OF STATISTICAL TURBULENCE THEORY TO
CONVECTIVE INSTABILITIES(U) SRI INTERNATIONAL MENLO
PARK CA C L RINO 01 NOV 86 DNA-TR-86-402

1/1


UNCLASSIFIED

DNA001-86-C-0002

F/G 4/1

NL

							END						
							9-87						
							DTIC						



DTIC FILE COPY

(12)

DNA-TR-86-402

AD-A183 492

THE APPLICATION OF STATISTICAL TURBULENCE THEORY TO CONVECTIVE INSTABILITIES

C. L. Rino
SRI International
333 Ravenswood Avenue
Menlo Park, CA 94025-3434

1 November 1986

Technical Report

DTIC
SELECTED
AUG 05 1987
CS D

CONTRACT No. DNA 001-86-C-0002

Approved for public release;
distribution is unlimited.

THIS WORK WAS SPONSORED BY THE DEFENSE NUCLEAR AGENCY
UNDER RDT&E RMC CODE B3220867663 RW RA 00001 25904D.

Prepared for
Director
DEFENSE NUCLEAR AGENCY
Washington, DC 20305-1000

87 8 4 067

DISTRIBUTION LIST UPDATE

This mailer is provided to enable DNA to maintain current distribution lists for reports. We would appreciate your providing the requested information.

- ☐ Add the individual listed to your distribution list.
- ☐ Delete the cited organization/individual.
- ☐ Change of address.

NAME: _____

ORGANIZATION: _____

OLD ADDRESS

CURRENT ADDRESS

TELEPHONE NUMBER: () _____

SUBJECT AREA(s) OF INTEREST:

DNA OR OTHER GOVERNMENT CONTRACT NUMBER: _____

CERTIFICATION OF NEED-TO-KNOW BY GOVERNMENT SPONSOR (if other than DNA):

SPONSORING ORGANIZATION: _____

CONTRACTING OFFICER OR REPRESENTATIVE: _____

SIGNATURE: _____

CUT HERE AND RETURN



Director
Defense Nuclear Agency
ATTN: [REDACTED] TITL
Washington, DC 20305-1000

Director
Defense Nuclear Agency
ATTN: [REDACTED]
Washington, DC 20305-1000

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE


ADA183492

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY N/A since Unclassified			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE N/A since Unclassified					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) SRI Project 1700			5. MONITORING ORGANIZATION REPORT NUMBER(S) DNA-TR-86-402		
6a. NAME OF PERFORMING ORGANIZATION SRI International		6b. OFFICE SYMBOL (if applicable)		7a. NAME OF MONITORING ORGANIZATION Director Defense Nuclear Agency	
6c. ADDRESS (City, State, and ZIP Code) 333 Ravenswood Avenue Menlo Park, California 94025-3434				7b. ADDRESS (City, State, and ZIP Code) Washington, DC 20305-1000	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (if applicable) RAAE/Wittwer		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER DNA 001-86-C-0002	
8c. ADDRESS (City, State, and ZIP Code)				10. SOURCE OF FUNDING NUMBERS	
				PROGRAM ELEMENT NO. 62715H	PROJECT NO. RW
11. TITLE (Include Security Classification) THE APPLICATION OF STATISTICAL TURBULENCE THEORY TO CONVECTIVE INSTABILITIES					
12. PERSONAL AUTHOR(S) Rino, Charles L.					
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM 850313 TO 860930		14. DATE OF REPORT (Year, Month, Day) 861101	
15. PAGE COUNT 32					
16. SUPPLEMENTARY NOTATION This work was sponsored by the Defense Nuclear Agency under RDT&E RMC Code B3220867663 RW RA 00001 25904D.					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Turbulent Plasmas		
17	02	1	Convective Plasma Instabilities		
20	09		Irregularity Power Spectral Density		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Statistical turbulence theories have been used effectively to characterize spectral density functions in fluid turbulence, and in turbulent plasmas, magnetic fields, and passive scalars in turbulent fluids. In this report, we review the applications to convective instabilities of the turbulence theories developed originally and independently by Kraichnan and Kadomtsev, but generalized for ionospheric applications by Sudan. These instabilities are believed to be the dominant structuring mechanism in equatorial spread F, artificial barium clouds, and in high-altitude nuclear plumes at late times. We have emphasized the ramifications of the cross-field anisotropy unique to convective instabilities, which have not been treated previously. We show that the limiting form of the one-dimensional spectral density functions measured perpendicular to the $E \times B$ drift direction is proportional to k^{-1} where k is the wave number.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT ☐ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT ☐ DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL Sandra E. Young			22b. TELEPHONE (Include Area Code) (202) 325-7042		22c. OFFICE SYMBOL DNA/CSTI

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE




UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

Conversion factors for U.S. Customary to metric (SI) units of measurement

angstrom	1.000 000 X E -10	meters (m)
atmosphere (normal)	1.013 25 X E +2	kilo pascal (kPa)
bar	1.000 000 X E +2	kilo pascal (kPa)
barn	1.000 000 X E -28	meter ² (m ²)
British thermal unit (thermochemical)	1.054 350 X E +3	joule (J)
calorie (thermochemical)	4.184 000	joule (J)
cal (thermochemical)/cm ²	4.184 000 X E -2	mega joule/m ² (MJ/m ²)
curie	3.700 000 X E +1	giga becquerel (GBq)
degree (angle)	1.745 329 X E -2	radian (rad)
degree Fahrenheit	$t_F = (t_C + 459.67)/1.8$	degree kelvin (K)
electron volt	1.602 19 X E -19	joule (J)
erg	1.000 000 X E -7	joule (J)
erg/second	1.000 000 X E -7	watt (W)
foot	3.048 000 X E -1	meter (m)
foot-pound-force	1.355 818	joule (J)
gallon (U.S. liquid)	3.785 412 X E -3	meter ³ (m ³)
inch	2.540 000 X E -2	meter (m)
jerk	1.000 000 X E +9	joule (J)
joule/kilogram (J/kg) (radiation dose absorbed)	1.000 000	Gray (Gy)
kilotons	4.183	terajoules
kup (1000 lbf)	4.448 222 X E +3	newton (N)
kup/inch ² (ksi)	6.894 757 X E +3	kilo pascal (kPa)
ktap	1.000 000 X E +2	newton-second/m ² (N-s/m ²)
micron	1.000 000 X E -6	meter (m)
mil	2.540 000 X E -5	meter (m)
mile (international)	1.609 344 X E +3	meter (m)
ounce	2.834 952 X E -2	kilogram (kg)
pound-force (lbs avoirdupois)	4.448 222	newton (N)
pound-force inch	1.129 848 X E -1	newton-meter (N-m)
pound-force/inch	1.751 268 X E +2	newton/meter (N/m)
pound-force/foot ²	4.788 026 X E -2	kilo pascal (kPa)
pound-force/inch ² (psi)	6.894 757	kilo pascal (kPa)
pound-mass (lbm avoirdupois)	4.535 924 X E -1	kilogram (kg)
pound-mass-foot ² (moment of inertia)	4.214 011 X E -2	kilogram-meter ² (kg-m ²)
pound-mass/foot ³	1.601 846 X E +1	kilogram/meter ³ (kg/m ³)
rad (radiation dose absorbed)	1.000 000 X E -2	*Gray (Gy)
roentgen	2.579 760 X E -4	coulomb/kilogram (C/kg)
shake	1.000 000 X E -8	second (s)
slug	1.459 390 X E +1	kilogram (kg)
torr (mm Hg, 0°C)	1.333 22 X E -1	kilo pascal (kPa)

- The becquerel (Bq) is the SI unit of radioactivity; 1 Bq = 1 event/s
- The Gray (Gy) is the SI unit of absorbed radiation.



A-1

TABLE OF CONTENTS

Section	Page
CONVERSION TABLE	111
1 INTRODUCTION	1
2 THE DIA APPROXIMATION	8
3 DETERMINATION OF SDF CHARACTERISTICS	12
4 DISCUSSION	15
5 LIST OF REFERENCES	18

SECTION 1

INTRODUCTION

The simplest mathematical model that contains the essential physics of a convectively unstable plasma is given by the coupled, nonlinear differential equations

$$\frac{\partial N}{\partial t} + (\hat{b} \times \nabla \phi / B) \cdot \nabla N = K \nabla^2 N \quad , \quad (1)$$

and

$$\nabla(N \nabla \phi) = D \cdot \nabla N \quad , \quad (2)$$

where $N(\rho, t)$ is the electron density in a plane normal to the magnetic field. The vector D/B is the the cross-field drift velocity caused by the combined forces of gravity, neutral winds, and electric fields. Thus, D can be thought of as an effective electric field. The potential, $\phi(\rho, t)$, which is established by Eq. (2) from the instantaneous electron density configuration, modifies the effective electric field in such a way that, as time evolves, small-scale structures are drawn out of the trailing edge of high-density regions and are thereby enhanced. The structure is ultimately removed by cross-field diffusion at a rate determined by the diffusion coefficient K .

The physics of this two-dimensional model and its potential limitations are discussed in detail in Perkins et al. [1973], Ossakow [1981], and Zalesak et al. [1985]. Here, we shall discuss the applications of statistical turbulence theory to determine the quasi-steady-state spectral density function (SDF) that characterizes the late-time evolution of plasma structure implied by Eqs. (1) and (2). As yet, no viable theory has emerged that can explain the measured kilometer-scale

spectral characteristics of naturally occurring equatorial F-region ionospheric irregularities that severely affect satellite radio signals [Aarons, 1982]. Yet, it is generally believed that the development and evolution of these irregularities can be modeled by using straightforward modifications of Eqs. (1) and (2) [Zalesak et al., 1982]. Our motivation in undertaking this study was to determine if formal turbulence theory can be modified to explain the results of recent data analyses.

Ott and Farley [1974] first pointed out that Eq. (1) is mathematically identical to the two-dimensional Navier-Stokes (NS) equation when it is written in terms of vorticity,

$$\mathbf{w} = \nabla \times \mathbf{v} \quad , \quad (3)$$

where $\mathbf{v}$ is the incompressible fluid velocity ($\nabla \cdot \mathbf{v} = 0$). If $\mathbf{v}$ is confined to a plane, it is easily shown that the momentum balance of inertial, pressure, and viscous forces can be written as

$$\frac{\partial \mathbf{w}}{\partial t} + \mathbf{v} \cdot \nabla \mathbf{w} = \nu \nabla^2 \mathbf{w} \quad , \quad (4)$$

where ν is the kinematic viscosity. Thus the momentum balance for two-dimensional fluids when expressed in terms of scalar vorticity is identical to the continuity equation for electron density; however, to complete the formal parallel we must identify the electron drift velocity

$$\mathbf{v}_e = \hat{\mathbf{a}}_z \times \nabla \phi / B \quad (5)$$

with the fluid velocity $\mathbf{v}$.

Because $\mathbf{v}$ is divergence free, it too can be derived from a potential function. If we let $\mathbf{v} = \hat{\mathbf{a}}_z \times \nabla \phi$, it follows from Eq (3) that

$$\nabla^2 \phi = -w \quad . \quad (6)$$

Thus, for inertially driven, two-dimensional turbulence, the potential function that generates the velocity structure is related to the scalar vorticity by Poisson's equation. Because of the profound differences between Eqs. (2) and (6), theories that address the turbulent structure of w cannot be applied directly to the convective plasma system described by Eqs. (1) and (2). Indeed, even if one were to accept ordinary turbulence as the source of the structure in the velocity field, the electron density must act as a passive scalar to map the inertially driven velocity structure [Kelley and Ott, 1978]. Ideally, one would proceed from a general plasma theory that accommodates both collisional and inertial forces [Kintner and Seyler, 1985]; however, the collisional limit is the most difficult to treat, and it appears to have some unique properties.

Consider that in Kolmogoroff's theory of fully developed turbulence, locally unstable eddies spontaneously break up into smaller eddies until they are removed by viscous dissipation. In sharp contrast, convectively unstable density irregularities evidently do not spontaneously break up or bifurcate. Rather, early in the nonlinear phase of the structure development, fingerlike striations develop with a comparatively small-size distribution. Evidently the striations continue to elongate without further significant change in their individual shapes [McDonald et al., 1981; Zalesak, et al., 1985]. This process has been referred to as "freezing." Clearly, the structure dynamics are vastly different in the collisional and inertial models, and we should expect these differences to manifest themselves in a viable theory.

Formal plasma turbulence theories have been developed by Kraichnan and Kadomtsev. Both theories use the Direct Interaction Approximation, which was developed by Kraichnan [1958, 1959]; his turbulence theory is usually referred to as the DIA. Kadomtsev's method, which was developed more intuitively, is usually referred to as the Weak Coupling Approximation (WCA) [Kadomtsev, 1965]. Sudan and Pfirsch [1984] have shown that the two methods ultimately give rise to formally identical mathematical models; however, the WCA form suggests a somewhat simpler

procedure for deriving spectral characteristics. Following Sudan and Pfirsch's approach, we shall start with the DIA because it proceeds directly from Eq. (1). We shall then recast the equations in the WCA form and follow Sudan's prescription for deriving the form of one-dimensional spectral-density function. The main difference is that we have attempted to relax the assumption of isotropy, which is critical in all the applications of the method to date [Sudan and Keskinen, 1979; Sudan, 1983; Sudan and Keskinen, 1984]. This is achieved by approximating the anisotropy, which manifests itself in the "frozen" striation distribution, by a one-dimensional transfer of scalar variance in the Fourier domain. This provides an alternate means of obtaining a one-dimensional conservation equation in place of the angle-averaged form that is normally used.

The first step is to transform Eq. (1) into the spatial Fourier domain. We assume that $N(\rho, t)$ and $\phi(\rho, t)$ admit spectral decompositions of the form

$$\begin{pmatrix} N(\rho, t) \\ \phi(\rho, t) \end{pmatrix} = \sum \begin{pmatrix} \hat{N}(\mathbf{k}, t) \\ \hat{\phi}(\mathbf{k}, t) \end{pmatrix} \exp \{i\mathbf{k} \cdot \rho\} \quad (7)$$

where

$$\mathbf{k} = (n\Delta K_x, m\Delta K_y) \quad (8)$$

for all integer values of n and m . Thus, N and ϕ are periodic on the lattice bounded by $L_x = 2\pi/\Delta K_x$ and $L_y = 2\pi/\Delta K_y$. As discussed by Kraichnan [1975], this imposes no limitations of practical significance. Also, because N and ϕ are real, $\hat{N}$ and $\hat{\phi}$ have the hermitian property

$$\begin{pmatrix} \hat{N}(\mathbf{k}) \\ \hat{\phi}(\mathbf{k}) \end{pmatrix} = \begin{pmatrix} \hat{N}^*(-\mathbf{k}) \\ \hat{\phi}^*(-\mathbf{k}) \end{pmatrix} \quad (9)$$

Substituting Eq. (7) into Eq. (1) and performing some standard Fourier manipulations gives the spectral domain form of Eq. (1), namely.

$$\frac{\partial \hat{N}(k)}{\partial t} + k^2 K \hat{N}(k) = \sum^{\Delta} (\hat{b} \times k' \times k''/B) \hat{\phi}(k') \hat{N}(k'') \quad . \quad (10)$$

The symbol Δ means that only those modes that satisfy the triangle equality

$$k = k' + k'' \quad (11)$$

contribute to the summation. We could have included the delta function $\delta(k - k' - k'')$ in the summation, but the alternate notation is more compact. In the interest of notational efficiency, we have also suppressed the independent time variable t .

Next, we multiply Eq. (10) by $\hat{N}^*(k)$ to obtain

$$\frac{\partial |\hat{N}(k)|^2}{\partial t} + 2Kk^2 |\hat{N}(k)|^2 = E(k) \quad , \quad (12)$$

where

$$E(k) = 2 \sum^{\Delta} (\hat{b} \cdot k' \times k''/B) \hat{\phi}(k') \hat{N}(k'') \quad . \quad (13)$$

Equations (12) and (13) characterize the temporal evolution of the spectral intensity for a single realization of the process. The nonlinear mixing term $E(k)$ has the important property

$$\sum E(k) = 0 \quad , \quad (14)$$

which does not depend on how $\hat{\phi}(k)$ is related to $\hat{N}(k)$. Thus, in Eqs. (1) and (4) the nonlinear interaction of the Fourier modes does not change the scalar variance

$$\Omega = \sum |\hat{N}(\mathbf{k})|^2 \quad (15)$$

or enstrophy (the name given to the variance of scalar vorticity) in the NS system. This conservation property is the basis for developing a statistical turbulence theory.

A formal statement of the conservation of scalar variance can be obtained from Eqs. (12) and (14) as

$$\frac{\partial \Omega}{\partial t} + 2K \sum k^2 |\hat{N}(\mathbf{k})|^2 = 0 \quad (16)$$

Thus, the structure evolution is such that the only change in scalar variance is through diffusion. Turbulence theory attempts to characterize the average spectral characteristics in the quasi steady state where the tendency for structure to form is essentially balanced by the structure removal process. In fluid turbulence, energy must be supplied continuously to sustain this steady state. In collisional plasma structuring, a reservoir of structure at low spatial frequencies can sustain structure growth. Thus, in our application of the DIA, we do not include a forcing function.

We envision many realizations of the process with small, randomly irregular changes in the initial conditions, but with similar structure reservoirs at small spatial wave numbers to sustain the growth of small-scale structure at late times. We expect the ensemble average of the spectral intensities from each realization, formally

$$\Phi(\mathbf{k}) = \langle |\hat{N}(\mathbf{k})|^2 \rangle \quad (17)$$

to yield an identifiable SDF, although it is well known that Eqs. (1) and (2) will not support a true steady state. If we attempt to evaluate Eq. (17) by using Eqs. (12) and (13), we encounter the third-order moment $\langle \hat{\phi}(\mathbf{k}') \hat{N}(\mathbf{k}'') \hat{N}^*(\mathbf{k}) \rangle$. Iterating the procedure will always introduce a moment of higher order. Thus, irrespective of the mathematical subtleties associated with statistical stationarity, the formal moment

equations are indeterminate. Direct interaction is a closure approximation in which the third-order moments are evaluated in terms of fourth-order moments containing the temporal response function of Eq. (1) to small changes in ϕ . The fourth-order moments are evaluated by using the well-known relation for Gaussian processes, although the Gaussian assumption itself is not critical to the development.

SECTION 2

THE DIA APPROXIMATION

The Direct Interaction approximation is discussed in considerable detail in Leslie [1973]. For our purposes here, we have simply followed the step-by-step procedure he presents in his Chapter 4.3. The computational details are omitted. As a preliminary, we first consider the response of Eq. (10) to a small perturbation in $\hat{\phi}$. To terms that are first order in $\delta\hat{\phi}$ and $\delta\hat{N}$, we have

$$\delta\hat{N}(\mathbf{k};t) = \sum_{\mathbf{k}'}^{\Delta} (\hat{\mathbf{b}} \cdot \mathbf{k}' \times \mathbf{k}''/B) \int_{-\infty}^t G(\mathbf{k};t,t') \delta\hat{\phi}(\mathbf{k};t') \hat{N}(\mathbf{k}'';t') dt' \quad , \quad (18)$$

where $G(\mathbf{k};t,t')$ is the solution to

$$\left[\frac{\partial}{\partial t} + Kk^2 \right] G(\mathbf{k};t,t') - \sum_{\mathbf{k}'}^{\Delta} (\hat{\mathbf{b}} \cdot \mathbf{k}' \times \mathbf{k}''/B) \hat{\phi}(\mathbf{k}';t) G(\mathbf{k}'';t,t') = \delta(t-t') \quad . \quad (19)$$

To apply the DIA, we first expand $\hat{N}$, $\hat{\phi}$, and $\hat{G}$ in series of the form

$$\hat{N} = \hat{N}^{(0)} + \epsilon \hat{N}^{(1)} \quad , \quad (20)$$

where ϵ is the order of the nonlinear terms in Eqs. (10) and (19). By substituting these expansions into Eqs. (10) and (19) and equating terms of equal order, we obtain equations for the first-order terms $\hat{N}^{(1)}$ and $G^{(1)}$ in terms of the zeroth-order terms. Consistent with this approximation, we let

$$\hat{\phi}(\mathbf{k}) = W(\mathbf{k}) \hat{N}(\mathbf{k}) \quad , \quad (21)$$

where $W(k) = W^*(-k)$ to preserve the hermitian property of $\hat{\phi}(k)$. This is tantamount to linearizing the potential equation, Eq. (2), but within the context of the DIA, this is a local approximation that preserves the essential nonlinearity in the problem.

The lowest order term in the expansion of $\hat{N}$ is assumed a Gaussian variable, and the corresponding term in the expansion of G is assumed exact. After considerable algebraic manipulations, we obtain the following equations for the time-dependent SDF:

$$\begin{aligned} \frac{1}{2} \frac{\partial S_N(k; t, t')}{\partial t} + Kk^2 S_N(k, t, t') &= \sum^A (\hat{b} \cdot k' \times k''/B)^2 f(k', k'') \\ &\times \left[\int_{-\infty}^{t'} G(k, t', t'') S_N(k'; t, t'') S_N(k''; t, t'') dt'' \right. \\ &\left. - \int_{-\infty}^t G(k', t, t'') S_N(k; t, t'') S_N(k''; t', t'') dt'' \right] \quad (22) \end{aligned}$$

where

$$f(k', k'') = \frac{1}{2} |W(k') - W(k'')|^2, \quad (23)$$

$$\begin{aligned} \left[\frac{\partial}{\partial t} + Kk^2 \right] G(k; t, t') &= - \sum^A (\hat{b} \cdot k' \times k''/B)^2 f(k', k'') \\ &\times \int_{t'}^t G(k''; t, t'') S_N(k'; t, t'') G(k; t', t'') dt'' + \delta(t - t') \quad (24) \end{aligned}$$

and

$$S_N(k; t, t') = \langle N(k; t) N^*(k; t') \rangle. \quad (25)$$

As the DIA equations are written here, they accommodate both the NS and convective plasma equations. For NS, it follows from Eq. (6) that

$$W(k) = -k^{-2} \quad , \quad (26)$$

which is exact. For a convectively unstable plasma, it follows from Eq. (2), essentially by dropping the nonlinear term, that

$$W(k) = -1 \frac{k \cdot D}{k^2} \quad . \quad (27)$$

Because Eq. (26) is isotropic, the functions in Eqs. (22) and (24) depend only on the magnitude of k and the equations can be reduced to a one-dimensional form by integrating over all directions. When Eq. (27) applies, however, the basic equations admit no obvious simplification.

Thus, we transform the complete DIA equations to their more compact WCA form. We first assume that the time dependences are functions only of the corresponding time differences, whereby Eqs. (22), (24), and (25) can be transformed to the temporal frequency domain. After some straightforward manipulations, we obtain the equivalent WCA equations

$$\begin{aligned} (\omega - iKk^2) S_N(k; \omega) = & - \sum^{\Delta} (\hat{b} \cdot k' \times k''/B)^2 f(k'; k'') \\ & \times \left\{ \frac{\int_{-\infty}^{\infty} S_N(k'; \omega') S_N(k''; \omega - \omega') d\omega'}{[\omega - iKk^2 + \Gamma(k; \omega)]^*} \right. \\ & \left. - S_N(k; \omega) \int_{-\infty}^{\infty} \frac{S_N(k''; \omega - \omega') d\omega'}{[\omega' - iKk'^2 + \Gamma(k'; \omega')] } \right\} \end{aligned} \quad (28)$$

and

$$\begin{aligned} \Gamma(k; \omega) = & \sum^{\Delta} (\hat{b} \cdot k' \times k''/B)^2 f(k'; k'') \\ & \times \int_{-\infty}^{\infty} \frac{S_N(k''; \omega - \omega') d\omega'}{[\omega' - iKk'^2 + \Gamma(k'; \omega')] } \quad . \end{aligned} \quad (29)$$

The new function $\Gamma(k;w)$ is related to $G(k;w)$ by the equation

$$G(k;w) = \frac{2\pi i}{[w - iKk^2 + \Gamma(k;w)]} . \quad (30)$$

Eq. (28) can be written the equivalent form

$$|w + iKk^2 + \Gamma(k;w)|^2 S_N(k;w) = \sum^{\Delta} (\hat{b} \cdot k' \times k'' / B)^2 f(k', k'') \\ \times \int_{-\infty}^{\infty} S_N(k'; w') S_N(k''; w - w') dw' , \quad (31)$$

which isolates the nonlinear damping function $\Gamma(k;w)$. The WCA equations are usually presented in this form [Eqs. (18) and (19) in Sudan and Keskinen, (1979)].

In the general forms presented here, neither the DIA nor the WCA equations can be solved, but we can establish an important property of Eqs. (22) and (28). Working from Eq. (28), we define the argument of the summation as $R(k, k', k'')$, which has the property

$$R(k, k', k'') + R(k', k, k'') + R(k, k'', k') = 0 . \quad (32)$$

It follows from Eq. (32) that the summation of the right-hand side of Eq. (28) over all k , allowing for the triangle equality constraint, is zero. Thus, the conservation property of the basic equations is preserved in the DIA or WCA equations for the spectral density function itself. This property is used to determine the spectral characteristics.

SECTION 3

DETERMINATION OF SDF CHARACTERISTICS

Sudan and Keskinen [1979] have argued that $S_N(k;w)$ will be highly peaked about $w = 0$ with a characteristic width that depends on $\Gamma(k;w)$. They have postulated a functional relation of the form

$$S(k;w) = \phi(k)g(w/\Gamma_k) \quad , \quad (33)$$

where Γ_k is the direction averaged value of $\Gamma(k;w)$. For the moment, let us consider only the fact that we have effectively assumed that the time or temporal frequency dependence of $S_N(k;w)$ is a multiplicative function. With this assumption alone, we can rewrite Eq. (28) as

$$\begin{aligned} \frac{\partial \phi(k)}{\partial t} + Kk^2 \phi(k) &= 2\theta(k) \sum_{\Delta} (\hat{b} \times k' \times k''/B)^2 f(k';k'') \phi(k') \phi(k'') \\ &- \phi(k) \Gamma(k) \end{aligned} \quad (34)$$

where

$$\theta(k) = 2\pi i \iint_{-\infty}^{\infty} \frac{g(w'/\Gamma_k)g(w-w')/\Gamma_k}{[w + iKk^2 + \Gamma(k;w)]^*} \frac{dw}{2\pi} \frac{dw'}{2\pi} \quad . \quad (35)$$

and $\Gamma(k)$ denotes the value of $-i\Gamma(k,w)$ integrated over all w . Both $\theta(k)$ and $\Gamma(k)$ are purely real.

At this point, the usual approach is to assume that $\phi(k)$ is independent of the direction of k , as would be strictly true when Eq. (26) holds. Indeed, this same property would then apply to $\Gamma(k)$ and $\theta(k)$. The remaining term in Eq. (34) can be evaluated and a conservation

relation formulated in terms of the magnitude k . One then characterizes the transfer of scalar variance from large to small wave numbers. It happens, however, that the second term in Eq. (34) dominates this process, and a differential equation of the form

$$\frac{\partial}{\partial t} \int_0^k \phi(k') dk' + \int_{\alpha k}^k \Gamma(k') \phi(k') dk' = \int_0^k k'^2 K \phi(k') dk' \quad , \quad (36)$$

where α is a small positive number results. The lower limit of the nonlinear damping term is nonzero because arbitrarily large structures are essentially background inhomogeneities.

In the stationary state, the time derivative is replaced by a constant. If $K = 0$, it follows that $\Gamma(k)\phi(k)$ must be a function, say $Y(k)$, with the property

$$Y(k) = \alpha Y(\alpha k) \quad , \quad (37)$$

so that the integral over αk to k is zero. The solution to Eq. (37) is proportional to k^{-1} . To determine the form of $\phi(k)$, we must establish a relation between $\Gamma(k)$ and $\phi(k)$. Sudan and Keskinen [1979] have derived an approximate form by using the Gaussian form

$$g(w/\Gamma_k) = [1/(2\pi\Gamma_k^2)]^{1/2} \exp\{-w^2/\Gamma_k^2\} \quad (38)$$

for $g(w/\Gamma_k)$. With appropriate assumptions, they were able to derive the approximate relation (for two dimensions)

$$\Gamma_k = D_r k^2 I_k^{1/2} \quad , \quad (39)$$

where I_k denotes the angle averaged value of $\phi_N(k)$. It follows that I_k is proportional to k^{-2} . This would seem to imply that in the "cascade" range, the one-dimensional spectral density function is proportional to k^{-1} , which seems to be close to the measured value. As we have noted,

however, this approach is inconsistent with the anisotropy of the collisional instability.

We suggest applying the argument in a different way. Because the problem is approximately one dimensional, with most of the small-scale structure characterized by Fourier modes along D, it seems reasonable to replace the angle averaged quantities by integrals over the direction orthogonal to D, which we take to lie along the x axis. We obtain a relation identical to Eq. (36) with $\phi(k)$ replaced by $\theta(k_y)$. It follows that in the cascade range,

$$\theta(k_y) \Gamma(k_y) = k_y^{-1} \quad (40)$$

and

$$\Gamma(k_y) = k_0 v k_y^{1/2} \theta^{1/2}(k_y) \quad (41)$$

from which it follows that $\theta(k_y) = k_y^{-1}$; however, because of the one-dimensional approximation, this result should be regarded as a limiting form.

SECTION 4

DISCUSSION

We have shown that a consistent application of statistical turbulence theory to convective plasma instabilities must accommodate the anisotropy of the structure, which is unique to the collisional limit. To get some handle on how this anisotropy might affect the theory, we have followed the general prescription suggested by Sudan, but attempted to relax the isotropy assumption by going to the opposite extreme of a one-dimensional transfer of scalar variance. For well-developed structures transverse to the dominant convection direction, the limiting form of the one-dimensional spectral density function is k^{-1} . Along the convection direction, the one-dimensional spectrum should be "shape-dominated" and, therefore, k^{-2} at long wavelengths, giving way to a much steeper power law where diffusion is effective. Indeed, in the diffusion range the spectrum should be isotropic, at least if classical cross-field diffusion is the dominant mechanism for structure removal.

Ott and Farley [1974] first pointed out that a k^{-1} one-dimensional spectral density function for density structure is the form that results from the purely dimensional arguments of turbulence theory; however, they questioned the existence of a truly inertiallike subrange because it implied no dependence on the initial gradient; moreover, the available data then seemed to imply a k^{-2} form. In our development, we have not addressed the question of the constants or other factors that determine the level of the spectrum in addition to its shape. The dependence on a length scale comes from the α in Eq. (36). For fluid turbulence, α is related to the Kolmogoroff constant, but the theory as it is applied here does not determine its value nor does it identify an actual physical scale that α can be identified with, although an outer scale or maximum size for transfer of scalar variance is reasonable.

We also note that Batchelor [1959], in treating the structure of passive scalar additives in turbulent fluids, showed that in the presence of a linear velocity shear the average one-dimensional spectrum for a conserved scalar governed by Eq. (1) is also k^{-1} . In fluid turbulence, this comes about when the viscous cutoff for velocity turbulence occurs at a larger scale than the diffusive cutoff for the scalar additive. This is referred to as the viscous convective subrange [Tennekes and Lumely, 1972]. The velocity shear is a consequence of the turbulent dissipation mechanism. To apply Batchelor's theory to convective instabilities, the presence of the velocity shears must be established independently. The direct application of the turbulence theory is more satisfying.

There is yet another turbulence theory argument that gives rise to limiting spectral forms that have power-law slopes less than two. These are discussed in Kintner and Seyler [1985], and they involve a dual cascade, which can occur in two dimensional fluids. Indeed, the fact that Eq. (4) conserves enstrophy introduces a second conserved quantity because energy is conserved as well. Kraichnan [1967] showed that under these conditions two types of cascade can occur. At scale sizes above a stirring scale where the structure is initiated, there is an upward cascade of enstrophy and scaling arguments give a k^{-1} form for the energy spectrum. Below the stirring scale, there is a downward cascade of energy and the conventional $k^{-5/3}$ spectrum results. To the extent that density is purely a passive scalar in this process and is related to velocity via Eqs. (21) and (27), its spectrum assumes the same two-component spectral form. The corresponding forms that apply when Eqs. (21) and (26), which are exact, hold are given in Table I of Kintner and Seyler, [1985]. The problem here is that the conditions under which Eq. (27) can be used give rise to anisotropy, in any case, would preclude the possibility of density behaving as a passive scalar as well. Indeed, Sudans's theory has yet to be generalized to accommodate more than a single conserved quantity. More important, the experimental

data, although showing two-component forms with fairly shallow low-frequency indices, have not shown the two-component form predicted by the dual cascade theory.

To summarize, the two-dimensional spectral density function for convective instabilities as modeled by Eqs. (1) and (2) is anisotropic at the longest wavelengths with a power-law index approaching -1 in the direction perpendicular to the convection direction and steepening to -2 or greater along the convection direction. Beyond the freezing scale where diffusion becomes effective, the spectrum would be more nearly isotropic. The ramifications of this model for propagation models are probably not severe because the slope change occurs only over a limited scale-size regime. As a means of testing this hypothesis, however, the propagation issues should be pursued.

SECTION 5

LIST OF REFERENCES

- Aarons, J., "Global Morphology of Ionospheric Scintillations," Proc. IEEE, Vol. 70, No. 4, pp. 360-378 (April 1982).
- Batchelor, G. K., "Small-Scale Variation of Convected Quantities Like Temperature in Turbulent Fluid--Part 1. General Discussion and the Case of Small Conductivity," J. Fluid Mech., Vol. 5, Part 1, pp. 113-133 (January 1959).
- Kadmotsev, B. B., Plasma Turbulence, Chap. III, (Academic, 1965).
- Kelley, M. C. and E. Ott, "Two-Dimensional Turbulence in Equatorial Spread F," J. Geophys. Res., Vol. 83, No. A9, pp. 4369-4372 (September 1978).
- Kintner, P. M. and C. E. Seyler, "The Status of Observations and Theory of High Latitude Ionospheric and Magnetospheric Plasma Turbulence," Space Science Reviews, pp. 91-129 (February 1985).
- Kraichnan, R. H., "Irreversible Statistical Mechanics of Incompressible Hydro-Magnetic Turbulence," Phys. Rev., Vol. 109, No. 5, pp. 1407-1422 (March 1958).
- Kraichnan, R. H., "The Structure of Isotropic Turbulence at Very High Reynolds Numbers," J. Fluid Mech., Vol. 5, Part 4, pp. 497-543 (May 1959).
- Kraichnan, R. H., "Inertial Ranges in Two-Dimensional Turbulence," The Physics of Fluids, Vol. 10, No. 7, pp. 1417-1423 (July 1967).
- Kraichnan, R. H., "Statistical Dynamics of Two-Dimensional Flow," J. Fluid Mech., Vol. 67, No. 1, pp. 155-175 (1975).
- Leslie, D. C., Developments in the Theory of Turbulence (Clarendon Press, 1973).
- McDonald, B. E., S. L. Ossakow, S. T. Zalesak, and N. J. Zubusky, "Scale Sizes and Lifetimes of F Region Striations as Determined by the Condition of Marginal Stability," J. Geophys. Res., pp. 5775 (1981).

- Ossakow, S. L., "Spread F Theories--A Review," J. Atmos. Terr. Phys., Vol. 43, No. 5/6, pp. 437-452 (May/June 1981).
- Ott, E. and D. T. Farley, "The k Spectrum of Ionospheric Irregularities," J. Geophys. Res., Vol. 83, No. A5, pp. 2066-2070 (June 1974).
- Perkins, F. W., N. J. Zabusky, and J. H. Doles III, "Deformation and Striation of Plasma Clouds in the Ionosphere, 1," J. Geophys. Res., Vol. 78, No. 4, pp. 697-710 (February 1973).
- Sudan, R. N. and D. Pfirsch, "On the Relation Between andirect Interaction Approximation' Theories of Turbulence," Lab. of Plasma Studies, LPS 296 (August 1984).
- Sudan, R. N., "Unified Theory of Type I and Type II Irregularities in the Equatorial Electrojet," J. Geophys. Res., Vol. 88, No. A6, pp. 4853-4860 (June 1983).
- Sudan, R. N. and M. J. Keskinen, "Unified Theory of the Power Spectrum of Intermediate Wavelength Ionospheric Electron Density Fluctuations," Amer. Geophys. Un., No. 4A8108, pp. 9840-9844 (1984).
- Sudan, R. N. and M. Keskinen, "Theory of Strongly Turbulent Two-Dimensional Convection of Low-Pressure Plasma, Phys. Fluids, Vol. 22, No. 12, pp. 2305-2314, 1977 (1979).
- Tennekes, H. and J. L. Lumley, A First Course in Turbulence, (MIT Press, Cambridge, Massachusetts, 1972)
- Zalesak, S. T., P. K. Chaturvedi, S. L. Ossakow, and J. A. Fedder, "Finite Temperature Effects on the Evolution of Ionospheric Barium Clouds in the Presence of a Conducting Background Ionosphere. 1. The Simplest Case: Incompressible Background Ionosphere, Equipotential Magnetic Field Lines, and an Altitude-Invariant Neutral Wind," J. Geophys. Res., Vol. 90, No. A5, pp. 4299-4310 (May 1985).
- Zalesak, S. T., S. L. Ossakow, and P. K. Chaturvedi, "Nonlinear Equatorial Spread F: The Effect of Neutral Winds and Background Pedersen Conductivity," J. Geophys. Res., Vol. 87, No. A1, pp. 151-166 (January 1982).

DISTRIBUTION LIST

DEPARTMENT OF DEFENSE

ASSISTANT SECRETARY OF DEFENSE
CMD, CONTROL, COMMUNICATIONS & INTEL
ATTN: DASD(I)

ASSISTANT TO THE SECRETARY OF DEFENSE
ATTN: EXECUTIVE ASSISTANT

DEF RSCH & ENGRG
ATTN: STRAT & SPACE SYS (OS)

DEFENSE ADVANCED RSCH PROJ AGENCY
ATTN: GSD R ALEWINE
ATTN: T TETHER

DEFENSE COMMUNICATIONS AGENCY
ATTN: A200 R CRAWFORD
ATTN: A320 P BIRD
ATTN: A730 G JONES

DEFENSE COMMUNICATIONS ENGINEER CENTER
ATTN: CODE R123 (TECH LIB)

DEFENSE INTELLIGENCE AGENCY
ATTN: DC-7B
ATTN: DIR
ATTN: DT-1B
ATTN: RTS-2B
ATTN: VP-TPO

DEFENSE NUCLEAR AGENCY
ATTN: NANF
ATTN: NAWF
ATTN: OPNA
3 CYS ATTN: RAAE
ATTN: RAAE K SCHWARTZ
ATTN: RAAE T WALSH
ATTN: RAAE
4 CYS ATTN: TITL

DEFENSE TECHNICAL INFORMATION CENTER
12 CYS ATTN: DD

FIELD COMMAND DEFENSE NUCLEAR AGENCY
ATTN: FCTT W SUMMA
ATTN: FCTXE

FIELD COMMAND DNA DET 2
LAWRENCE LIVERMORE NATIONAL LAB
ATTN: FC-1

JOINT CHIEFS OF STAFF
ATTN: C3S

JOINT DATA SYSTEM SUPPORT CTR
ATTN: C-312 R MASON

JOINT STRAT TGT PLANNING STAFF
ATTN: JK (ATTN: DNA REP)
ATTN: JKCS
ATTN: JLWT (THREAT ANALYSIS)
ATTN: JPEP
ATTN: JPPFD
ATTN: JPSS
ATTN: JPTM

NATIONAL SECURITY AGENCY
ATTN: B432 C GOEDEKE

STRATEGIC DEFENSE INITIATIVE ORGANIZATION
ATTN: KE
ATTN: SLKT
ATTN: SN
ATTN: SY

DEPARTMENT OF THE ARMY

ARMY LOGISTICS MANAGEMENT CTR
ATTN: DLSIE

ASSISTANT CHIEF OF STAFF FOR AUTO & COMM
ATTN: DAMO-C4 P KENNY

DEP CH OF STAFF FOR OPS & PLANS
ATTN: DAMO-RQC (C2 DIV)

HARRY DIAMOND LABORATORIES
2 CYS ATTN: SCHLD-NW-P
ATTN: SLCHD-NW-R R WILLIAMS (22000)

U S ARMY ATMOSPHERIC SCIENCES LAB
ATTN: SLCAS-AE-E

U S ARMY COMMUNICATIONS R&D COMMAND
ATTN: DRDCO-COM-RY W KESSELMAN

U S ARMY FOREIGN SCIENCE & TECH CTR
ATTN: DRXST-SD

U S ARMY MATERIAL COMMAND
ATTN: DRCLDC J BENDER

U S ARMY MISSILE COMMAND
ATTN: DRSMI-YSO J GAMBLE

U S ARMY NUCLEAR & CHEMICAL AGENCY
ATTN: LIBRARY

U S ARMY SATELLITE COMM AGENCY
ATTN: AMCPM-SC-3

DNA-TR-86-402 (DL CONTINUED)

U S ARMY STRATEGIC DEFENSE CMD
ATTN: DASD-H-SAV

U S ARMY STRATEGIC DEFENSE COMMAND
ATTN: ATC-O W DAVIES
ATTN: ATC-R W DICKSON

U S ARMY TRADOC SYS ANALYSIS ACTVY
ATTN: ATAA-PL
ATTN: ATAA-TCC F PAYAN JR
ATTN: ATAA-TDC

US ARMY WHITE SANDS MISSILE RANGE
ATTN: STEWS-TE-N K CUMMINGS

DEPARTMENT OF THE NAVY

COMMAND & CONTROL PROGRAMS
ATTN: OP 941

JOINT CRUISE MISSILES PROJECT OFC (PM-3)
ATTN: JCMG-707

NAVAL AIR SYSTEMS COMMAND
ATTN: PMA 271

NAVAL ELECTRONICS ENGRG ACTVY, PACIFIC
ATTN: CODE 250 D OBRYHIM

NAVAL INTELLIGENCE SUPPORT CTR
ATTN: NISC-50

NAVAL OCEAN SYSTEMS CENTER
ATTN: CODE 532
ATTN: CODE 54 J FERGUSON

NAVAL RESEARCH LABORATORY
ATTN: CODE 4180 J GOODMAN
ATTN: CODE 4700 S OSSAKOW
ATTN: CODL 4720 J DAVIS
ATTN: CODE 4732 B RIPIN
ATTN: CODE 4750 P RODRIGUEZ
ATTN: CODE 4780 J HUBA

NAVAL SURFACE WEAPONS CENTER
ATTN: CODE H-21

NAVAL UNDERWATER SYS CENTER
ATTN: CODE 3411, J KATAN

OFC OF THE DEPUTY CHIEF OF NAVAL OPS
ATTN: NOP 941D
ATTN: OP 654(STRAT EVAL & ANAL BR)
ATTN: OP 981N

SPACE & NAVAL WARFARE SYSTEMS CMD
ATTN: CODE 3101 T HUGHES
ATTN: CODE 501A
ATTN: PD 50TD
ATTN: PDE-110-X1 B KRUGER

ATTN: PDE-110-11021 G BRUNHART
ATTN: PME 106-4 S KEARNEY
ATTN: PME-106 F W DIEDERICH

STRATEGIC SYSTEMS PROGRAM OFFICE (PM-1)
ATTN: NSP-L63 (TECH LIB)
ATTN: NSP-2141
ATTN: NSP-2722

THEATER NUCLEAR WARFARE PROGRAM OFC
ATTN: PMS-42331F (D SMITH)

DEPARTMENT OF THE AIR FORCE

AIR FORCE CTR FOR STUDIES & ANALYSIS
ATTN: AFCSA/SAMI (R GRIFFIN)
ATTN: AFCSA/SASC

AIR FORCE GEOPHYSICS LABORATORY
ATTN: LID/J RAMUSSEN
ATTN: LIS J BUCHAU
ATTN: LS
ATTN: LS/R O'NIEL
ATTN: LSI/ H GARDINER
ATTN: LYD/K CHAMPION

AIR FORCE SATELLITE CTRL FACILITY
ATTN: WE

AIR FORCE SPACE DIVISION
ATTN: YA
ATTN: YG
ATTN: YK
2 CYS ATTN: YN

AIR FORCE SPACE TECHNOLOGY CENTER
ATTN: XP

AIR FORCE TECHNICAL APPLICATIONS CTR
ATTN: TN

AIR FORCE WEAPONS LABORATORY, NTAAB
ATTN: NTN
ATTN: SUL

AIR FORCE WRIGHT AERONAUTICAL LAB/AAAD
ATTN: W HUNT

AIR UNIVERSITY LIBRARY
ATTN: AUL-LSE

BALLISTIC MISSILE OFFICE/DAA
ATTN: ENSN

DEPUTY CHIEF OF STAFF/AF-RDQM
ATTN: AF/RDQI

DEPUTY CHIEF OF STAFF/AFRDS
ATTN: AFRDS (SPACE SYS & C3 DIR)

ELECTRONIC SYSTEMS DIVISION/SC

ATTN: SCS-1E
ATTN: SCS-2 LTC ALEXANDER

ROME AIR DEVELOPMENT CENTER, AFSC

ATTN: OCDS R SCHNEIBLE
ATTN: TSLD

SPACE COMMAND/DC

ATTN: XP

STRATEGIC AIR COMMAND/NRI-STINFO

ATTN: NRI/STINFO

STRATEGIC AIR COMMAND/XPFC

ATTN: XPFS

STRATEGIC AIR COMMAND/XPQ

ATTN: XPQ

DEPARTMENT OF ENERGY

DEPARTMENT OF ENERGY

ATTN: DP-233

EG&G, INC

ATTN: D WRIGHT

UNIVERSITY OF CALIFORNIA

LAWRENCE LIVERMORE NATIONAL LAB

ATTN: L-31 R HAGER
ATTN: L-53 TECH INFO DEPT. LIBRARY

LOS ALAMOS NATIONAL LABORATORY

ATTN: D SAPPENFIELD
ATTN: D SIMONS
ATTN: J WOLCOTT
ATTN: MS J ZINN
ATTN: R JEFFRIES
ATTN: R W WHITAKER
ATTN: T KUNKLE

SANDIA NATIONAL LABORATORIES

ATTN: D HARTLEY 8300

SANDIA NATIONAL LABORATORIES

ATTN: A D THORNBROUGH 400
ATTN: D DAHLGREN 6440
ATTN: ORG 1231 T P WRIGHT
ATTN: ORG 314 W D BROWN
ATTN: ORG 332 R C BACKSTROM
ATTN: SPACE PROJECT DIV
ATTN: TECH LIB 3141 (RPTS RECEIVING CLRK)

OTHER GOVERNMENT

CENTRAL INTELLIGENCE AGENCY

ATTN: OSWR/NED
ATTN: OSWR/SSD FOR K FEUERPFETL

U S DEPARTMENT OF COMMERCE

ATTN: A JEAN (UNCL ONLY)
ATTN: W UTLAUT

US DEPARTMENT OF STATE

ATTN: PM/TMP

DEPARTMENT OF DEFENSE CONTRACTORS

AEROSPACE CORP

ATTN: D OLSEN
ATTN: E RODRIGUEZ
ATTN: I GARFUNKEL
ATTN: J KLUCK
ATTN: J STRAUS
ATTN: K S CHO
ATTN: R SLAUGHTER
ATTN: T SALMI

AEROSPACE CORP

ATTN: S MEWATERS

ANALYTICAL SYSTEMS ENGINEERING CORP

ATTN: SECURITY

AUSTIN RESEARCH ASSOCIATES

ATTN: J THOMPSON

AUTOMETRIC, INC

ATTN: C LUCAS

BDM CORP

ATTN: A VITELLO
ATTN: L JACOBS

BERKELEY RSCH ASSOCIATES, INC

ATTN: C PRETTIE
ATTN: J WORKMAN
ATTN: S BRECHT

BOEING CO

ATTN: G HALL
ATTN: D CLAUSON

BR COMMUNICATIONS

ATTN: J MCLAUGHLIN

CALIFORNIA RESEARCH & TECHNOLOGY, INC

ATTN: M ROSENBLATT

CHARLES STARK DRAPER LAB, INC

ATTN: A TETESKI

COMMUNICATIONS SATELLITE CORP

ATTN: G HYDE

COMPUTER SCIENCES CORP

ATTN: F EISENBARTH

DNA-TR-86-402 (DL CONTINUED)

CORNELL UNIVERSITY

ATTN: D FARLEY JR
ATTN: M KELLY

ELECTROSPACE SYSTEMS, INC

ATTN: H LOGSTON
ATTN: P PHILLIPS

EOS TECHNOLOGIES, INC

ATTN: B GABBARD
ATTN: W LELEVIER

GENERAL ELECTRIC CO

ATTN: A STEINMAYER
ATTN: C ZIERDT

GEO CENTERS, INC

ATTN: E MARRAM

GTE GOVERNMENT SYSTEMS CORPORATION

ATTN: W I THOMPSON, III

HARRIS CORP

ATTN: E KNICK

HSS, INC

ATTN: D HANSEN

IBM CORP

ATTN: H ULANDER

INSTITUTE FOR DEFENSE ANALYSES

ATTN: E BAUER
ATTN: H WOLFHARD

JAYCOR

ATTN: J SPERLING

JOHNS HOPKINS UNIVERSITY

ATTN: C MENG
ATTN: J D PHILLIPS
ATTN: J NEWLAND
ATTN: K POTOCKI
ATTN: R STOKES
ATTN: T EVANS

KAMAN SCIENCES CORP

ATTN: E CONRAD

KAMAN TEMPO

ATTN: B GAMBILL
ATTN: DASIAC
ATTN: R RUTHERFORD
ATTN: W MCNAMARA

KAMAN TEMPO

ATTN: DASIAC

LOCKHEED MISSILES & SPACE CO, INC

ATTN: J KUMER
ATTN: R SEARS

LOCKHEED MISSILES & SPACE CO, INC
2 CYS ATTN: D CHURCHILL

M I T LINCOLN LAB

ATTN: D TOWLE
ATTN: I KUPIEC

MAXIM TECHNOLOGIES, INC

ATTN: J LEHMAN
ATTN: J MARSHALL
ATTN: J SO
ATTN: N CIANOS

MCDONNELL DOUGLAS CORP

ATTN: R HALPRIN

METEOR COMMUNICATIONS CORP

ATTN: R LEADER

MISSION RESEARCH CORP

ATTN: B R MILNER
ATTN: C LAUER
ATTN: C RINO
ATTN: D ARCHER
ATTN: D KNEPP
ATTN: F FAJEN
ATTN: F GUIGLIANO
ATTN: G MCCARTOR
ATTN: K COSNER
ATTN: R BIGONI
ATTN: R BOGUSCH
ATTN: R DANA
ATTN: R HENDRICK
ATTN: R KILB
ATTN: R MORGANSTERN
ATTN: S GUTSCHE
ATTN: TECH LIBRARY

MITRE CORPORATION

ATTN: A KYMMEL
ATTN: C CALLAHAN
ATTN: D RAMPTON, PH.D
ATTN: DR D RAMPTON
ATTN: M R DRESP
ATTN: R DRESP

MITRE CORPORATION

ATTN: J WHEELER
ATTN: M HORROCKS
ATTN: R C PESCI
ATTN: W FOSTER

NORTHWEST RESEARCH ASSOC, INC

ATTN: E FREMOUW

DNA-TR-86-402 (DL CONTINUED)

PACIFIC-SIERRA RESEARCH CORP

ATTN: E FIELD JR
ATTN: F THOMAS
ATTN: H BRODE, CHAIRMAN SAGE

PHOTOMETRICS, INC

ATTN: I L KOFSKY

PHYSICAL RESEARCH INC

ATTN: H FRITZ
ATTN: J JORDANO

PHYSICAL RESEARCH, INC

ATTN: R DELIBERIS
ATTN: T STEPHENS

PHYSICAL RESEARCH, INC

ATTN: J DEVORE
ATTN: J THOMPSON
ATTN: W SCHLUETER

R & D ASSOCIATES

ATTN: B LAMB
ATTN: C GREIFINGER
ATTN: F GILMORE
ATTN: G HOYT
ATTN: H ORY
ATTN: M GANTSWEG
ATTN: M GROVER
ATTN: R TURCO
ATTN: W KARZAS

R & D ASSOCIATES

ATTN: B WEBSTER

R & D ASSOCIATES

ATTN: G GANONG

RAND CORP

ATTN: C CRAIN
ATTN: E BEDROZIAN
ATTN: P DAVIS

RAND CORP

ATTN: B BENNETT

ROCKWELL INTERNATIONAL CORP

ATTN: S QUILICI

SCIENCE APPLICATIONS INTL CORP

ATTN: C SMITH
ATTN: D HAMLIN
ATTN: E STRAKER
ATTN: L LINSON

SCIENCE APPLICATIONS INTL CORP

ATTN: R LEADABRAND

SCIENCE APPLICATIONS INTL CORP

ATTN: J COCKAYNE

SCIENCE APPLICATIONS INTL CORP

ATTN: D TELAGE
ATTN: M CROSS

SRI INTERNATIONAL

ATTN: D MCDANIEL (UNCL ONLY)
ATTN: W CHESNUT
ATTN: W JAYE

STEWART RADIANCE LABORATORY

ATTN: R HUPPI

TECHNOLOGY INTERNATIONAL CORP

ATTN: W BOQUIST

TELECOMMUNICATION SCIENCE ASSOCIATES

ATTN: R BUCKNER

TOYON RESEARCH CORP

ATTN: J GARBARINO
ATTN: J ISE

TRW ELECTRONICS & DEFENSE SECTOR

ATTN: DR D GRYBOS
ATTN: R PLEBUCH

UTAH STATE UNIVERSITY

ATTN: A STEED
ATTN: D BURT
ATTN: K BAKER, DIR ATMOS & SPACE SCI
ATTN: L JENSEN, ELEC ENG DEPT

VISIDYNE, INC

ATTN: J CARPENTER

END

9-87

DTIC