

REPORT NO. T27-87

FREEZING TIMES, REWARMING TIMES, AND LOWEST TEMPERATURES IN EXPERIMENTAL FROSTBITE OF HANFORD MINIATURE SWINE

AD-A185 051

**U S ARMY RESEARCH INSTITUTE
OF
ENVIRONMENTAL MEDICINE
Natick, Massachusetts**

JULY 1987

**DTIC
ELECTE
SEP 24 1987**



Approved for public release; distribution unlimited.

**UNITED STATES ARMY
MEDICAL RESEARCH & DEVELOPMENT COMMAND**

The findings in this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

DISPOSITION INSTRUCTIONS

Destroy this report when no longer needed.

Do not return to the originator.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION U.S. Army Research Institute of Environmental Medicine		6b. OFFICE SYMBOL (if applicable) SGRD-UE-CR	7a. NAME OF MONITORING ORGANIZATION U.S. Army Research Institute of Environmental Medicine		
6c. ADDRESS (City, State, and ZIP Code) Kansas Street Natick, MA 01760-5007			7b. ADDRESS (City, State, and ZIP Code) Kansas Street Natick, MA 01760-5007		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c. ADDRESS (City, State, and ZIP Code)			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO. 62777A	PROJECT NO. 3E1627 77A879	TASK NO. BB
			WORK UNIT ACCESSION NO. 121 DAOC633		
11. TITLE (Include Security Classification) Freezing Times, Rewarming Times, and Lowest Temperatures in Experimental Frostbite of Hanford Miniature Swine					
12. PERSONAL AUTHOR(S) Polly Schoning, D.V.M., Ph.D., Murray P. Hamlet, D.V.M.					
13a. TYPE OF REPORT Technical Report		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) 1978, 14 July 1987	
15. PAGE COUNT					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Frostbite was produced in the skin of five Hanford Miniature Swine exposed to chilled air (-75°C) for 1, 3, 5, 10, or 20 minutes. A hypodermic probe with a copper-constantan thermocouple was inserted into the skin to monitor the temperature. Average freezing time - the time to reach 0°C, or lower - was approximately 2 minutes. The average lowest temperature for 1 minute freezes ranged from -8.3 to -9.2°C and for the 3 minute freezes ranged from -13.0 to -18.2°C. Excluding pig 5, the 5, 10, and 20 minute freezes had a smaller range, from -18.9 to -24.3. Average rewarming times were shortest for 1 minute freezes and longest for 20 minute freezes. The ranges within freezes were minimal: 2.8-3.8, 4.0-4.9, 5.0-6.3, 6.2-7.2, and 7.6-8.9 minutes (+.5 min.) for the 1,3,5,10, and 20 minute freezes respectively.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT ☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT. ☐ DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL POLLY SCHONING, D.V.M., Ph.D.			22b. TELEPHONE (Include Area Code) (617) 651-4869		22c. OFFICE SYMBOL SGRD-UE-CR

DISCLAIMER STATEMENT

The views, opinions, and/or findings contained in this report are those of the authors and should not be construed as official Department of the Army position, policy, or decision, unless so designated by other official documentation. In conducting the research described in this report, the investigators adhered to the "Guide for Laboratory Animal Facilities and Care," as promulgated by the Committee on the Guide for Laboratory Animal Facilities and Care of the Institute of Laboratory Animal Resources, National Academy of Sciences, National Research Council.

Accession For	
NTIS CRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	



Technical Report

T27-87

FREEZING TIMES, REWARMING TIMES, AND LOWEST TEMPERATURES
IN EXPERIMENTAL FROSTBITE OF HANFORD MINIATURE SWINE

by

Polly Schoning DVM, PhD and Murray Hamlet DVM

US Army Research Institute of Environmental Medicine
Natick, Massachusetts 01760

TABLE OF CONTENTS

	<u>Page</u>
List of Tables	iv
Forward	v
Acknowledgements	vi
Abstract	vii
Introduction	1
Materials and Methods	1
Results	2
Discussion	2
Conclusions	3
References	7
Tables	5-11

LIST OF TABLES

	<u>Page</u>
Table 1 - Length of time in minutes ($\pm .5$ min.) for skin of Hanford Miniature Swine to reach freezing when exposed to -75°C air. Each number represents a frostbite lesions.	5
Table 2 - Lowest temperatures ($-^{\circ}\text{C}$) for skin of Hanford Miniature Swine exposed to -75°C air for 1, 3, 5, 10, or 20 minutes after -0°C or lower was recorded.	7
Table 3 - Length of time in minutes ($\pm .5$ min.) for skin of Hanford Miniature Swine frozen for 1, 3, 5, 10, or 20 minutes to rewarm to 25°C when exposed to 25°C air.	9
Table 4 - Freezing time averages in (minutes $\pm .5$ min) for skin of Hanford Miniature Swine to reach freezing when exposed to -75°C air.	11
Table 5 - Lowest temperature averages ($-^{\circ}\text{C}$) for 1, 3, 5, 10, or 20 minute freezes in 5 Hanford Miniature Swine exposed to -75°C air.	11
Table 6 - Rewarming time averages in (minutes) ($\pm .5$ min.) for 1, 3, 5, 10, and 20 minute freezes in 5 Hanford Miniature Swine exposed to -75°C air.	11

FORWARD

This experiment was done in 1986 as part of the U. S. Army Summer Faculty Research and Engineering Program administered by Battelle. In addition, Kansas State University has played a major role in providing support for continuation of the project. Kansas State has provided salary to the principal investigator, laboratory space, and clerical and photographic support.

ACKNOWLEDGEMENT

The authors thank Dr. George Silver, Ms. Audrey Armstrong, and Sgts. Rodrigo Mariano, Mark Sharp, and George Staruch for their technical assistance.

Abstract

Frostbite was produced in the skin of five Hanford Miniature Swine exposed to chilled air (-75°C) for 1, 3, 5, 10, or 20 minutes. A hypodermic probe with a copper-constantan thermocouple was inserted into the skin to monitor the temperature. Average freezing time - the time to reach 0°C , or lower - was approximately 2 minutes. The average lowest temperature for 1 minute freezes ranged from -8.3 to -9.2°C and for the 3 minute freezes ranged from -13.0 to -18.2°C . Excluding pig 5, the 5, 10, and 20 minute freezes had a smaller range, from -18.9 to -24.3 . Average rewarming times were shortest for 1 minute freezes and longest for 20 minute freezes. The ranges within freezes were minimal: 2.8-3.8, 4.0-4.9, 5.0-6.3, 6.2-7.2, and 7.6-8.9 minutes(± 0.5 min.) for the 1, 3, 5, 10, and 20 minute freezes respectively.

Introduction

Third degree frostbite has been consistently reproduced in Hanford Miniature Swine by exposing them to chilled air (-75°C).¹ Morphology and development of those changes have been described.^{2,3} In the same experiment, freezing times, lowest temperatures, and rewarming times were recorded.

This report documents and discusses the following:

- 1) length of time for skin to reach 0°C when exposed to -75°C air
- 2) lowest temperature skin reaches when frozen for 1, 3, 5, 10, or 20 minutes
- 3) length of time for skin frozen for 1, 3, 5, 10, or 20 minutes to rewarm to 25°C when exposed to 25°C air

Materials and Methods

Experimental pigs and procedures were described in detail previously.^{1,2} Briefly, frostbite lesions were produced by cold air (-75°C) delivered to 1-inch circular areas of the skin of five pigs. A hypodermic probe with a copper-constantan thermocouple was inserted into the skin, parallel to the surface for an approximate distance of 1.0 cm. Temperatures were monitored every 30 seconds. Once the thermocouple reached 0°C or lower, the cold air was continued for 1, 3, 5, 10, or 20 minutes. At the end of that time, the skin was rewarmed with warm air (25°C) until the computer recorded a skin temperature of 25°C .

For this report the following terms were used:

freezing time - time, to the nearest 30 seconds, for skin to reach 0°C when exposed to -75°C air.

lowest temperature - recorded during freezing skin with -75°C for 1, 3, 5, 10, or 20 minutes.

rewarming time - time, to the nearest 30 seconds, for skin frozen for 1, 3, 5, 10, or 20 minutes to rewarm to 25°C when exposed to 25°C air.

Results

Freezing times, lowest temperatures, and rewarming times for each of 40 lesions per pig (200 total) are listed in Tables 1 - 3.

Freezing time averages (Table 4) are for 40 lesions per pig. This table is small because the length of time for skin to reach freezing is unrelated to the duration it will be frozen.

Lowest temperature averages (Table 5) and rewarming time averages (Table 6) are for each duration of freeze, for each pig. Each average represents 8 lesions.

Discussion

From the time pig skin was exposed to -75°C chilled air, until it reached 0°C was approximately 2 minutes. Averages for pigs 1-4 had a narrow range of 1.9 to 2.2 minutes ($\pm .5$ min.). Pig 5 had a low average of 1.3 minutes ($\pm .5$ min.). The reason for this finding is not known. Possible explanations are that the thermocouple may have been inserted

more superficially, or the environmental or animal temperatures may have had some influence. Since the computer recorded the temperature only every 30 seconds, freezing may have occurred just after a computer recording enough times to make a difference.

As would be expected, lowest temperature averages became lower the longer the skin was frozen. One minute freezes ranged from -8.3 to -9.2°C and 3 minute freezes ranged from -13.0 to -18.2°C . The 5, 10, and 20 minute freezes had a smaller range from -18.9 to -24.3°C , excluding pig 5.

The average lowest temperatures for pig 5 were consistently lower than those for the other four pigs. This finding adds support to the hypothesis that the thermocouple was probably inserted more superficially in pig 5.

Rewarming times were shortest for skin frozen for 1 minute and longest for skin frozen for 20 minutes. The shortest average thaw was 2.8 minutes ($\pm .5$ min.) and the longest thaw was 8.9 minutes ($\pm .5$ min.). The ranges within freezes were minimal for all groups.

Comments

One of the biggest drawbacks of this project was the hypodermic probe, because it was inserted manually. Any change in its depth could result in a change in temperature recorded.

Although 5 pigs is a small sample, evaluation of 200 lesions from those pigs is acceptable as a baseline. This time and temperature study should add support to histopathologic studies of frostbite.

These two studies together are the controls necessary to evaluate topical ointments used to treat or prevent frostbite.

References

1. Schoning P, Hamlet MP: Experimental third degree frostbite in Hanford Miniature Swine. Technical Report T24-87, 1987.
2. Schoning P, Hamlet MP: Experimental frostbite in Hanford Miniature Swine. I. Epithelial changes. submitted for publication, 1987.
3. Schoning P, Hamlet MP: Experimental frostbite in Hanford Miniature Swine. II. Vascular changes. submitted for publication, 1987.
4. Sharp MW, Ahle NW, Mariano RA, Sawyer W: Development of a tissue freezing and rewarming devise that uses compressed air as a medium. Technical Report T13/87:1-9, 1987.

Table 1 - Length of time in minutes (± 0.5 min.) for skin of Hanford Miniature Swine to reach freezing when exposed to -75°C air. Each number represents a frostbite lesion.

pig 1	1.5	3	2	2	2.5
	3	2.5	3.5	2	2.5
	1.5	1	2	2.5	2
	4	2.5	2.5	1.5	2.5
	1	1	2	1.5	1
	1	1.5	1.5	1	1.5
	4	1.5	1.5	5.5	4.5
	1.5	1.5	1.5	5	1
pig 2	3	2	2	2	2
	3	3	2.5	1.5	2
	2	4	6.5	2.5	2
	1	1	1	.5	1
	2	2	1	1.5	1.5
	1	1	2.5	1.5	4
	2.5	1	.5	4	1.5
	1	1	.5	1.5	2.5
pig 3	1	2	2	1.5	2
	1.5	2.5	.5	2	4.5
	2.5	2	1.5	2.5	1.5
	1	3.5	.5	3	3
	2	1	4.5	2.5	8.5
	1.5	1.5	.5	2	1
	1.5	1.5	1	1.5	1
	.5	2	1.5	1.5	2
pig 4	1.5	1.5	5.5	5.5	1.5
	1	1	2	3	3
	1.5	6	1	3.5	1.5
	2.5	.5	2.5	2	4.5
	1.5	1	1.5	3.5	2.5
	1	1.5	.5	1	1.5
	1	1	1	.5	1
	1	.5	1	1.5	1.5
pig 5	2.5	1.5	1.5	1.5	3
	2	2.5	1	1.5	*5
	1	1	1	1	23
	.5	1	1.5	1.5	2
	1.5	1.5	1.5	1	1
	.5	1.5	1	1	1.5
	.5	1	1.5	1	2
	.5	1	1.5	.5	1

* This number was not included in calculating averages.

Table 2 - Lowest temperatures ($^{\circ}\text{C}$) for skin of Hanford Miniature Swine exposed to -75°C air for 1, 3, 5, 10, or 20 minutes after -0°C or lower was recorded.

		<u>duration of freeze (minutes)</u>				
		<u>1</u>	<u>3</u>	<u>5</u>	<u>10</u>	<u>20</u>
pig 1		12.9	13.4	20.2	20.7	22.9
		7.6	14.4	15.9	27.9	18.8
		9.3	19.3	23.6	21.1	24.2
		2.9	11.1	16.7	30.8	22.8
		15.9	26.4	17.5	24.2	26.7
		7.4	15.7	15.7	23.9	30.9
		2.8	16.2	20.1	8.2	18.5
		7.9	9.9	21.5	8.7	29.9
pig 2		4.3	15.9	14.2	23.6	20.4
		6.8	9.7	18.7	24.6	20.2
		5.6	8.1	6.5	18.9	25.9
		12.2	21.4	22.0	34.2	33.7
		8.3	12.7	27.2	23.8	22.7
		11.8	22.1	14.4	23.9	14.5
		4.3	18.7	31.3	8.3	22.8
		13.9	20.6	26.4	26.0	22.9
pig 3		9.8	15.6	21.6	26.3	25.5
		9.5	13.8	21.4	18.4	13.7
		8.4	11.9	18.1	20.4	22.7
		12.6	7.0	20.0	12.8	17.1
		5.5	23.3	7.6	16.6	8.7
		10.2	11.6	27.9	18.7	30.8
		5.8	11.4	20.8	22.5	26.2
		11.0	9.6	17.7	24.9	22.1
pig 4		10.2	10.1	18.8	8.8	27.9
		13.6	16.7	19.6	15.0	16.6
		9.3	4.6	27.7	10.4	22.9
		3.8	22.1	11.6	25.6	14.7
		6.2	17.9	21.8	13.0	20.2
		11.2	17.0	20.9	31.0	27.3
		6.9	14.4	22.1	29.8	25.4
		10.7	21.6	24.4	27.0	31.7
pig 5		7.7	22.2	23.0	28.9	20.6
		7.3	10.4	27.5	28.3	31.4
		7.5	15.9	29.8	30.9	20.3
		9.2	15.0	29.3	26.5	25.7
		8.5	15.6	22.9	23.4	26.6
		7.6	13.8	26.9	32.3	24.8
		9.6	28.8	20.0	27.8	27.9
		16.5	23.5	23.3	31.6	27.1

Table 3 - Length of time in minutes (± 0.5 min.) for skin of Hanford Miniature Swine frozen for 1, 3, 5, 10, or 20 minutes to rewarm to 25°C when exposed to 25°C air.

		<u>duration of freeze (minutes)</u>				
		<u>1</u>	<u>3</u>	<u>5</u>	<u>10</u>	<u>20</u>
pig 1	3	3	4.5	8	6.5	9.5
	3.5	3.5	4.5	7.5	7.5	8
	3.5	3.5	3.5	5	6.5	7
	6	6	5.5	7.5	7.5	10.5
	3	3	4	6.5	5.5	7
	3.5	3.5	3.5	6	6.5	9
	5	5	3.5	4	8	10.5
	3	3	4	5.5	9.5	10
pig 2	3.5	3.5	4.5	5	6	9.5
	3.5	3.5	4	5.5	5.5	7.5
	2.5	2.5	5	5.5	5	6
	2.5	2.5	3.5	5	5.5	8.5
	3	3	4	4.5	6	5.5
	2.5	2.5	4	6.5	9	10
	3.5	3.5	3.5	4.5	8	12
	2.5	2.5	3.5	3.5	10	12
pig 3	3	3	6	6	7	9.5
	2.5	2.5	5.5	4	4.5	5.5
	3.5	3.5	4.5	5	6.5	5.5
	3	3	7	5.5	10	12
	2.5	2.5	3.5	5.5	5	6
	3	3	4.5	5	7.5	8.5
	3	3	4	5	9	10
	2	2	4.5	6.5	9	12.5
pig 4	3	3	5	7.5	8	8
	2	2	4.5	6	8	6
	2.5	2.5	5	4	4	5
	4.5	4.5	4.5	6.5	9.5	12.5
	3	3	3.5	4.5	7	6.5
	3	3	4.5	6	6.5	9
	2.5	2.5	4.5	6	7	7
	3	3	4	5	10	16
pig 5	5	5	6.5	6.5	5.5	6
	4.5	4.5	4.5	5.5	5.5	6.5
	2.5	2.5	4	4.5	5.5	4
	3	3	4.5	6	7.5	11
	3.5	3.5	4.5	5	5.5	6.5
	1.5	1.5	5.5	5	7	7
	2.5	2.5	5	6	6.5	9.5
	2	2	4.5	5	6.5	10.5

Table 4 - Freezing time averages in (minutes $\pm .5\text{min}$) for skin of Hanford Miniature Swine to reach freezing when exposed to -75°C air.

	<u>minutes</u>
pig 1	2.2
pig 2	2.0
pig 3	2.0
pig 4	1.9
pig 5	1.3

Table 5 - Lowest temperature averages ($-^{\circ}\text{C}$) for 1, 3, 5, 10, or 20 minute freezes in 5 Hanford Miniature Swine exposed to -75°C air.

	<u>duration of freeze (minutes)</u>				
	<u>1</u>	<u>3</u>	<u>5</u>	<u>10</u>	<u>20</u>
pig 1	8.3	15.8	18.9	20.7	24.3
pig 2	8.4	16.2	20.1	22.9	22.9
pig 3	9.1	13.0	19.4	20.1	20.9
pig 4	9.0	15.6	20.9	20.1	23.3
pig 5	9.2	18.2	25.3	28.7	25.6

Table 6 - Rewarming time averages in (minutes) ($\pm .5\text{min.}$) for 1, 3, 5, 10, and 20 minute freezes in 5 Hanford Miniature Swine exposed to -75°C air.

	<u>duration of freeze (minutes)</u>				
	<u>1</u>	<u>3</u>	<u>5</u>	<u>10</u>	<u>20</u>
pig 1	3.8	4.1	6.3	7.2	8.9
pig 2	2.9	4.0	5.0	6.9	8.9
pig 3	2.8	4.9	5.3	7.3	8.7
pig 4	2.9	4.4	5.7	7.5	8.8
pig 5	3.1	4.9	5.4	6.2	7.6