

AD-A185 366


FREIDLIN-MENTZELL TYPE ESTIMATES AND THE LAW OF THE
ITERATED LOGARITHM FD. (U) NORTH CAROLINA UNIV AT
CHAPEL HILL CENTER FOR STOCHASTIC PROC. T MORI ET AL.
MAY 87 TR-184 AFOSR-TR-87-1106 F/G 12/3

1/1

UNCLASSIFIED

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

DTIC FILE COPY

AD-A185 366

①

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE


REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY NA			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; Distribution Unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE NA					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. 184			5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 87-1106		
6a. NAME OF PERFORMING ORGANIZATION University of North Carolina		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION AFOSR/NM		
6c. ADDRESS (City, State and ZIP Code) Statistics Dept. Phillips Hall 039-A Chapel Hill, NC 27514			7b. ADDRESS (City, State and ZIP Code) Building 410 Bolling AFB, DC 20332-6448		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR		8b. OFFICE SYMBOL (If applicable) NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F49620 85 C 0144		
8c. ADDRESS (City, State and ZIP Code) Building 410, Bolling AFB, DC 20332-6448			10. SOURCE OF FUNDING NOS.		
			PROGRAM ELEMENT NO 6.1102F	PROJECT NO. 2304	TASK NO. A5
			WORK UNIT NO.		
11. TITLE (Include Security Classification) Freidlin-Wentzell type estimates and the law of the iterated logarithm for a class of					
12. PERSONAL AUTHOR(S) stochastic processes related to -ymmetric statistics Mori, T. and Oodaira, H.					
13a. TYPE OF REPORT preprint		13b. TIME COVERED FROM 9/86 TO 9/87		14. DATE OF REPORT (Yr., Mo., Day) May 1987	
				15. PAGE COUNT 10	
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB. GR.	Keywords and phrases:		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Analogues of Freidlin and Wentzell's estimates for diffusion processes and the functional law of the iterated logarithm are obtained for a class of stochastic processes represented by multiple Wiener integrals with respect to two parameter Wiener processes, which arise as the limit processes of sequences of normalized symmetric statistics.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT ☐ DTIC USERS ☐			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL			22b. TELEPHONE NUMBER (Include Area Code)		22c. OFFICE SYMBOL

DTIC
ELECTE
OCT 07 1987
S E D

CENTER FOR STOCHASTIC PROCESSES

Department of Statistics
University of North Carolina
Chapel Hill, North Carolina



FREIDLIN-WENTZELL TYPE ESTIMATES AND THE LAW OF THE
ITERATED LOGARITHM FOR A CLASS OF STOCHASTIC
PROCESSES RELATED TO SYMMETRIC STATISTICS

by

Toshio Mori

and

Hiroshi Oodaira

Technical Report No. 184

May 1987

- 1 -

FREIDLIN-WENTZELL TYPE ESTIMATES AND THE LAW OF THE
ITERATED LOGARITHM FOR A CLASS OF STOCHASTIC
PROCESSES RELATED TO SYMMETRIC STATISTICS

Toshio Mori
Department of Mathematics
Yokohama City University

and

Hiroshi Oodaira*
Center for Stochastic Processes
Department of Statistics
University of North Carolina at Chapel Hill
and
Department of Applied Mathematics
Yokohama National University

Abstract

Analogues of Freidlin and Wentzell's estimates for diffusion processes and the functional law of the iterated logarithm are obtained for a class of stochastic processes represented by multiple Wiener integrals with respect to two parameter Wiener processes, which arise as the limit processes of sequences of normalized symmetric statistics. 

*Research supported by Air Force Office of Scientific Research Grant No.

F49620 85C 0144.



Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

1. Introduction and results. Let $h = h(u_1, \dots, u_m)$ be a square integrable symmetric function on $[0,1]^m$ and assume that h is canonical, i.e., it satisfies the condition

$$\int_0^1 h(u_1, u_2, \dots, u_m) du_1 = 0 \quad \text{for all } u_2, \dots, u_m \in [0,1].$$

Let $\{X_j\}$ be a sequence of independent identically distributed random variables uniformly distributed over $[0,1]$. Consider the following random sequence of normalized symmetric statistics

$$Y_n(t) = n^{-m/2} \sum_{1 \leq i_1 < \dots < i_m \leq [nt]} h(X_{i_1}, \dots, X_{i_m}), \quad 0 \leq t \leq 1,$$

in $D[0,1]$, the space of right continuous functions on $[0,1]$ having left limits with Skorohod's J_1 topology. A. Mandelbaum and M.S. Taqqu [3] showed that the random sequence $\{Y_n(t)\}$ converges weakly in $D[0,1]$ to the following process X :

$$X(t) = \int_{[0,1]^m} \dots \int_{[0,1]^m} h(u_1, \dots, u_m) 1_t(v_1) \dots 1_t(v_m) \cdot W(du_1, dv_1) \dots W(du_m, dv_m), \quad 0 \leq t \leq 1,$$

where the right hand side is an m -ple Wiener integral with respect to a two parameter Wiener process $\{W(u,v), 0 \leq u, v \leq 1\}$ and $1_t(\cdot)$ is the indicator function of $[0,t]$. X has continuous paths a.s. and note also that it can be written as

$$X(t) = \int_{[0,1]^m} h(u_1, \dots, u_m) W_t(du_1) \dots W_t(du_m), \quad 0 \leq t \leq 1,$$

with $W_t(u) = W(u,t)$.

The purpose of this note is firstly to prove certain large deviations results, i.e., asymptotic estimates of Freidlin-Wentzell type, for the above process X , and secondly to remark that the functional law of the iterated logarithm for X can be derived by the same arguments.

Let $C_N = C([0,1]; \mathbb{R}^N)$ be the space of $\mathbb{R}^N$ -valued continuous functions x on $[0,1]$ vanishing at the origin, with the norm $\|x\|_C = \sup_{0 \leq t \leq 1} |x(t)|$ and the metric $d(\cdot, \cdot)$, where $|\cdot|$ stands for the Euclidean norm in $\mathbb{R}^N$. Let $B = B(t) = (B_1(t), 1 \leq i \leq N)$.

$0 \leq t \leq 1$, be an N -dimensional standard Brownian motion with $B(0)=0$, and let H_N denote the reproducing kernel Hilbert space (RKHS) associated with B , i.e., the Hilbert space consisting of absolutely continuous function φ on $[0,1]$ such that $\varphi(0)=0$ and its derivative $\dot{\varphi}$ is square integrable. Its norm $||\varphi||_H$ is given by $||\varphi||_H = ||\dot{\varphi}||_2 = (\int_0^1 |\dot{\varphi}|^2)^{1/2}$. H_N is a subspace of C_N and the sets $K_r = \{\varphi \in H_N : ||\varphi||_H \leq r\}$, $r > 0$, are compact in C_N .

Define a mapping A from $L^2[0,1] \otimes H_1$ (the tensor product of $L^2[0,1]$ and RKHS H_1) to C_1 by

$$Af(t) = \int \dots \int_{[0,1]^m} h(u_1, \dots, u_m) f(u_1, t) \dots f(u_m, t) du_1 \dots du_m, \quad 0 \leq t \leq 1$$

for $f \in L^2[0,1] \otimes H_1$, and let G denote the class of functions

$$G = \{g = Af, f \in L^2[0,1] \otimes H\}.$$

Let $|||\cdot|||$ be the norm of $L^2[0,1] \otimes H_1$, which is given, e.g., by

$$|||f|||^2 = \int_0^1 \int_0^1 \left(\frac{\partial}{\partial t} f(u, t) \right)^2 du dt. \quad \text{Define}$$

$$D(g) = \inf \{ |||f||| : g = Af, f \in L^2[0,1] \otimes H_1 \} \quad \text{for } g \in G,$$

and

$$G_r = \{g = Af : |||f||| \leq r\}, \quad r > 0.$$

The main result of this note is the following

Theorem 1. (i) For any $g \in G$ and for any $\delta, \delta' > 0$, there is a number $\alpha_1 = \alpha_1(\delta, \delta')$,

$D(g)$ such that

$$P(||X/\alpha - g||_C < \delta) \geq \exp[-(\alpha^{2/m}/2)(D^2(g) + \delta')]$$

for all $\alpha \geq \alpha_1$, and

(ii) for any $\delta, \delta', r > 0$, there is a number $\alpha_2 = \alpha_2(\delta, \delta', r)$ such that

$$P(d(X/\alpha, G_r) > \delta) \leq \exp[-(\alpha^{2/m}/2)(r^2 - \delta')]$$

for all $\alpha \geq \alpha_2$.

In the proof of Theorem 1 we need the following Freidlin-Wentzell type

estimates for N-dimensional Brownian motion B.

Theorem A (i) For any $\varphi \in H_N$ and for any $\delta, \delta' > 0$, there is a number

$\alpha_1 = \alpha_1(\delta, \delta', \|\varphi\|_H)$ such that

$$P(\|B/\alpha - \varphi\|_C < \delta) \geq \exp[-(\alpha^2/2)(\|\varphi\|_H^2 + \delta')]$$

for all $\alpha \geq \alpha_1$ and

(ii) for any $\delta, \delta', r > 0$, there is a number $\alpha_2 = \alpha_2(\delta, \delta', r)$ such that

$$P(d(B/\alpha, K_r) > \delta) \leq \exp[-(\alpha^2/2)(r^2 - \delta')]$$

for all $\alpha \geq \alpha_2$.

Theorem A is a special case of general result on Gaussian processes, and we shall use the following easy consequences of Theorem A (see [2]).

Let F be a continuous mapping from C_N to C_1 , homogeneous with degree $p > 0$, i.e., satisfying the condition $F(c \cdot) = c^p F(\cdot)$ for any $c > 0$.

Theorem B (i) For any $\varphi \in H_N$ and for any $\delta, \delta' > 0$,

$$P(\|F(B)/\alpha - F(\varphi)\|_C < \delta) \geq \exp[-(\alpha^{2/p}/2)(\|\varphi\|_H^2 + \delta')]$$

for all sufficiently large α , and

(ii) for any $\delta, \delta', r > 0$,

$$P(d(F(B)/\alpha, F(K_r)) > \delta) \leq \exp[-(\alpha^{2/p}/2)(r^2 - \delta')]$$

for all sufficiently large α .

Theorem C

$$\lim_{\alpha \rightarrow \infty} (1/\alpha^{2/p}) \log P(\|F(B)\|_C > \alpha) = -b^2/2,$$

where

$$\begin{aligned} b^2 &= \inf\{\|\varphi\|_H^2: \|F(\varphi)\|_C > 1\} \\ &= \sup\{r^2: \sup(\|F(\varphi)\|_C: \varphi \in K_r) < 1\}. \end{aligned}$$

The arguments used in the proof of Theorem 1, combined with Strassen's law of the iterated logarithm for Brownian motion, yield also the following functional iterated logarithm law for the process X .

Theorem 2. Define the random sequence $\{Z_n\}$ in C_1 by

$$Z_n(t) = X(nt)/(2n \log_2 n)^{m/2}, \quad 0 \leq t \leq 1, \quad n \geq 3,$$

where $\log_2 = \log \log$. Then, with probability 1 $\{Z_n\}$ is relatively compact and the set of its limit points coincides with G_1 .

Remark. The above theorem is an improvement of a recent result of H. Dehling [1], in the sense that a moment condition on h is weakened. However, it should be noted that Dehling [1] proves more generally the functional law of the iterated logarithm for $\{Y_n(t)\}$. The methods of proofs are different.

2. Proof of Theorem 1

Let $\{e_i, i \geq 0\}$ be a complete orthonormal sequence (CONS) in $L^2[0,1]$ with $e_0=1$. Then $\{e_{i_1}(u_1) \dots e_{i_m}(u_m), i_1, \dots, i_m \geq 0\}$ is a CONS in $L^2([0,1]^m)$ and $h \in L^2([0,1]^m)$, symmetric and canonical, can be expanded as

$$h(u_1, \dots, u_m) = \sum_{1 \leq i_1, \dots, i_m} c_{i_1, \dots, i_m} e_{i_1}(u_1) \dots e_{i_m}(u_m), \quad c_{i_1, \dots, i_m} \in \mathbb{R}.$$

For $N \geq 1$ let

$$h_N(u_1, \dots, u_m) = \sum_{1 \leq i_1, \dots, i_m \leq N} c_{i_1, \dots, i_m} e_{i_1}(u_1) \dots e_{i_m}(u_m),$$

and define the process $X_N = \{X_N(t), 0 \leq t \leq 1\}$ by

$$X_N(t) = \int \dots \int_{[0,1]^m} h_N(u_1, \dots, u_m) W_t(du_1) \dots W_t(du_m).$$

Then we have

Lemma 1

$$P(\|X - X_N\|_C > z) \leq C \exp(-M_N z^{2/m})$$

for all sufficiently large $z > 0$, where C is a finite constant and the positive constant M_N can be made arbitrarily large by taking N large.

Proof. The lemma follows from a result of Plikusas [5] on multiple Wiener integrals and Lemmas 6.2 and 6.3 of [4]. Indeed it is enough to note that the exponential bound obtained by Plikusas holds also for multiple Wiener integrals with respect to two parameter Wiener processes and that, putting $Z_N = X - X_N$, we have

$$E|Z_N(t+s) - Z_N(t)|^2 \leq C_N \cdot s \quad \text{for } 0 \leq t < t+s \leq 1,$$

with a finite constant C_N and

$$\|E|Z_N(t)|^2\|_C = m! \|h - h_N\|_2^2,$$

where $\|\cdot\|_2$ is the norm of $L^2([0,1]^m)$. Now

$$X_N(t) = \sum_{1 \leq i_1, \dots, i_m \leq N} c_{i_1 \dots i_m} \int_{[0,1]^m} e_{i_1}(u_1) \dots e_{i_m}(u_m) W_t(du_1) \dots W_t(du_m)$$

and, since $\{e_i\}$ is orthonormal, each term of the right hand side can be written as a product of Hermite polynomials, i.e.,

$$\begin{aligned} & \int_{[0,1]^m} e_{i_1}(u_1) \dots e_{i_m}(u_m) W_t(du_1) \dots W_t(du_m) \\ &= H_{p_1} \left(\int_0^1 e_{q_1}(u) W_t(du) \right) \dots H_{p_r} \left(\int_0^1 e_{q_r}(u) W_t(du) \right), \end{aligned}$$

if there are $p_1 e_{q_1}(\cdot), \dots, p_r e_{q_r}(\cdot)$ among $e_{i_1}(\cdot), \dots, e_{i_m}(\cdot)$ with $p_1 + \dots + p_r = m$.

$0 < p_1, \dots, p_r$, where $H_p(\cdot)$ is the p -th Hermite polynomial with leading coefficient

1. Note also that

$$B = \left(\int_0^1 e_1(u) W_t(du), \dots, \int_0^1 e_N(u) W_t(du) \right)$$

is an N -dimensional Brownian motion.

Define a mapping T_N from C_N to C_1 by

$$T_N(x)(\cdot) = \sum_{1 \leq i_1, \dots, i_m \leq N} c_{i_1, \dots, i_m} x_{q_1}^{p_1}(\cdot) \dots x_{q_r}^{p_r}(\cdot), \text{ for } x = (x_1, \dots, x_N) \in C_N.$$

Then

$$\begin{aligned} X_N(t) - T_N(B)(t) &= \sum_{1 \leq i_1, \dots, i_m \leq N} c_{i_1, \dots, i_m} \{H_{p_1}(\int_0^1 e_{q_1}(u) W_t(du)) \dots H_{p_r}(\int_0^1 e_{q_r}(u) W_t(du)) \\ &\quad - (\int_0^1 e_{q_1}(u) W_t(du))^{p_1} \dots (\int_0^1 e_{q_r}(u) W_t(du))^{p_r}\}, \end{aligned}$$

i.e., $X_N - T_N(B)$ is a finite linear combination of polynomials of degree $\leq m-2$.

Applying Theorem C to each term, we obtain

Lemma 2. For any $\delta > 0$ we have

$$P(|(1/\alpha)|X_N - T_N(B)|_C > \delta) \leq C' \cdot \exp[-C'' \alpha^{2/(m-2)}]$$

for sufficiently large α , where C' and C'' are finite constants.

Since T_N is clearly continuous and homogeneous with degree m , we get from Theorem B

Lemma 3. (i) For any $\varphi \in H_N$ and for any $\delta, \delta' > 0$,

$$P(|T_N(B)/\alpha - T_N(\varphi)|_C < \delta) \geq \exp[-(\alpha^{2/m}/2)(\|\varphi\|_H^2 + \delta')]$$

for sufficiently large α , and

(ii) for any $\delta, \delta', r > 0$,

$$P(d(T_N(B)/\alpha, T_N(K_r)) > \delta) \leq \exp[-(\alpha^{2/m}/2)(r^2 - \delta')]$$

for sufficiently large α .

Let $\{\psi_j, j \geq 0\}$ be a CONS in H_1 . Then $\{e_i \psi_j, i, j \geq 0\}$ is a CONS in $L^2[0,1] \otimes H_1$, and any $f \in L^2[0,1] \otimes H_1$ can be expanded as $f(u, t) = \sum_{i,j=0}^{\infty} c_{ij} e_i(u) \psi_j(t)$ with $\sum_{i,j=0}^{\infty} |c_{ij}|^2 < \infty$. Put $\varphi_i(t) = \sum_{j=0}^{\infty} c_{ij} \psi_j(t)$. Then $\varphi_i \in H_1$ for all $i \geq 0$.

$$f(u, t) = \sum_{i=0}^{\infty} e_i(u) \varphi_i(t) \text{ and } |||f|||^2 = \sum_{i=0}^{\infty} ||e_i||_2^2 ||\varphi_i||_H^2 = \sum_{i=0}^{\infty} ||\varphi_i||_H^2.$$

Now, let $g = Af \in G$ with $f \in L^2[0,1] \otimes H_1$ and define

$$g_N(t) = \int \dots \int_{[0,1]^m} h_N(u_1, \dots, u_m) f(u_1, t) \dots f(u_m, t) du_1 \dots du_m.$$

By Schwarz's inequality,

$$\begin{aligned} |g(t) - g_N(t)| &\leq ||h - h_N||_2 \cdot (\int_0^1 f^2(u, t) du)^{m/2} \\ &= ||h - h_N||_2 \cdot (\sum_{i=0}^{\infty} \varphi_i(t))^m. \end{aligned}$$

and so

$$\begin{aligned} ||g - g_N||_C &\leq ||h - h_N||_2 \cdot (\sum_{i=0}^{\infty} ||\varphi_i||_C^2)^{m/2} \\ &\leq ||h - h_N||_2 \cdot (\sum_{i=0}^{\infty} ||\varphi_i||_H^2)^{m/2} \\ &= ||h - h_N||_2 \cdot |||f|||^m. \end{aligned}$$

Note that

$$\begin{aligned} g_N(t) &= \sum_{1 \leq i_1, \dots, i_m \leq N} c_{i_1 \dots i_m} \int \dots \int_{[0,1]^m} e_{i_1}(u_1) \dots e_{i_m}(u_m) \cdot \\ &\quad f(u_1, t) \dots f(u_m, t) du_1 \dots du_m \\ &= \sum_{1 \leq i_1, \dots, i_m \leq N} c_{i_1 \dots i_m} (\int_0^1 e_{i_1}(u) f(u, t) du) \dots (\int_0^1 e_{i_m}(u) f(u, t) du) \\ &= \sum_{1 \leq i_1, \dots, i_m \leq N} c_{i_1 \dots i_m} \varphi_{i_1}(t) \dots \varphi_{i_m}(t) \\ &= \sum_{1 \leq i_1, \dots, i_m \leq N} c_{i_1 \dots i_m} (\varphi_{q_1}(t))^{p_1} \dots (\varphi_{q_r}(t))^{p_2} \\ &= T_N(\varphi)(t), \text{ where } \varphi = (\varphi_1, \dots, \varphi_N) \in H_N. \end{aligned}$$

and also that, for any given $\varphi = (\varphi_1, \dots, \varphi_N) \in H_N$, if we put $f(u, t)$

$$= \sum_{i=1}^N e_i(u) \varphi_i(t), \text{ then } f \in L^2[0,1] \otimes H_1, |||f||| = ||\varphi||_H \text{ and } Af = T_N(\varphi).$$

From the above we immediately obtain the following

Lemma 4 (i) For any $g = Af \in G$ with $f \in L^2[0,1] \otimes H_1$ and for any $\delta > 0$, there is

an element g_N of the form $g_N = T_N(\varphi)$ with $\varphi \in H_N$, $\|\varphi\|_H \leq \|f\|$ such that $\|g - g_N\|_C < \delta$ for all sufficiently large N .

(11) $T_N(K_r) \subset G_r$ for all $r > 0$, $N \geq 1$.

Proof of Theorem 1 (i) Let $g \in G$, $\delta, \delta' > 0$ be given, and assume that g is of the form $g = Af$, $f \in L^2[0,1] \otimes H_1$. Choose N large enough that $\|g - g_N\|_C < \delta/4$ with $g_N = T_N(\varphi)$, $\varphi \in H_N$ (Lemma 4 (1)), and $M_N > (4/\delta)^{2/m} (\|f\|^2 + \delta')$ (cf. Lemma 1). Then, by Lemmas 1, 2 and 3(1),

$$\begin{aligned} P(\|X/\alpha - g\|_C < \delta) &\geq P(\|T_N(B)/\alpha - T_N(\varphi)\|_C < \delta/4) \\ &\quad - P(\|X - X_N\|_C > \alpha\delta/4) \\ &\quad - P(\|X_N - T_N(B)\|_C > \alpha\delta/4) \\ &\geq \exp[-(\alpha^{2/m}/2)(\|\varphi\|_H^2 + \delta'/3)] \\ &\quad - C \cdot \exp\{-M_N(\alpha\delta/4)^{2/m}\} \\ &\quad - C' \cdot \exp[-C' \cdot \alpha^{2/(m-2)}] \end{aligned}$$

for all sufficiently large α . Noting that $\|\varphi\|_H \leq \|f\|$, we thus obtain

$$P(\|X/\alpha - g\|_C < \delta) \geq \exp[-(\alpha^{2/m}/2)(\|f\|^2 + 2\delta'/3)]$$

for all sufficiently large α , and the assertion follows from the definition of $D(g)$.

Proof of Theorem 1 (ii) Given δ, δ' , $r > 0$, choose N large enough that

$M_N > (3/\delta)^{2/m}(r^2 - \delta')$ (cf. Lemma 1). Then, by Lemmas 1, 2, 3(11) and 4(11),

$$\begin{aligned} P(d(X/\alpha, G_r) > \delta) &\leq P(d(X/\alpha, T_N(K_r)) > \delta) \\ &\leq P(d(T_N(B)/\alpha, T_N(K_r)) > \delta/3) \\ &\quad + P(\|X - X_N\|_C > \alpha\delta/3) + P(\|X_N - T_N(B)\|_C > \alpha\delta/3) \\ &\leq \exp[-(\alpha^{2/m}/2)(r^2 - \delta'/2)] \\ &\quad + C \cdot \exp[-M_N(\alpha\delta/3)^{2/m}] + C' \cdot \exp[-C' \cdot \alpha^{2/(m-2)}] \\ &\leq \exp[-(\alpha^{2/m}/2)(r^2 - \delta')] \end{aligned}$$

for all sufficiently large α . This completes the proof of Theorem 1.

3. Proof of Theorem 2

It follows from Lemmas 1 and 2 that

$$P(|X - T_N(B)|_C > z) \leq \exp[-Mz^{2/m}]$$

for all sufficiently large z and N , where M is any given positive constant.

Note also that both processes $X(\cdot)$ and $T_N(B)(\cdot)$ are self-similar with parameter $m/2$, i.e., $X(c\cdot)$ and $c^{m/2}X(\cdot)$, and also $T_N(B)(c\cdot)$ and $c^{m/2}T_N(B)(\cdot)$, have the same finite dimensional distributions for any $c > 0$. Using these facts and the first Borel-Cantelli lemma, we get by the standard argument

Lemma 5 For any $\epsilon > 0$, with probability 1

$$\limsup_{n \rightarrow \infty} \|Z_N - T_N(B)(nt)/(2n \log_2 n)^{m/2}\|_C \leq \epsilon$$

for sufficiently large N .

Note that

$$T_N(B)(nt)/(2n \log_2 n)^{m/2} = T_N(B_n)(t),$$

where $B_n(t) = B(nt)/(2n \log_2 n)^{1/2}$. Thus, by Strassen's law of the iterated logarithm for B and the continuous mapping theorem, we have

Lemma 6 For any $N \geq 1$, with probability 1 the random sequence $\{T_N(B_n), n \geq 3\}$ in C_1 is relatively compact and the set of its limit points is $T_N(K_1) = \{T_N(\varphi); \varphi \in K_1\}$.

Theorem 2 follows from Lemmas 4, 5 and 6.

References

- [1] Dehling, H.: The functional law of the iterated logarithm for von-Mises-functionals and multiple Wiener integrals. Preprint, 1984 (revised 1987).
- [2] Kallianpur, G. and Oodaira, H.: Freidlin-Wentzell type estimates for abstract Wiener spaces. Sankhya, Ser. A, 40, 116-137 (1978).
- [3] Mandelbaum, A. and Taqqu, M.S.: Invariance principle for symmetric statistics. Ann. Statistics, 12, 483-496 (1984).
- [4] Mori, T. and Oodaira, H.: The law of the iterated logarithm for self-similar processes represented by multiple Wiener integrals. Probab. Th. Rel. Fields, 71, 367-391 (1986).
- [5] Plikusas, A.: Some properties of the multiple Ito $\hat{\int}$ integral (in Russian). Liet. Mat. Rink., 21, 163-173 (1981).

END

11-87

DTIC