

AO-A106 735

MEASURING THE DEPENDENCE BETWEEN TWO POINT PROCESSES

1/1

THROUGH CONFIDENCE I. (U) FLORIDA STATE UNIV


TALLAHASSEE DEPT OF STATISTICS H DOSS SEP 87

UNCLASSIFIED

FSU-STATISTICS-M767 AFOSR-TR-87-1531

F/G 12/3

NL



1.0  
1.1  
1.25  
1.4  
1.6  
1.8  
2.0  
2.2  
2.5  
2.8  
3.2  
3.6  
4.0  
4.5  
5.0  
5.6  
6.3  
7.1  
8.0  
9.0  
10.0  
11.2  
12.5  
14.0  
16.0  
18.0  
20.0  
22.4  
25.0  
28.0  
31.5  
36.0  
40.0  
45.0  
50.0  
56.0  
63.0  
71.0  
80.0  
90.0  
100.0

AD-A186 735

OTC FILE COPY

## REPORT DOCUMENTATION PAGE

|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |  |                                                                                                                                                                                                             |  |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|
| 1a. REPORT SECURITY CLASSIFICATION<br><b>UNCLASSIFIED</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |  | 1b. RESTRICTIVE MARKINGS                                                                                                                                                                                    |  |
| 2a. SECURITY CLASSIFICATION AUTHORITY<br><b>NA</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |  | 3. DISTRIBUTION/AVAILABILITY OF REPORT<br>Approved for Public Release; Distribution Unlimited                                                                                                               |  |
| 2b. DECLASSIFICATION/DOWNGRADING SCHEDULE<br><b>NA</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |  | 5. MONITORING ORGANIZATION REPORT NUMBER(S)<br><b>AFOSR-TR-87-21531</b>                                                                                                                                     |  |
| 4. PERFORMING ORGANIZATION REPORT NUMBER<br>FSL Technical Report No. M-767                                                                                                                                                                                                                                                                                                                                                                                                                                                |  | 7a. NAME OF MONITORING ORGANIZATION<br>AFOSR/NM                                                                                                                                                             |  |
| 6a. NAME OF PERFORMING ORGANIZATION<br>Florida State University                                                                                                                                                                                                                                                                                                                                                                                                                                                           |  | 7b. ADDRESS (City, State and ZIP Code)<br>Bldg. 410<br>Bolling AFB, DC 20332-6448                                                                                                                           |  |
| 6b. ADDRESS (City, State and ZIP Code)<br>Department of Statistics<br>Tallahassee, FL 32306-3033                                                                                                                                                                                                                                                                                                                                                                                                                          |  | 9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER                                                                                                                                                             |  |
| 8a. NAME OF FUNDING/SPONSORING ORGANIZATION<br>AFOSR                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |  | 8b. OFFICE SYMBOL (If applicable)<br>NM                                                                                                                                                                     |  |
| 8c. ADDRESS (City, State and ZIP Code)<br>Bldg. 410<br>Bolling AFB, DC                                                                                                                                                                                                                                                                                                                                                                                                                                                    |  | 10. SOURCE OF FUNDING NOS.<br>PROGRAM ELEMENT NO. 6.1102F<br>PROJECT NO. 2304<br>TASK NO. 115<br>WORK UNIT NO.                                                                                              |  |
| 11. TITLE (Include Security Classification) Measuring the Dependence Between Two Point Processes Through Confidence Intervals for the Second Order Distribution                                                                                                                                                                                                                                                                                                                                                           |  |                                                                                                                                                                                                             |  |
| 12. PERSONAL AUTHOR(S)<br>Hani Poss                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |  |                                                                                                                                                                                                             |  |
| 13a. TYPE OF REPORT<br>Technical Journal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |  | 13b. TIME COVERED<br>FROM TO                                                                                                                                                                                |  |
| 14. DATE OF REPORT (Yr., Mo., Day)<br>September, 1987                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |  | 15. PAGE COUNT<br>20                                                                                                                                                                                        |  |
| 16. SUPPLEMENTARY NOTATION<br>Also issued as Stanford University Technical Report, Supported by Office of Naval Research Grant No. N00014-86-K-0156.                                                                                                                                                                                                                                                                                                                                                                      |  |                                                                                                                                                                                                             |  |
| 17. COSATI CODES<br>FIELD GROUP SUB GR                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |  | 18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)<br>Bivariate point process; Ripley's k-function; cross-intensity function; stationary point process; stationary sequence. |  |
| 19. ABSTRACT (Continue on reverse if necessary and identify by block number)<br><br>To assess the dependence structure in a stationary bivariate point process the second-order distribution (Ripley 1976, 1977) <sup>3</sup> can be very useful. We prove that the natural estimates of this distribution, based on a realization $A_1 < A_2 < \dots < A_{n_A}, B_1 < B_2 < \dots < B_{n_B}$ are asymptotically normal, and we present a method for constructing approximate confidence intervals for this distribution. |  |                                                                                                                                                                                                             |  |
| 20. DISTRIBUTION/AVAILABILITY OF ABSTRACT<br>UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>                                                                                                                                                                                                                                                                                                                                         |  | 21. ABSTRACT SECURITY CLASSIFICATION<br>UNCLASSIFIED                                                                                                                                                        |  |
| 22a. NAME OF RESPONSIBLE INDIVIDUAL<br>Frank Prochen                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |  | 22b. TELEPHONE NUMBER (Include Area Code)<br>(904) 644-3218                                                                                                                                                 |  |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |  | 22c. OFFICE SYMBOL<br>AFOSR/NM                                                                                                                                                                              |  |


**Measuring the Dependence Between Two Point Processes  
Through Confidence Intervals for the Second Order Distribution**

by

**AFOSR-TR- 87 - 1531**

Hani Doss

Department of Statistics  
Florida State University  
Tallahassee, FL 32306



FSU Technical Report No. M-767  
AFOSR Technical Report No. 87-210

September, 1987

|                    |       |
|--------------------|-------|
| Accession For      |       |
| NTIS               | CRA&I |
| DTIC               | TAB   |
| Unannounced        |       |
| Justification      |       |
| By                 |       |
| Distribution       |       |
| Availability Codes |       |
| DIA                | AFOSR |
| A-1                |       |

Key words and phrases: Bivariate point process; Ripley's K-function; cross-intensity function; stationary point process; stationary sequence.

AMS 1980 subject classifications. Primary 62M09, 62M07; secondary 62G05, 62G10.

Abbreviated title: Estimation of Dependence Between Point Processes.

Research supported by the Air Force Office of Scientific Research Grant No. ~~AFOSR~~ 85-C-0007.

87 10 15 051

**Measuring the Dependence Between Two Point Processes  
Through Confidence Intervals for the Second Order Distribution**

by

**Hani Doss**

**ABSTRACT**

To assess the dependence structure in a stationary bivariate point process the second-order distribution (Ripley 1976, 1977) can be very useful. We prove that the natural estimates of this distribution, based on a realization  $A_1 < A_2 < \dots < A_{n_A}, B_1 < B_2 < \dots < B_{n_B}$  are asymptotically normal, and we present a method for constructing approximate confidence intervals for this distribution.

## 1. INTRODUCTION AND SUMMARY.

Let  $(N_A, N_B)$  be a stationary bivariate point process on  $\mathbb{R}$ . This article is concerned with statistical methods for discovering and quantifying an association between the two processes from a realization  $A_1 < A_2 < \dots < A_{n_A}, B_1 < B_2 < \dots < B_{n_B}$  over a long period of time  $T$ . The paper is motivated by certain problems that arise in neurophysiology, which are very briefly described as follows (for further details see e.g. Bryant, Ruiz Marcos, and Segundo, 1973).

Two neurons,  $A$  and  $B$ , are monitored over a period of time  $T$  during which each neuron fires a sequence of impulses. The problem is to determine whether or not the impulse times are associated. An association between  $N_A$  and  $N_B$  may be construed as evidence that either the two neurons are communicating, or that they both share input from a third source.

Another problem arises in certain neurophysiological studies of learning and memory. An animal is to be taught (trained) to perform a certain task. Now consider two "connected" neurons,  $A$  and  $B$ , which are essential in the performance of this task. Record the impulse times during a period before the learning experience, obtaining a realization of  $(N_A^{Bef.}, N_B^{Bef.})$ , and during a period of time *well after* the learning experience, obtaining a realization of  $(N_A^{Aft.}, N_B^{Aft.})$ . The processes  $N_A^{Bef.}$  and  $N_B^{Bef.}$  may be *dependent*. The problem is to determine whether or not this dependence is "stronger" for the processes  $N_A^{Aft.}$  and  $N_B^{Aft.}$ . A neurophysiologist may consider a change in the strength of the dependence as evidence that learning has taken place.

The two problems have very different statistical character. Let  $S$  be a statistic that "measures" the dependence between two point processes. The first problem is one of testing the hypothesis that  $N_A$  and  $N_B$  are independent, and requires only knowledge of the distribution of  $S$  under the assumption that  $N_A$  and  $N_B$  are independent. The second problem is much more difficult: to compare  $S$  across two situations we must know the distribution of  $S$  when the two point processes are dependent.

In a more general context, Ripley (1976, 1977) introduced a measure  $K$ , defined on an appropriate space, that summarizes the second-order properties of the process. Before describing this measure, we need to state some assumptions and introduce some notation. Let  $N_i(s, t)$  denote the number of events of type  $i$  occurring in the interval  $(s, t]$ , for  $i = A, B$ . Assume that each process has no multiple occurrences, and that the intensities

$$\lambda_i = \lim_{h \rightarrow 0} \frac{1}{h} P\{N_i(t, t+h) > 0\} \quad \text{for } i = A, B \quad (1)$$

are finite. (The existence of these limits was proved by Khintchine, 1960). The  $\lambda_i$ 's then have an interpretation as mean number of occurrences per unit of time: for  $t_1 < t_2$ ,

$$E N_i(t_1, t_2) = \lambda_i(t_2 - t_1) \quad \text{for } i = A, B \quad (2)$$

(this follows from Dobrushin's Lemma and Korolyuk's Theorem; see Leadbetter, 1968).

We now give an informal description of the measure  $K$ , adapted to the present context.

The measure  $K$  is defined on the Borel subsets of  $\mathbb{R}$ , and for  $t_1 < t_2$ , writing  $K(t_1, t_2)$  for  $K\{(t_1, t_2)\}$ , we have

$$\begin{aligned} K(t_1, t_2) &= \frac{1}{\lambda_A} E\{N_A(t_1, t_2) \mid \text{a } B \text{ point at } t = 0\} \\ &\quad \left( = \frac{1}{\lambda_B} E\{N_B(-t_2, -t_1) \mid \text{an } A \text{ point at } t = 0\} \right). \end{aligned} \quad (3)$$

Note that if  $N_A$  and  $N_B$  are independent, then

$$K(t_1, t_2) = t_2 - t_1, \quad (4)$$

regardless of the values  $\lambda_A$  and  $\lambda_B$ ; thus,  $K$  represents Lebesgue measure on  $\mathbb{R}$ .

Ripley proposed the estimate of  $K(t_1, t_2)$  given by

$$\hat{K}(t_1, t_2) = \frac{T}{n_A n_B} \sum_{i=1}^{n_B} \sum_{j=1}^{n_A} I\{A_j - B_i \in (t_1, t_2)\}, \quad (5)$$

where  $I\{\cdot\}$  denotes the indicator function (actually, the estimate proposed by Ripley has an edge correction for points near the boundary of the period of observation; this edge correction will not concern us).

Previous work on the estimator  $\hat{K}$  is concerned with spatial processes. The results center on using  $\hat{K}$  to test that a single process is Poisson (Ripley, 1977; Chapter 8 of Ripley, 1981; Silverman, 1976) and on using  $\hat{K}$  to test for independence of two processes (Lotwick and Silverman, 1982; Diggle and Milne, 1983).

In this paper we study the asymptotic properties of  $\hat{K}(t_1, t_2)$ . The contributions of this paper are two-fold:

first, a proof that under certain regularity conditions, as  $n_B \rightarrow \infty$

$$\sqrt{n_B}(\hat{K}(t_1, t_2) - K(t_1, t_2)) \xrightarrow{d} \mathcal{N}(0, \sigma^2(t_1, t_2)), \quad (6)$$

and second, a method for constructing consistent estimates of  $\sigma^2(t_1, t_2)$ . (7)

Besides providing the basis for a test of independence between  $N_A$  and  $N_B$ , (6) and (7) enable one to test whether or not  $K(t_1, t_2)$  has changed in the experimental situation described earlier.

The cross-intensity function defined by

$$\lambda_{AB}(u) = \lim_{h_1, h_2 \rightarrow 0} \frac{1}{h_1 h_2} P\{N_A(u+t, u+t+h_1) > 0; N_B(t, t+h_2) > 0\} \quad (8)$$

is related to  $K$  by

$$K(t_1, t_2) = \frac{1}{\lambda_A \lambda_B} \int_{t_1}^{t_2} \lambda_{AB}(u) du. \quad (9)$$

Under the independence hypothesis,  $\lambda_{AB}(u) = \lambda_A \lambda_B$  for all  $u$ . Brillinger (1976) considered the random function

$$J_{AB}^T(u) = \sum_{i=1}^{n_B} \sum_{j=1}^{n_A} I\{A_j - B_i \in (u-h, u+h)\} \quad (10)$$

and showed that under suitable regularity, if  $h \rightarrow 0$  and  $T \rightarrow \infty$  in such a way that  $hT$  remains constant, then for  $u_k^T \rightarrow u_k, |u_k^T - u_{k'}^T| > 2h, 1 \leq k < k' \leq M, J_{AB}^T(u_k^T)$  are asymptotically independent Poisson random variables with means  $2hT\lambda_{AB}(u_k)$ , for  $k = 1, \dots, M$ . Thus,  $\hat{\lambda}_{AB}(u) = \frac{J_{AB}^T(u)}{2hT}$  can be used to estimate  $\lambda_{AB}(u)$  at a finite number of points.

In practice one would graph the two functions  $\hat{\lambda}_{AB}$  and  $\hat{K}$  over a finite range, say  $[-L, L]$  (i.e. graph  $\hat{K}(-L, t)$  for  $-L \leq t \leq L$ ). Although from a mathematical viewpoint  $\lambda_{AB}$  and  $K$  contain essentially the same information, the statistical properties of their estimates are quite different: estimation of  $\lambda_{AB}$  is akin to estimating a density, and from Brillinger's result the variance of  $\hat{\lambda}_{AB}$  is of the order  $\frac{1}{hT}$ ; on the other hand, estimation of  $K$  is akin to estimating a distribution function, and from (6), the variance of  $\hat{K}$  is of the smaller order  $\frac{1}{n_B}$ . A graph of  $\hat{\lambda}_{AB}$  may, however, indicate features (spikes, location of maxima and minima, etc.) that cannot be easily seen in the graph of  $\hat{K}$ . Clearly the two approaches are complementary.

## 2. ASYMPTOTIC DISTRIBUTION OF THE $\hat{K}$ - FUNCTION.

Let

$$U_{AB}(t_1, t_2) = E\{N_A(t_1, t_2) \mid \text{a } B \text{ point at } t = 0\}. \quad (11)$$

We may estimate  $U_{AB}(t_1, t_2)$  by

$$\hat{U}_{AB}(t_1, t_2) = \frac{1}{n_B} \sum_{i=1}^{n_B} \sum_{j=1}^{n_A} I\{A_j - B_i \in (t_1, t_2)\}. \quad (12)$$

Letting

$$\hat{\lambda}_i = \frac{n_i}{T} \quad \text{for } i = A, B, \quad (13)$$

we note that

$$K(t_1, t_2) = \frac{1}{\lambda_A} U_{AB}(t_1, t_2), \quad \text{and} \quad \hat{K}(t_1, t_2) = \frac{1}{\hat{\lambda}_A} \hat{U}_{AB}(t_1, t_2). \quad (14)$$

To prove asymptotic normality of  $\hat{K}$  (Theorem 2) we will prove joint asymptotic normality of  $(\hat{U}_{AB}(t_1, t_2), \hat{\lambda}_A)$ . We will in fact find it necessary to first prove joint asymptotic normality of  $(\hat{U}_{AB}(t_1, t_2), \hat{\lambda}_A/\hat{\lambda}_B, 1/\hat{\lambda}_B)$ . The delta-method (i.e. a first term Taylor expansion) applied to the function  $f(x, y, z) = \frac{xz}{y}$  then yields the asymptotic normality of  $\hat{K}(t_1, t_2)$ . We also obtain the joint asymptotic normality  $(\hat{U}_{AB}(t_1, t_2), \hat{\lambda}_A, \hat{\lambda}_B)$  by applying the delta-method to the function  $g(x, y, z) = (x, y/z, 1/z)$ .

We now need to give the statistical setting of our asymptotic investigation. The functions  $U_{AB}(t_1, t_2)$  and  $K(t_1, t_2)$  involve the notion of the Palm measure. That is for  $\varepsilon > 0$ , we consider the conditional distribution of the process  $(N_A, N_B)$  given that there is a  $B$  point in the interval  $(0, \varepsilon)$ , and take the limiting distribution of  $(N_A, N_B)$  as  $\varepsilon \rightarrow 0$ . Intuitively, this corresponds to selecting a  $B$  point "arbitrarily", and considering the process with that point labeled the origin. This notion is discussed for univariate processes by Leadbetter (1972) and for bivariate processes by Wisniewski (1972). We will assume that the process is observed during a period of length  $T$  starting immediately after the occurrence of an "arbitrary"  $B$  point, say  $B_0$  (thus, we will be working with the Palm measure). This mode of sampling is called semisynchronous sampling by Cox and Lewis (1972); see Wisniewski (1972) for some fundamental properties related to it. Also, for the sake of convenience, we will assume that the period of observation ends with a  $B$  point.

We now consider two subfields of the  $\sigma$ -field on which the Palm measure is defined. Let  $\mathcal{F}_{-\infty}^{B_0}$  denote the  $\sigma$ -field generated by the events

$$\{B_{k_1} \in (B_0 + v_1, B_0 + w_1), \dots, B_{k_m} \in (B_0 + v_m, B_0 + w_m);$$

$$N_A(B_0 + r_1, B_0 + s_1) = h_1, \dots, N_A(B_0 + r_n, B_0 + s_n) = h_n\}$$

for  $v_i < w_i \leq 0, k_i = -1, -2, \dots, i = 1, \dots, m; r_j < s_j \leq 0, h_j = 0, 1, 2, \dots, j = 1, \dots, n$ , and  $m$  and  $n$  nonnegative integers. For  $u > 0$ , let  $\mathcal{F}_{B_0+u}^{\infty}$  denote the  $\sigma$ -field generated by the events

$$\{B_{k_1} \in (B_0 + v_1, B_0 + w_1), \dots, B_{k_m} \in (B_0 + v_m, B_0 + w_m);$$

$$N_A(B_0 + r_1, B_0 + s_1) = h_1, \dots, N_A(B_0 + r_n, B_0 + s_n) = h_n\}$$

for  $u \leq v_i < w_i, k_i = 1, 2, \dots, i = 1, \dots, m; u \leq r_j < s_j, h_j = 0, 1, 2, \dots, j = 1, \dots, n$ , and  $m$  and  $n$  nonnegative integers. Let

$$\alpha(u) = \sup\{|P(E_1 \cap E_2) - P(E_1)P(E_2)|; E_1 \in \mathcal{F}_{-\infty}^{B_0}, E_2 \in \mathcal{F}_{B_0+u}^{\infty}\}. \quad (15)$$

If  $\alpha(u) \rightarrow 0$  as  $u \rightarrow \infty$ , then the distant future is virtually independent of the past. We will actually need stronger conditions on  $\alpha(\cdot)$ .

Let  $\beta > 0, \eta > 1, 0 < \tau < 1$  be any constants satisfying

$$\left(\eta - \frac{\eta + 1}{\beta + 1}\right) \tau > 1. \quad (16)$$

Assumptions:

$$A1 \quad \int_0^{\infty} [\alpha(t)]^{\tau} t^{\beta} dt < \infty$$

$$A2 \quad \sup_{-\infty < j < \infty} E\left\{|N_B(j, j+1)|^{\eta} \mid \text{a } B \text{ point at } t = 0\right\} = D < \infty$$

$$A3 \quad E\left\{|N_A(t_1, t_2)|^{4(1+\frac{\tau}{1-\tau})} \mid \text{a } B \text{ point at } t = 0\right\} < \infty$$

$$A4 \quad E\left[(B_1 - B_0)^{4(1+\frac{\tau}{1-\tau})}\right] < \infty$$

$$A5 \quad E\left[N_A(B_0, B_1)\right]^{4(1+\frac{\tau}{1-\tau})} < \infty$$

Before stating our theorems we discuss our assumptions, and compare them with those of Brillinger (1976). Any theorem giving asymptotic normality of the normalized partial sums of a

stationary sequence  $\{T_i\}$  must assume a moment condition on  $T_1$  and also a mixing condition on  $\{T_i\}$ . In general, weakening of the moment condition must be compensated by strengthening of the mixing condition, and vice versa. Assumptions A3, A4, and A5 provide moment conditions on the sequences  $\{U_i\}$ ,  $\{N_A(B_{i-1}, B_i)\}$ , and  $\{(B_i - B_{i-1})\}$ , respectively. Assumption A2 ensures that the  $B$  process "moves along" rapidly enough so that A1, the mixing condition imposed on the point process, translates into a mixing condition for the sequences  $\{U_i\}$ ,  $\{N_A(B_{i-1}, B_i)\}$ , and  $\{(B_i - B_{i-1})\}$ . Relationship (16) describes in a technical way the interplay between the mixing rate on the point process and the moment condition on the sequences  $\{U_i\}$ ,  $\{N_A(B_{i-1}, B_i)\}$ , and  $\{(B_i - B_{i-1})\}$ .

The conditions assumed by Brillinger (1976) neither imply nor are implied by A1-A5 of the present paper. Brillinger assumes a mixing condition on the bivariate point process and also that the "second order moments"  $\lambda_{ij}(\cdot)$  ( $i, j = A, B$ ) exist and are continuous (he also assumes existence and continuity of the "third and fourth order moments" ; see equation (2.2) of his paper). This condition on  $\lambda_{AB}(\cdot)$  is not satisfied by the following process:  $N_B$  is a Poisson process, and  $N_A$  is  $N_B$  shifted to the right by 1 unit. In this case,  $\lambda_{AB}(1) = \infty$ . This process does however satisfy A1-A5. Conversely, it is easy to find processes  $(N_A, N_B)$  satisfying all of Brillinger's conditions, but not those of the present paper. Perhaps the simplest example is the following. Let  $N_A$  and  $N_B$  be independent,  $N_A$  being a Poisson process, and  $N_B$  being an equilibrium renewal process on  $(-\infty, \infty)$  (for a definition and a construction see pp. 517-519 of Karlin and Taylor, 1975) with interarrival distribution having a first moment but no second moment. Then A4 is violated, and it is not difficult to check that this process satisfies all of Brillinger's conditions.

**THEOREM 1.** Assume A1 and A2. Let  $U_{AB}(t_1, t_2)$ ,  $\hat{U}_{AB}(t_1, t_2)$ , and  $\hat{\lambda}_i$  for  $i = A, B$  be defined by (11), (12), and (13), respectively.

(i) Under A3, we have as  $n_B \rightarrow \infty$

$$\sqrt{n_B}(\hat{U}_{AB}(t_1, t_2) - U_{AB}(t_1, t_2)) \xrightarrow{d} \mathcal{N}(0, \gamma^2(t_1, t_2)).$$

Furthermore, any estimate  $\hat{\gamma}^2(t_1, t_2)$  of the form (25), satisfying (27) and (28), is a consistent estimate of  $\gamma^2(t_1, t_2)$ .

(ii) Under A4 and A5 we have as  $n_B \rightarrow \infty$

$$\sqrt{n_B}(\hat{\lambda}_A - \lambda_A, \hat{\lambda}_B - \lambda_B)' \xrightarrow{d} \mathcal{N}(0, \Lambda).$$

Furthermore, any estimate  $\hat{\Lambda}$  of the form (57) (refer to equations (49) and (51)-(56)), satisfying (27) and (28) is a consistent estimate of  $\Lambda$ .

(iii) Under A3-A5 we have as  $n_B \rightarrow \infty$

$$\sqrt{n_B}(\hat{U}_{AB}(t_1, t_2) - U_{AB}(t_1, t_2), \hat{\lambda}_A - \lambda_A, \hat{\lambda}_B - \lambda_B)' \xrightarrow{d} \mathcal{N}(0, \Sigma(t_1, t_2)).$$

Furthermore, any estimate  $\hat{\Sigma}(t_1, t_2)$  of the form (56) (refer to equations (49) and (51)-(55)), satisfying (27) and (28), is a consistent estimate of  $\Sigma(t_1, t_2)$ .

PROOF of (i). We begin by showing asymptotic normality. Let  $U'_i$  and  $U_i$  be defined by

$$U'_i = \sum_{j=-\infty}^{\infty} I\{A_j \in (B_i + t_1, B_i + t_2)\} I\{B_0 \leq A_j \leq B_{n_B}\}$$

and

$$U_i = \sum_{j=-\infty}^{\infty} I\{A_j \in (B_i + t_1, B_i + t_2)\}.$$

Note that

$$\hat{U}_{AB}(t_1, t_2) = \frac{1}{n_B} \sum_{i=1}^{n_B} U'_i. \quad (17)$$

It is clear that  $\sum_{i=1}^{n_B} U_i - \sum_{i=1}^{n_B} U'_i = 0_p(1)$ . Thus, it suffices to prove the result with  $U_i$ 's instead of  $U'_i$ 's in (17). Observe that the sequence  $\{U_i\}_{i=-\infty}^{\infty}$  is stationary, with mean  $U_{AB}(t_1, t_2)$  and finite variance (by A3). The  $U_i$ 's may be far from independent: for small  $k$ ,  $U_i$  and  $U_{i+k}$  may be nearly identical. If, however,  $U_i$  and  $U_{i+k}$  are "nearly independent" for large  $k$ , then one can still hope to have a Central Limit Theorem effect. The proof consists of translating A1, the mixing condition on the point process, into a mixing condition on  $\{U_i\}$  that allows the application of an appropriate central limit theorem for stationary sequences.

Let  $\mu(k)$  be defined for  $k = 1, 2, \dots$  by

$$\mu(k) = \sup\{|P(E_1 \cap E_2) - P(E_1)P(E_2)|; E_1 \in \sigma(\dots, U_{-1}, U_0), E_2 \in \sigma(U_k, U_{k+1}, \dots)\} \quad (18)$$

(Here,  $\sigma(\dots, U_{-1}, U_0)$  denotes the  $\sigma$ -field generated by  $\{\dots, U_{-1}, U_0\}$ , and similarly for  $\sigma(U_k, U_{k+1}, \dots)$ ). The function  $\mu(\cdot)$  is called the mixing coefficient of the sequence  $\{U_i\}$ . Our goal is to prove that  $\sum_{k=1}^{\infty} [\mu(k)]^r < \infty$ . It will be more convenient however, to show instead that  $\sum_{k=1}^{\infty} [\mu(2k)]^r < \infty$ . The two conditions are equivalent since  $\mu(\cdot)$  is nonincreasing.

Let  $k \geq 1$  be fixed, let  $E_1 \in \sigma(\dots, U_{-k})$ ,  $E_2 \in \sigma(U_k, \dots)$ , and consider  $P(E_1 \cap E_2)$ . Let

$$C_{-k} = \{B_{-k} - B_0 \leq -[k^{1/(\beta+1)}]\} \quad \text{and} \quad C_k = \{B_k - B_0 \geq [k^{1/(\beta+1)}]\}.$$

We may write

$$P(E_1 \cap E_2) = P\{(E_1 \cap C_{-k}) \cap (E_2 \cap C_k)\} + P\{(E_1 \cap E_2) \cap (C_{-k}^c \cup C_k^c)\}, \quad (19)$$

where  $^c$  denotes complementation.

Consider the first term on the right side of (19). For all large  $k$ , since  $B_{-k} - B_0 \leq -k^{1/(\beta+1)}$  implies that  $B_{-k} + t_2 \leq B_0$ , we have  $E_1 \cap C_{-k} \in \mathcal{F}_{-\infty}^{B_0}$ . Furthermore,  $E_2 \cap C_k \in \mathcal{F}_{B_0 + [k^{1/(\beta+1)}] + t_1}^\infty$ . Therefore,

$$P\{(E_1 \cap C_{-k}) \cap (E_2 \cap C_k)\} \leq P(E_1)P(E_2) + \alpha([k^{1/(\beta+1)}] + t_1). \quad (20)$$

The second term on the right side of (19) is obviously less than or equal to  $P(C_{-k}^c) + P(C_k^c)$ . These last two probabilities are dealt with in the same way. Consider  $P(C_k^c)$ . Observe that

$$P(C_k^c) \leq P\{\text{one of the intervals } (B_0 + j, B_0 + j + 1), j = 0, 1, \dots, [k^{1/(\beta+1)}] - 1 \text{ has at least } [k^{\beta/(\beta+1)}] \text{ points}\}. \quad (21)$$

By A2, Chebyshev's inequality and Boole's inequality, the right side of (21) is less than or equal to  $[k^{1/(\beta+1)}] D [k^{\beta/(\beta+1)}]^{-\eta}$ . Combining this with (20) and handling the opposite inequality in a similar way, we obtain

$$\mu(2k) \leq \alpha(k^{1/(\beta+1)} + t_1) + k^{1/(\beta+1)} D [k^{\beta/(\beta+1)}]^{-\eta}.$$

Assumption A1 implies that

$$\sum_{k=1}^{\infty} \{\alpha(k^{1/(\beta+1)} + t_1)\}^\tau < \infty. \quad (22)$$

Combining (22) and (16) we obtain that  $\sum_{k=1}^{\infty} [\mu(2k)]^\tau < \infty$ , and hence that

$$\sum_{k=1}^{\infty} [\mu(k)]^\tau < \infty. \quad (23)$$

Assumption A3 implies that

$$E[U_1^{2(1+\frac{1}{1-\tau})}] < \infty.$$

This, together with (23) allows us to apply Theorem 1.7 of Ibragimov (1962) to conclude that the series  $\sum_{h=1}^{\infty} \text{Cov}(U_0, U_h)$  converges absolutely, and that as  $n_B \rightarrow \infty$

$$\sqrt{n_B}(\hat{U}_{AB}(t_1, t_2) - U_{AB}(t_1, t_2)) \xrightarrow{d} \mathcal{N}(0, \gamma^2(t_1, t_2)),$$

where

$$\gamma^2(t_1, t_2) = \text{Var } U_0 + 2 \sum_{h=1}^{\infty} \text{Cov}(U_0, U_h). \quad (24)$$

Consider next the estimation of  $\gamma^2(t_1, t_2)$ . Let  $\nu_h = \text{Cov}(U_0, U_h)$  for  $h = 0, 1, 2, \dots$ , so that  $\gamma^2(t_1, t_2) = \nu_0 + 2 \sum_{h=1}^{\infty} \nu_h$ . Let

$$\hat{\gamma}^2(t_1, t_2) = \hat{\nu}_0 + 2 \sum_{h=1}^M c_h \hat{\nu}_h \quad (25)$$

where  $\hat{\nu}_h$  is the sample covariance at lag  $h$ :

$$\hat{\nu}_h = \frac{1}{n_B - h} \sum_{i=0}^{n_B - h - 1} (U_i - \bar{U})(U_{i+h} - \bar{U}) \quad (26)$$

(here  $\bar{U} = \frac{1}{n_B} \sum_{i=1}^{n_B} U_i$ ). We will assume that

$$M = M_{n_B} \text{ satisfies } \frac{M_{n_B}}{(n_B)^{\frac{1}{3}}} \rightarrow 0 \text{ as } n_B \rightarrow \infty, \quad (27)$$

and that the constants  $c_h = c_h^{(n_B)}$  satisfy the following:

$$\begin{aligned} \text{for each } n_B = 4, 5, 6, \dots, \quad 1 \geq c_1^{(n_B)} \geq c_2^{(n_B)} \geq \dots \geq c_M^{(n_B)} = 0, \\ \text{and for fixed } h, \quad c_h^{(n_B)} \rightarrow 1 \text{ as } n_B \rightarrow \infty. \end{aligned} \quad (28)$$

The choice of constants  $M$  and  $c_1, c_2, \dots, c_M$  is discussed at the end of this section. We will show under (27) and (28) that

$$E|\hat{\gamma}^2(t_1, t_2) - \gamma^2(t_1, t_2)| \rightarrow 0. \quad (29)$$

This will imply that  $\hat{\gamma}^2(t_1, t_2)$  converges to  $\gamma^2(t_1, t_2)$  in probability.

Since  $E\hat{\nu}_h$  is not in general equal to  $\nu_h$ , it is more convenient to first work with

$$\tilde{\nu}_h = \frac{1}{n_B - h} \sum_{i=0}^{n_B - h - 1} (U_i - U_{AB}(t_1, t_2))(U_{i+h} - U_{AB}(t_1, t_2)) \quad (30)$$

and

$$\tilde{\gamma}^2(t_1, t_2) = \tilde{\nu}_0 + 2 \sum_{h=1}^M c_h \tilde{\nu}_h. \quad (31)$$

Since  $E\tilde{\nu}_h = \nu_h$ , if we define

$$\gamma_{n_B}^2(t_1, t_2) = \nu_0 + 2 \sum_{h=1}^M c_h \nu_h, \quad (32)$$

we have

$$E\tilde{\gamma}^2(t_1, t_2) = \gamma_{n_B}^2(t_1, t_2). \quad (33)$$

From now on we drop the arguments  $t_1$  and  $t_2$  whenever convenient. The triangle inequality gives

$$E|\hat{\gamma}^2 - \gamma^2| \leq E|\hat{\gamma}^2 - \tilde{\gamma}^2| + E|\tilde{\gamma}^2 - \gamma_{n_B}^2| + |\gamma_{n_B}^2 - \gamma^2|. \quad (34)$$

Thus, our objective is to show that each of the three terms on the right side of (34) converges to 0 as  $n_B \rightarrow \infty$ . It is easy to see that under (27) and (28),  $|\gamma_{n_B}^2 - \gamma^2| \rightarrow 0$  as  $n_B \rightarrow \infty$ .

We now consider  $E|\tilde{\gamma}^2 - \gamma_{n_B}^2|$ . Let  $c_0 = \frac{1}{2}$ . We have

$$\begin{aligned} E|\tilde{\gamma}^2 - \gamma_{n_B}^2| &= E|2 \sum_{h=0}^M c_h(\tilde{\nu}_h - \nu_h)| \leq 2 \sum_{h=0}^M E|\tilde{\nu}_h - \nu_h| \\ &\leq 2 \sum_{h=0}^M \left\{ E(\tilde{\nu}_h - \nu_h)^2 \right\}^{\frac{1}{2}} = 2 \sum_{h=0}^M \{ \text{Var } \tilde{\nu}_h \}^{\frac{1}{2}}. \end{aligned} \quad (35)$$

We now examine the variance terms. It is well-known (and easy to see) that if  $\{Y_i\}$  is a stationary sequence, then

$$\text{Var} \left( \frac{1}{n} \sum_{i=1}^n Y_i \right) = \frac{1}{n} \sum_{|\ell| < n} \left( 1 - \frac{|\ell|}{n} \right) \text{Cov}(Y_i, Y_{i+\ell}). \quad (36)$$

Equation (30) shows that  $\tilde{\nu}_h$  is of the form  $\tilde{\nu}_h = \frac{1}{n} \sum_{i=1}^n Y_i$ , where  $Y_i = (U_i - \bar{U})(U_{i+h} - \bar{U})$ , with  $\{Y_i\}$  stationary. We can thus obtain the exact variance of  $\tilde{\nu}_h$ :

$$\text{Var } \tilde{\nu}_h = \frac{1}{n_B - h} \sum_{|\ell| < n_B - h} \left( 1 - \frac{|\ell|}{n_B - h} \right) \sigma_h(\ell), \quad (37)$$

where

$$\sigma_h(\ell) = \text{Cov}((U_i - U_{AD})(U_{i+h} - U_{AD}), (U_{i+\ell} - U_{AD})(U_{i+h+\ell} - U_{AD})). \quad (38)$$

To obtain a useful bound on  $\text{Var } \tilde{\nu}_h$  we will show that

$$|\sigma_h(\ell)| \leq \left[ \mu((|\ell| - h)_+) \right]^r C, \quad (39)$$

where  $a_+ = \max\{a, 0\}$  and  $C$  is a constant not depending on  $h$  or  $\ell$ . The key ingredient in the proof of (39) is the use of a lemma of Ibragimov (1962) that gives an explicit bound for the covariance of two random variables  $\xi$  and  $\eta$  satisfying  $\xi \in \sigma(\dots, U_{-1}, U_0)$  and  $\eta \in \sigma(U_k, U_{k+1}, \dots)$ , in terms of  $\mu(k)$  and certain moments of  $\xi$  and  $\eta$ .

For any  $\ell$  such that  $|\ell| > h$ , Lemma 1.3 of Ibragimov (1962) implies that

$$\sigma_h(\ell) \leq |\mu(|\ell| - h)|^r \left\{ 4 + 6 E|(U_{\ell} - U_{AB})(U_{\ell+h} - U_{AB})|^{2(1 + \frac{1}{1-r})} \right\}. \quad (40)$$

The Cauchy-Schwarz inequality implies that

$$E|(U_{\ell} - U_{AB})(U_{\ell+h} - U_{AB})|^{2(1 + \frac{1}{1-r})} \leq E|(U_{\ell} - U_{AB})|^{4(1 + \frac{1}{1-r})}. \quad (41)$$

By A3 the right side of (41) is a finite constant not depending on  $h$  or  $\ell$ . This proves (39) for  $|\ell| > h$ . For  $|\ell| \leq h$  the proof is even simpler, and is omitted. We now combine (37) and (39) to obtain

$$\begin{aligned} \text{Var } \hat{\nu}_h &\leq \frac{1}{n_B - h} \sum_{|\ell| < n_B - h} |\sigma_h(\ell)| \leq \frac{1}{n_B - h} \left( \sum_{|\ell| \leq h} |\sigma_h(\ell)| + \sum_{|\ell| > h} \sigma_h(\ell) \right) \\ &\leq \frac{1}{n_B - h} \left( (2h + 1)C + \sum_{j=1}^{\infty} \mu(j)^r C \right). \end{aligned} \quad (42)$$

Next, we use (23) and (27) to see that there exists an integer  $J$ , not depending on  $h$  or  $\ell$ , such that

$$\text{Var } \hat{\nu}_h \leq n_B^{-\frac{1}{2}} \quad \text{for all } n_B \geq J. \quad (43)$$

We now return to  $E|\hat{\gamma}^2 - \gamma_{n_B}^2|$  (see (35)). Combining (27) and (43) we arrive at

$$E|\hat{\gamma}^2 - \gamma_{n_B}^2| \rightarrow 0 \quad \text{as } n_B \rightarrow \infty. \quad (44)$$

To complete the proof of (29) we need only show that  $E|\hat{\gamma}^2 - \gamma^2| \rightarrow 0$  as  $n_B \rightarrow \infty$ . We have

$$E|\hat{\gamma}^2 - \gamma^2| \leq E\left| 2 \sum_{h=0}^M c_h(\hat{\nu}_h - \nu_h) \right| \leq 2 \sum_{h=0}^M E|\hat{\nu}_h - \nu_h|. \quad (45)$$

We will presently show that

$$E|\hat{\nu}_h - \nu_h| \leq \frac{C'}{n_B}, \quad (46)$$

where  $C'$  is a constant not depending on  $h$ .

Referring to equations (26) and (30), it is easy to see that

$$\hat{\nu}_h = \hat{\nu}_h + (\bar{U}^{(1)} - U_{AB})(U_{AB} - U) + (U^{(2)} - U_{AB})(U_{AB} - U) + (U - U_{AB})^2. \quad (47)$$

where

$$\bar{U}^{(1)} = \frac{1}{n_B - h} \sum_{i=0}^{n_B - h - 1} U_i, \quad \bar{U}^{(2)} = \frac{1}{n_B - h} \sum_{i=h}^{n_B - 1} U_i.$$

Thus,  $E|\bar{U}_h - \bar{U}| \leq E|(\bar{U}^{(1)} - U_{AB})(U_{AB} - \bar{U})| + E|(\bar{U}^{(2)} - U_{AB})(U_{AB} - \bar{U})| + E(\bar{U} - U_{AB})^2$ .

Consider first  $E(\bar{U} - U_{AB})^2$ . From (36) we have

$$E(\bar{U} - U_{AB})^2 = \frac{1}{n_B} \sum_{|\ell| < n_B} (1 - \frac{|\ell|}{n_B}) \nu_\ell \leq \frac{C''}{n_B},$$

where  $C'' = \nu_0 + 2 \sum_{\ell=1}^{\infty} |\nu_\ell| < \infty$ . Similarly,

$$E(\bar{U}^{(1)} - U_{AB})^2 \leq \frac{C''}{n_B - h}, \quad \text{and} \quad E(\bar{U}^{(2)} - U_{AB})^2 \leq \frac{C''}{n_B - h}.$$

The Cauchy-Schwarz inequality now yields (46), which together with (27) imply that  $E|\hat{\gamma}_h - \tilde{\gamma}_h| \rightarrow 0$  as  $n_B \rightarrow \infty$ . This completes the proof of (i).

We next prove (iii), since the proof of (ii) proceeds in a very similar way.

PROOF OF (iii). For  $\xi_1, \xi_2, \xi_3 \in \mathbb{R}$ , let  $X_i = \xi_1 U_i + \xi_2 N_A(B_i, B_{i+1}) + \xi_3 (B_{i+1} - B_i)$ . The sequence  $\{X_i\}_{i=-\infty}^{\infty}$  is stationary, and if  $\pi(\cdot)$  denotes its mixing coefficient, it is clear that (23) holds for  $\pi(\cdot)$  as well. This gives a central limit theorem for  $\{X_i\}$ . It is simple to argue that  $E N_A(B_0, B_1) = \lambda_A/\lambda_B$  and that  $E(B_1 - B_0) = 1/\lambda_B$ . By the Cramér-Wold device we now have that

$$\sqrt{n_B} \left( \frac{\sum_{i=1}^{n_B} U_i}{n_B} - U_{AB}, \frac{n_A}{n_B} - \frac{\lambda_A}{\lambda_B}, \frac{T}{n_B} - \frac{1}{\lambda_B} \right)' \quad (48)$$

is asymptotically normal with mean 0 and covariance matrix, say  $\Psi$ . To identify  $\Psi$  and describe consistent estimates of it, it is convenient to introduce additional notation. Let

$$V_i^{(1)} = U_i, \quad V_i^{(2)} = N_A(B_{i+1}, B_i), \quad V_i^{(3)} = B_{i+1} - B_i \quad \text{for } i = \dots - 2, -1, 0, 1, 2, \dots \quad (49)$$

Let

$$\psi_h^{(pq)} = \text{Cov}(V_0^{(p)}, V_h^{(q)}) \quad \text{for } p, q = 1, 2, 3, \quad h = \dots - 2, -1, 0, 1, 2, \dots \quad (50)$$

Note that  $\psi_h^{(11)} = \nu_h$ . It is clear that the asymptotic covariance matrix of (48) is equal to the matrix whose  $pq$ th entry is  $\Psi^{(pq)} = \sum_{h=-\infty}^{\infty} \psi_h^{(pq)}$  (note that  $\psi_h^{(pq)}$  is not necessarily equal to  $\psi_{-h}^{(qp)}$ , unless  $p = q$ ). Next, for  $p, q = 1, 2, 3$ , define

$$\hat{\psi}_h^{(pq)} = \begin{cases} \frac{1}{n_B - h} \sum_{i=0}^{n_B - h - 1} (V_i^{(p)} - \bar{V}^{(p)})(V_{i+h}^{(q)} - \bar{V}^{(q)}) & \text{for } h = 0, 1, 2, \dots, M \\ \frac{1}{n_B - h} \sum_{i=h}^{n_B - 1} (V_i^{(p)} - \bar{V}^{(p)})(V_{i+h}^{(q)} - \bar{V}^{(q)}) & \text{for } h = -1, -2, -3, \dots, -M \end{cases} \quad (51)$$

where

$$\bar{V}^{(p)} = \frac{1}{n_B} \sum_{i=0}^{n_B-1} V_i^{(p)}. \quad (52)$$

Let

$$\hat{\Psi}^{(pq)} = \sum_{h=-M}^M c_{|h|} \hat{\psi}_h^{(pq)} \quad \text{for } p, q = 1, 2, 3, \quad (53)$$

where  $M$  and  $\{c_h\}$  satisfy (27) and (28), respectively. Also, let

$$\hat{\Psi} = \text{matrix whose } pq\text{th entry is } \hat{\Psi}_h^{(pq)}. \quad (54)$$

The same argument that was used in the proof of (i) now applies and we see that  $\hat{\Psi}$  converges to  $\Psi$  componentwise in probability as  $n_B \rightarrow \infty$ .

Consider now the function  $g(x, y, z) = (x, y/z, 1/z)$ , which maps  $(\sum_{i=1}^{n_B} U_i/n_B, n_A/n_B, T/n_B)$  into  $(\sum_{i=1}^{n_B} U_i/n_B, \hat{\lambda}_A, \hat{\lambda}_B)$ . The derivative of  $g$  evaluated at  $(x, y, z)$  is

$$Dg(x, y, z) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{z} & \frac{-y}{z^2} \\ 0 & 0 & \frac{-1}{z^2} \end{pmatrix}. \quad (55)$$

An application of the delta method (with the function  $g$ ) to the vector (48) gives the asymptotic normality result asserted in Part (iii) of the theorem, with

$$\sum(t_1, t_2) = \sum = Dg(U_{AB}, \lambda_A/\lambda_B, 1/\lambda_B) \Psi [Dg(U_{AB}, \lambda_A/\lambda_B, 1/\lambda_B)]'.$$

Defining  $\widehat{\sum} = \widehat{\sum}(t_1, t_2)$  by

$$\widehat{\sum} = Dg(\hat{U}_{AB}, \hat{\lambda}_A/\hat{\lambda}_B, 1/\hat{\lambda}_B) \hat{\Psi} [Dg(\hat{U}_{AB}, \hat{\lambda}_A/\hat{\lambda}_B, 1/\hat{\lambda}_B)]', \quad (56)$$

it is clear that under (27) and (28)  $\widehat{\sum}$  converges to  $\sum$  componentwise in probability as  $n_B \rightarrow \infty$ .  
**PROOF of ii.** As was mentioned above, the proof of (ii) is very similar to that of (iii). We now consider the variance estimates. Let  $\sum^{(pq)}$  and  $\widehat{\sum}^{(pq)}$  denote the  $pq$ th entries of  $\sum$  and  $\widehat{\sum}$ , respectively, and let

$$\hat{\Lambda} = \begin{pmatrix} \widehat{\sum}^{(22)} & \widehat{\sum}^{(23)} \\ \widehat{\sum}^{(32)} & \widehat{\sum}^{(33)} \end{pmatrix}, \quad (57)$$

with a similar definition for  $\Lambda$ . Obviously, this is the same  $\Lambda$  that appears in the statement of Part (ii) of Theorem 1. It is clear that under (27) and (28), as  $n_B \rightarrow \infty$ ,  $\hat{\Lambda} \rightarrow \Lambda$  componentwise in probability. This completes the proof of Theorem 1. ■

**THEOREM 2.** Let  $K(t_1, t_2)$  and  $\hat{K}(t_1, t_2)$  be defined by (3) and (5), respectively, and assume A1-A5. Then, as  $n_B \rightarrow \infty$

$$\sqrt{n_B}(\hat{K}(t_1, t_2) - K(t_1, t_2)) \xrightarrow{d} \mathcal{N}(0, \sigma^2(t_1, t_2)).$$

Furthermore, any estimate  $\hat{\sigma}^2(t_1, t_2)$  of the form (58) (refer to equations (49) and (51)-(54)), satisfying (27) and (28), is a consistent estimate of  $\sigma^2(t_1, t_2)$ .

PROOF. We apply the delta method to the vector (48) with the function  $f(x, y, z) = \frac{xz}{y}$ . The derivative of  $f$  at the point  $(x, y, z)$  is  $Df(x, y, z) = (\frac{z}{y}, \frac{-xz}{y^2}, \frac{x}{y})$ . Evaluated at  $(U_{AB}, \lambda_A/\lambda_B, \frac{1}{\lambda_B})$  and  $(\hat{U}_{AB}, \hat{\lambda}_A/\hat{\lambda}_B, \frac{1}{\hat{\lambda}_B})$ , this is  $(\frac{1}{\lambda_A}, K \frac{\lambda_B}{\lambda_A}, K \lambda_B)$  and  $(\frac{1}{\hat{\lambda}_A}, \hat{K} \frac{\hat{\lambda}_B}{\hat{\lambda}_A}, \hat{K} \hat{\lambda}_B)$ , respectively. The asymptotic normality asserted in the theorem follows from the asymptotic normality of the vector (48), with

$$\sigma^2(t_1, t_2) = (\frac{1}{\lambda_A}, K \frac{\lambda_B}{\lambda_A}, K \lambda_B) \Psi (\frac{1}{\lambda_A}, K \frac{\lambda_B}{\lambda_A}, K \lambda_B)'$$

and it is clear that under (27) and (28), if  $\hat{\sigma}^2(t_1, t_2)$  is defined by

$$\hat{\sigma}^2(t_1, t_2) = (\frac{1}{\hat{\lambda}_A}, \hat{K} \frac{\hat{\lambda}_B}{\hat{\lambda}_A}, \hat{K} \hat{\lambda}_B) \hat{\Psi} (\frac{1}{\hat{\lambda}_A}, \hat{K} \frac{\hat{\lambda}_B}{\hat{\lambda}_A}, \hat{K} \hat{\lambda}_B)' \quad (58)$$

then as  $n_B \rightarrow \infty$ ,  $\hat{\sigma}^2(t_1, t_2)$  converges to  $\sigma^2(t_1, t_2)$  in probability. This completes the proof of Theorem 2. ■

Results giving the asymptotic normality of estimates of  $\lambda_A$  and  $\lambda_B$  (under varying sets of assumptions) already exist in the literature; see e.g. Theorem 8.6 of Daley and Vere-Jones (1972). It was necessary to establish joint asymptotic normality of  $\hat{\lambda}_A$  and  $\hat{U}_{AB}(t_1, t_2)$  in order to obtain asymptotic normality of  $\hat{K}(t_1, t_2)$ .

We now discuss the choice of the constants  $M$  and  $c_1, c_2, \dots, c_M$ , which enter into the estimates  $\hat{\gamma}^2$  and  $\hat{\Psi}$  given by (25) and (53), respectively, and for the sake of simplicity, our discussion is in terms of  $\hat{\gamma}^2$  only. It is appropriate to discuss the choice of these constants within the framework of spectral density estimation. Defining

$$f(\omega) = \frac{1}{2\pi} \nu_0 + \frac{1}{\pi} \sum_{h=1}^{\infty} \nu_h \cos \omega h,$$

we see that  $\gamma^2 = 2\pi f(0)$ . To estimate  $f(\omega)$  we must effectively estimate  $\nu_h$  for each  $h$ . For fixed  $n_B$ ,  $\nu_h$  may be estimated for  $h = 0, 1, \dots, n_B - 1$ . However, because of the fact that for fixed  $n_B$

the variance of  $\hat{\nu}_h$  increases with  $h$ , it is standard to consider estimates of the form

$$\hat{f}(\omega) = \frac{1}{2\pi} \hat{\nu}_0 + \frac{1}{\pi} \sum_{h=1}^M c_h \hat{\nu}_h \cos \omega h$$

where  $M$  is much smaller than  $n_B$ , and  $c_h$  decreases as  $h$  increases. For a given value of  $M$  the constant  $c_h$  are usually given by  $c_h = w(\frac{h}{m})$  for some function  $w$  (called the "lag window") defined on  $[0, 1]$ , satisfying the following:  $w(0) = 1, w(1) = 0$ , and  $w$  decreases smoothly. Two commonly used choices are the Blackman-Tukey and the Parzen windows; see pp 514-516 of Anderson (1971) for a definition of these. Also, see Chapter 9 of Anderson (1971) for a general discussion of estimation of the spectral density. It is clear that as  $M$  increases the bias of  $\hat{f}(\omega)$  decreases while its variance increases. For both the Blackman-Tukey and the Parzen windows as well as for most of the commonly used windows, a value of  $M$  of the order  $n_B^{\frac{1}{2}}$  is usually used, since (under certain conditions on the stationary series) this minimizes the asymptotic mean squared error. See Section 9.3.4 of Anderson (1971). Thus, condition (27) is not at all restrictive.

It should be noted that the part of the proof of Theorem 1 that gives the consistency of  $\hat{f}(0)$  applies equally well to the estimates  $\hat{f}(\omega)$  for any  $\omega$ , and similarly for the spectral density estimates of the series  $\{N_A(B_{i+1}, B_i)\}$  and  $\{B_{i+1} - B_i\}$ .

It was necessary to give a proof of the consistency of  $\hat{f}(0)$  because the currently available consistency results for spectral density estimates are valid under conditions on  $\{U_i\}$  that are not implied by Assumptions A1- A5 (e.g. existence of all moments in Brillinger, 1975;  $\{U_i\}$  is a linear process as in Anderson, 1971, and in Hannan, 1970).

### 3. DISCUSSION.

The methods described in this paper enable the construction of asymptotic confidence intervals for  $K(t_1, t_2)$ , for fixed values of  $t_1$  and  $t_2$ . The function  $K(\cdot, \cdot)$  will usually be of interest over a continuum of values, say  $-L \leq t_1 < t_2 \leq L$ , where  $L$  is some number much smaller than  $T$ . One can plot  $\hat{K}(-L, t)$  for  $-L \leq t \leq L$  or, what is sometimes more useful, plot  $\hat{K}(t - \frac{d}{2}, t + \frac{d}{2})$  for  $-L + \frac{d}{2} \leq t \leq L - \frac{d}{2}$ . Here,  $d$  is some small number representing the experimenter's guess at the duration or likely duration of the effect of a  $B$  point on the  $A$  process. The function  $K(t - \frac{d}{2}, t + \frac{d}{2})$  is identically equal to  $d$  if  $N_A$  and  $N_B$  are independent.

We may form the bands

$$\hat{K}(-L, t) \pm z^{(\alpha/2)} \hat{\sigma}(-L, t) / \sqrt{n_B} \quad -L \leq t \leq L$$

and

$$\hat{K}(t - \frac{d}{2}, t + \frac{d}{2}) \pm z^{(\alpha/2)} \hat{\sigma}(t - \frac{d}{2}, t + \frac{d}{2}) / \sqrt{n_B} \quad -L + \frac{d}{2} \leq t \leq L - \frac{d}{2}$$

where  $\hat{\sigma}(t_1, t_2)$  is an estimate of  $\sigma(t_1, t_2)$  and  $z^{(\alpha/2)}$  is the upper  $\frac{\alpha}{2} \bullet 100$  percentile point of a standard normal variable. These bands of course are not simultaneous confidence bands. To form simultaneous confidence bands one would need to carry out two distinct steps:

(i) Establish weak convergence of the processes

$$V_{n_B}(t) = \sqrt{n_B} (\hat{K}(-L, t) - K(-L, t))$$

and

$$W_{n_B}(t) = \sqrt{n_B} (\hat{K}(t - \frac{d}{2}, t + \frac{d}{2}) - K(t - \frac{d}{2}, t + \frac{d}{2}))$$

to Gaussian processes  $V(t)$  and  $W(t)$ , respectively.

(ii) Obtain  $v^{(\alpha)}$  and  $w^{(\alpha)}$ , the upper  $\alpha \bullet 100$  percentile points of  $\sup_{-L \leq t \leq L} |V(t)|$  and  $\sup_{-L + \frac{d}{2} \leq t \leq L - \frac{d}{2}} |W(t)|$ , respectively.

The bands

$$\hat{K}(-L, t) \pm v^{(\alpha)} / \sqrt{n_B} \quad -L \leq t \leq L$$

and

$$\hat{K}(t - \frac{d}{2}, t + \frac{d}{2}) \pm w^{(\alpha)} / \sqrt{n_B} \quad -L + \frac{d}{2} \leq t \leq L - \frac{d}{2}$$

are then asymptotic simultaneous confidence bands.

A proof of weak convergence appears extremely difficult. Although desirable from a theoretical point of view, weak convergence is not useful statistically unless the distribution of the supremum of the absolute value of the limiting process can be obtained. Unfortunately, this is in general a very difficult problem even if the Gaussian process is stationary (see Cressie and Davis, 1981).

## REFERENCES

- Brillinger, D. R. (1975). Time Series, Data Analysis and Theory. Holt, New York.
- Brillinger, D. R. (1976). Estimation of the second-order intensities of a bivariate stationary point process. J. Roy. Statist. Soc. Ser. B **38**, 60-66.
- Bryant, H. L., Ruiz Marcos, A., and Segundo, J. P. (1973). Correlations of neuronal spike discharges produced by monosynaptic connections and of common inputs. J. Neurophysiol. **36**, 205-225.
- Cox, D. R. and Lewis, P. A. W. (1972). Multivariate point processes. Proc. 6th Berkeley Symp. **2**, 401-448.
- Cressie, N. and Davis, R. W. (1981). The supremum distribution of another Gaussian process. J. Appl. Prob. **18**, 131-138.
- Daley, D. J. and Vere-Jones, D. (1972). A summary of the theory of point processes. In Stochastic Point Processes: Statistical Analysis, Theory and Applications, (P. A. W. Lewis, ed.) 299-383. Wiley, N.Y.
- Diggle, P. J., and Milne, R. K. (1983). Bivariate Cox processes: some models for bivariate spatial point patterns. J. Roy. Statist. Soc. Ser. B **45**, 11-21.
- Hannan, E. J. (1970). Multiple Time Series. Wiley, New York.
- Ibragimov, I. A. (1962). Some limit theorems for stationary processes. Theory Prob. and Applic. **7**, 349-382.
- Karlin, S. and Taylor, H. (1975). A First Course in Stochastic Processes, 2nd ed. Academic Press, New York.
- Khintchine, A. Ya. (1960). Mathematical Methods in the theory of Queueing. Griffin, London.
- Leadbetter, M. R. (1968). On three basic results in the theory of stationary point processes. Proc. Amer. Math. Soc. **19**, 115-117.
- Leadbetter, M. R. (1972). On basic results of point process theory. Proc. 6th Berkeley Symp. **3** 449-462.
- Lotwick, H. W., and Silverman, B. W. (1982). Methods for analysing spatial processes, of several types of points. J. Roy. Statist. Soc. Ser. B **44**, 406-413.
- Ripley, B. D. (1976). The second-order analysis of stationary point processes. J. Appl. Prob. **13**, 255-266.

- Ripley, B. D. (1977). Modelling spatial patterns. (With discussion.) J. Roy. Statist. Soc. Ser. B 39, 172-212.
- Ripley, B. D. (1981). Spatial Statistics. Wiley, New York.
- Silverman, B. W. (1976). Limit theorems for dissociated random variables. Adv. Appl. Prob. 8, 806-819.
- Wisniewski, T. K. M. (1972). Bivariate stationary point processes, fundamental relations and first recurrence times. Adv. Appl. Prob. 4, 296-317.

END

JAN.

1988

DTIC