

AD-A197 254 RECENT PROGRESS IN THE P AND H-P VERSION OF THE FINITE ELEMENT METHOD(U) MARYLAND UNIV COLLEGE PARK INST FOR PHYSICAL SCIENCE AND TECHNOLOGY I BABUSKA JUL 87 1/1

AD-A197 254 RECENT PROGRESS IN THE P AND H-P VERSION OF THE FINITE ELEMENT METHOD(U) MARYLAND UNIV COLLEGE PARK INST FOR PHYSICAL SCIENCE AND TECHNOLOGY I BABUSKA JUL 87 1/1

AD-A197 254 RECENT PROGRESS IN THE P AND H-P VERSION OF THE FINITE ELEMENT METHOD(U) MARYLAND UNIV COLLEGE PARK INST FOR PHYSICAL SCIENCE AND TECHNOLOGY I BABUSKA JUL 87 1/1

UNCLASSIFIED BN-1067 N00014-85-K-0169 F/G 12/1 NL

UNCLASSIFIED BN-1067 N00014-85-K-0169 F/G 12/1 NL

UNCLASSIFIED BN-1067 N00014-85-K-0169 F/G 12/1 NL

UNCLASSIFIED BN-1067 N00014-85-K-0169 F/G 12/1 NL

[illegible]

1	2	3	4	5	6	7	8	9	10
1	2	3	4	5	6	7	8	9	10



DTIC FILE COPY



COLLEGE PARK CAMPUS




RECENT PROGRESS IN THE p AND h-p VERSION
OF THE FINITE ELEMENT METHOD

I. Babuška
Institute for Physical Science and Technology
University of Maryland, College Park, MD 20742

BN-1067

July 1987



INSTITUTE FOR PHYSICAL SCIENCE
AND TECHNOLOGY

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER BN-1067	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Recent Progress in the p and h-p Versions of the Finite Element Method		5. TYPE OF REPORT & PERIOD COVERED Final life of the contract
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) I. Babuska		8. CONTRACT OR GRANT NUMBER(s) ONR N00014-85-K-0169
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Maryland Institute for Physical Science and Technology College Park, MD 20742		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Department of the Navy Office of Naval Research Arlington, VA 22217		12. REPORT DATE July 1987
		13. NUMBER OF PAGES 17
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release: distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The paper gives the state of the art and literature related to the development of the p and h-p version of the finite element method.		

DD FORM 1473

EDITION OF 1 NOV 65 IS OBSOLETE

S/N 0102-LF-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

by
I. Babuška
Institute for Physical Science and Technology
University of Maryland, College Park, MD 20742

1. INTRODUCTION

The finite element method has become the main tool in computational mechanics. The success is manifested by the development of over five hundred user-oriented finite element program systems, ~~see e.g. [33]~~. The literature on the subject is overwhelming. To date there are over two hundred monographs and conference proceedings ^[40] and new monographs and proceedings are continuously appearing. Various forms of the finite element method are used in practice for the numerical treatment of elliptic, parabolic, hyperbolic, linear and nonlinear partial differential equations, integral and integrodifferential equations, etc. Any class of problems has its own specific features. ~~In this paper we will only deal~~ with the class of partial differential equation of elliptic type. For the sake of simplicity ~~we will~~ ^{the author} elaborate on a characteristic model problem and illustrative results and make ^{only} additional comments of more general nature although the results ~~we refer~~ ^{referred} to are general.

2. THE MODEL PROBLEM AND ITS PROPERTIES

We restrict ourself to the most simple model problem. Let $\Omega \subset \mathbb{R}^2$, $x = (x_1, x_2)$ be a bounded (simple connected) domain with the boundary $\partial\Omega = \Gamma$ consisting of simple arcs

$$\Gamma_1 = \{x_1 = f_{1,1}(t), x_2 = f_{1,2}(t), t \in I\}$$

where $I = (-1, 1)$ and $\Gamma = \bigcup_{i=1}^M \bar{\Gamma}_i$. We will assume that $f_{1,1}$ and $f_{1,2}$ are analytic functions on $\bar{I}$ and $f_{1,1}^2 + f_{1,2}^2 > \alpha > 0$. The vertices of Ω will be denoted by A_i , $i = 1, \dots, M$ and the internal angles at A_i by ω_i .

Let us be interested in the model problem and its standard (weak solution):

$$-\Delta u = f \text{ on } \Omega$$

(2.1)



A-1

$$u = \varphi \text{ on } r^0 = \bigcup_{j \in Q} \bar{\Gamma}_j \quad (2.2a)$$

$$\frac{\partial u}{\partial n} = \psi \text{ on } r^1 = r - r^0 \quad (2.2b)$$

Here Q is a subset of $\{1, \dots, M\} = M$. For simplicity we assume that $r^0 \neq \emptyset$.

The performance of any numerical method strongly depends on the properties of the (exact) solution of the solved problem especially on its smoothness. The more information is available, the better method could be designed.

It is very advantageous to characterize the set of solutions of (2.1), (2.2) under consideration in the terms of countably normed spaces.

Let $A_i = (x_{1,i}, x_{2,i})$ and $r_i^2 = (x_1 - x_{1,i})^2 + (x_2 - x_{2,i})^2$. Define $\phi_\beta(x) = \prod_{i=1}^M r_i^{\beta_i}(x)$, $\beta = (\beta_1, \dots, \beta_M)$, $0 < \beta_i < 1$, and for any integer k let $\phi_{\beta \pm k}(x) = \prod_{i=1}^M r_i^{\beta_i \pm k}(x)$. Then we let $B_\beta^2(\Omega) = \{u \in H^1(\Omega) |$

$$|\phi_{\beta+k-2} D^\alpha u|_{L_2(\Omega)} \leq C d^k k!, \quad k = 2, 3, \dots, |\alpha| = k, C, d \text{ independent of } k\}.$$

If $u \in B_\beta^2(\Omega)$ then it is analytic on $\bar{\Omega} = \bigcup_{i=1}^M A_i$ and has specific behavior

in the neighborhood of A_i , $i = 1, \dots, M$. In [7] [8] we have proven

Theorem 2.1. Let f be analytic on $\bar{\Omega}$, φ be analytic on $\bar{\Gamma}_j$, $j \in Q$ and continuous $\bar{\Gamma}^0$, ψ be analytic on Γ_j , $j \in M - Q$. Then for (2.1) and (2.2) $u \in B_\beta^2(\Omega)$ with $\beta_i > \bar{\beta}_i$; $\bar{\beta}_i$ depends on ω_i and the type of boundary condition on Γ_{i-1} and Γ_i . □

If, for example, $M = 5$, $Q = \{1, 2, 3\}$ we get $\bar{\beta}_1 = \max(0, 1 - \frac{\pi}{4\omega_1})$, $\bar{\beta}_2 = \max(0, 1 - \frac{\pi}{2\omega_2})$, $\bar{\beta}_3 = \max(0, 1 - \frac{\pi}{2\omega_3})$, $\bar{\beta}_4 = \max(0, 1 - \frac{\pi}{4\omega_4})$, $\bar{\beta}_5 = \max(0, 1 - \frac{\pi}{2\omega_5})$.

Remark 2.1. In [9] we precisely characterized the traces of functions from $B_\beta^2(\Omega)$ and gave full characterization of the sets of f , φ and ψ which guarantee that the solution u of (2.1) (2.2) belongs to B_β^2 .

Remark 2.2. Theorem 2.1 also holds when the differential equation in (2.1)

has analytic coefficients on $\bar{\Omega}$ (see [7]).

Remark 2.3. The eigenfunctions of the eigenvalue problem (2.1) (2.2) also belong to $B^2_{\beta}(\Omega)$ (see [10]).

Remark 2.4. Theorem 2.1 also holds for strongly elliptic system of differential equations as elasticity equations (see [11]).

In practice, e.g. in the field of structural mechanics, the problems of partial differential equations are characterized by piecewise analytic data and hence theorem 2.1 is very well suited for the applications.

3. THE FINITE ELEMENT METHOD

There are various different forms of the finite element method. We will consider here only the basic class of finite element methods (for our model problem).

Let $T = \{\tau_i\}$ be a partition of Ω into (in general curvilinear) triangles or quadrilaterals called elements τ_i . In the case when T is a triangulation we are making the standard assumptions. For the general case we refer to [8], [9], [36]. We will formulate here the assertions in the case of triangulation only although they hold in general. Let $H(p, T) = \{u \in H^1(\Omega) \mid u|_{\tau_i}, \tau_i \in T \text{ is a polynomial of degree } p\}$ be the finite element space. If τ_i is a rectangle, then polynomials are of degree p in both variables. If the elements are curvilinear, then $u|_{\tau_i}$ are the standard "pull-back" polynomials.

Remark 3.1. We have assumed that the degree of polynomials are the same over all elements. The theory is developed for general case when the degree p can be different on different elements.

Let $H_0(p, T) = H(p, T) \cap H_0^1(\Omega)$ where $H_0^1(\Omega) = \{u \in H^1(\Omega) \mid u = 0 \text{ on } r^0\}$ and $H_0(p, T)$ be the restriction of $H(p, T)$ on r^0 . We will assume that a projection operator $P_0(p, T)$ of function φ into $H_0(p, T)$ be given and we denote $\varphi_{p, T} = P_0(p, T)\varphi$. A concrete possible form of $P_0(p, T)$ will be given later.

The finite element method consists now in finding $u_{FE} = u(p, T) \in H(p, T)$ so that

$$1) \quad u(p, T) = \varphi_{p, T} \quad \text{on } \Gamma^0$$

$$2) \quad \iint_{\Omega} \left(\frac{\partial u_{FE}}{\partial x_1} \frac{\partial v}{\partial x_1} + \frac{\partial u_{FE}}{\partial x_2} \frac{\partial v}{\partial x_2} \right) dx_1 dx_2$$

$$= \int_{\Gamma_1} \psi v \, ds + \iint_{\Omega} f v \, dx_1 dx_2$$

holds for any $v \in H_0(p, T)$.

We will be interested in the accuracy of the finite element solution measured in the energy norm. Define $e = u - u_{FE}$ and let

$$|e(T, p)|_E^2 = \iint_{\Omega} \left(\left(\frac{\partial e}{\partial x_1} \right)^2 + \left(\frac{\partial e}{\partial x_2} \right)^2 \right) dx_1 dx_2$$

be the error measured in the energy norm.

Two kind of operators $P_0(T, p)$ can be considered. Let $\gamma \subset \Gamma^0$ be the side of the triangle τ with endpoints A, B and assume that $\gamma = [-1, 1]$. $A = -1$, $B = 1$. Then $\varphi_{p, T}|_{\gamma} = l(x) + w(x)$ where $l(x)$ is linear function on γ such that $\varphi_{p, T}|_{\gamma}(\pm 1) = \varphi(\pm 1)$ and a) in the case of H^1 -

projection: $P_0^1(\tau, p)$: $\varphi' = \sum_{k=0}^{\infty} a_k l'_k$, $w'(x) = \sum_{k=0}^{p-1} a_k l'_k$, where l_k are the Legendre polynomials; b) in the case of $H^{1/2}$ -projection $P_0^{1/2}(T, p)$: $\varphi(x) = \sum_{k=0}^{\infty} a_n T_k(x)$, $w(x) = \sum_{k=0}^p a_k T_k(x)$, where $T_k(x)$ are the Tchebyshev polynomials.

As we have seen there is a large freedom in the selection of $H(p, T)$ namely the degree p and the partition T and in the selection of the operators $P_0(T, p)$. We expect that $|e|_E \rightarrow 0$ if $\dim H_0(p,) \rightarrow \infty$.

It is convenient to distinguish in this context three versions of the finite element method.

a) The h-version. Here a sequence (family) $H(p, T_1)$ is considered when p is fixed (usually $p = 1, 2$) and the mesh T_1 is successively refined so that the size h of the elements of T_1 goes to zero.

b) The p-version. Here the mesh T is kept fixed and $p \rightarrow \infty$ uniformly on selectively.

c) The h-p version. In this version the mesh is simultaneously

refined and the degree p increased uniformly or selectively.

The h-version of the finite element method is the standard one and extensive literature is devoted to it. The p-version is a recent development. The first theoretical paper about the p-version [26] and the h-p version [6] appeared in 1981 and various results were obtained since then. Some of them will be mentioned later.

There are many codes, research and commercial utilizing the h-version. The only commercial code using p and h-p versions is the code PROBE which was developed recently by NOETIC Technologies, St. Louis [54]. PROBE solves two dimensional problems of linear elasticity, stationary heat problems and thermoelasticity problems. The three dimensional extension of PROBE will be released in 1988. Three dimensional research code STRIPE was developed by Swedish Aeronautical Research Institute, see e.g. [1]. These codes have in addition various features as adaptive approaches, various a-posteriori error estimation, etc.

4. THE BASIC PROPERTIES OF THE p AND h-p VERSIONS OF THE FINITE ELEMENT METHOD

After 1980 the p and h-p versions of the finite element method was studied in detail from various point of view. We will mention here some essential illustrative results.

In one dimensional setting the versions were analyzed in detail in [34]. Here, among other, the optimal meshes and p-distribution has been established with upper and lower bounds of the errors for the three basic finite element versions.

In two dimensional setting the following theorem is characteristic for the performance of the h-p version. (For details see [8], [9], [36].)

Theorem 4.1. Let the solution u of the problems (2.1), (2.2) belongs to the set $B^2_{\beta}(\Omega)$. Then there is a sequence of meshes T_1 and the degrees p_1 such that

$$\|e\|_E < C e^{-\frac{3}{\alpha\sqrt{N_1}}}, \quad \alpha > 0 \quad (4.1)$$

where $N_1 = \dim H_0(p_1, T_1)$ is the number of degrees-of-freedom for the h-p version. □

In one dimension the rate is $Ce^{-\frac{2}{\alpha}\sqrt{N}}$. It has been proven in [34] that the optimal mesh is a geometric one with the factor $(\sqrt{2} - 1)^2 = .17$. The experience shows that the geometric mesh with the factor $= .15$ is also optimal in two dimension. Theorem 4.1 together with Theorem 2.1 shows that practically in any problem of structural mechanics the exponential rate of convergence can be achieved.

For the p-version the following theorem is another typical one (for more, see [21], [22], [23], [24]).

Theorem 4.2. Let us consider the problem (2.1) (2.2) and let $\bar{\beta} = \max(\bar{\beta}_1)$ given in Theorem 2.1. Then for the p-version we have

$$|e|_E \leq CN^{-(1-\bar{\beta})} \quad (4.2)$$

while for the h-version with uniformed mesh

$$|e|_E \geq CN^{-\left(\frac{1-\bar{\beta}}{2}\right)}. \quad (4.3)$$

□

In Theorems 4.1 and 4.2 either the projection P_0^1 or $P_0^{\frac{1}{2}}$ could be used provided φ, ψ are sufficiently smooth. The difference between these two operators occurs for the p-version when the boundary condition φ is unsmooth, e.g. $\varphi \in H^\delta(\Gamma)$, $\frac{1}{2} < \delta < 3/4$. In this case the $H^{\frac{1}{2}}$ projection has to be used. For the h-p version there is no difference in the asymptotic rate but some difference occurs in the constant of the estimates. For the analysis of the influence of the operator P_0 on the accuracy we refer to [12].

Remark 4.1. So far we have assumed that the domain Ω is bounded. Nevertheless the exponential rate of convergence (4.1) holds also for the problem on $\Omega^C = R^2 - \Omega$ when f has bounded support. Here the infinite elements and properly selected shape functions have to be used. For more, see [15].

Remark 4.2. The h, p and h-p versions have different aspects with respect to the pollution problem. In presence of a singular behaviour of the solution (e.g. in the neighborhood of the entrant corner of the domains)

the L_∞ error is very large in the element consisting the corner. This effect disappears in elements which are separated away from the singularity by few elements. This effect is essential for a proper mesh design in practical computation. For details we refer to [14].

So far we have dealt with the problem (2.1) and (2.2) of second order. For the analyses of the finite element solution for the problems of order $2k$ we refer in the case of the h-p version to [35] and the p-version to [39] [53]. For the basic analysis of the p-version in 3 dimensions, we refer to [30] and [31]. The eigenvalue problem is, as is well known, directly related to the "source" problem we addressed earlier. See e.g. [16] and [17]. In the case of the eigenvalue problem (in our case)

$$\Delta u = \lambda u$$

$$u = 0 \text{ on } \Gamma^0$$

$$\frac{\partial u}{\partial n} = 0 \text{ on } \Gamma^{(1)}.$$

The eigenfunctions belong to $B_B^2(\Omega)$ and hence

$$|\lambda - \lambda_{FE}| \leq C e^{-2\alpha\sqrt{N}^3}$$

$$\|u_{FE}(\lambda_h) - u(\lambda)\|_E \leq C e^{-\alpha\sqrt{N}^3}.$$

For more details, see e.g. [10].

5. IMPLEMENTATION AND COMPUTATIONAL COMPLEXITY

There are some essential features of implementation of the p and h-p versions. See e.g. [56]. The elements are of a hierarchical type which leads to augmentation (bordering) of the local stiffness matrices when p is increased. This also allows to change very flexibly the degree of the shape functions from one element to the other one. The shape elements are (in two dimensions) of nodal type, side type and internal type and are based (in PROBE) on the integrals for the Legendre polynomials. This is important for numerical stability aspects. The computation (in two dimension) of the local stiffness matrix on a rectangle requires (when optimally programmed) $O(p^4)$ operations. The computation of the local stiffness matrices takes much more effort for the p-version (with high p) than for the h-version and

hence the p-version is very well suited for parallel computations. The sparsity of the global stiffness matrix is also smaller for the p version than the h-version. Hence, the complexity of the computation for the same number of degrees-of-freedom is higher for the p-version than the h-version. Nevertheless, it is essential to relate the achieved accuracy to the computational effort. For an analysis of the computational complexity and computer time comparison, we refer to [19] and [20]. The results show that the h-p version with higher degree p is preferable for solutions which are not overly unsmooth or have singular behaviour in a-priori known areas as in the neighborhood of the corners. If the solution is very unsmooth, e.g. if the coefficients in the differential equations are uniformly rough, for example, measurable only, then only low accuracy is practically achievable by any method and h-version with low p is preferable. In general, for very low accuracies the low order elements are preferable, for the modest one higher degree elements have to be preferred.

6. THE PROBLEM OF THE MESH DESIGN

One of the most laborious part of the finite element analysis, especially in three dimensions, is the mesh generation also when sophisticated mesh generators are used. The use of large elements (possibly of high degrees) which are describing only the geometry, greatly simplifies the user's work, also if possibly on expense of the computer time. (It is necessary to realize that the relation between manpower cost and computation cost is going steadily up.) The option of a change of the degrees of elements increases significantly the flexibility of the program and gives the user effective tool for the quality control. The p and h-p version programs give such options. It is advantageous therefore to create directly or indirectly the proper mesh and to achieve then the desired accuracy by an appropriate choice of the degrees, which can be made, for example, in an adaptive mode. The goal is to achieve the same combination of the degrees and mesh refinement which would be obtained for the given accuracy by the h-p version directly. To achieve this goal two avenues could be followed, the expert system and the adaptive approach. The expert system, see e.g. [18], [45], advises the user how to design the mesh and element degrees for the requested accuracy and provides the user with a mesh generator. The expert system is interactive, follows the progress of computation, gives the user

on his request various desired information for an effective computation and engineering analysis. The experience (see [18] [19]) is that the cost of the expert activity for a mesh and degree design is at most 20% of the total cost.

The adaptive approach (see e.g. [46], [34], Part 3) which is possible to see as an "automatic" expert system makes various decisions for the users. Both approaches have some common parts but the concepts are significantly different. We refer also to [46] for various additional aspects.

7. THE ROBUSTNESS

An effective method has to perform uniformly well for a broad class of input data. The elasticity problems can be in practice nearly singular as, for example, in the case of nearly incompressible material, various plate and shell theories, in the case of thin domains, etc. The h-version suffers in these cases by the "locking" problem which has to be overcome by various special approaches as reduced integration, etc. (see e.g. [28]). Problems of these types are usually avoided when the p and h-p versions are used. The convergence rate then (in contrast to the h-versions) is uniform with respect to the Poisson ratio when higher degrees of elements are used. See e.g. [49], [50] [64].

8. THE QUALITY CONTROL OF THE SOLUTION

It is essential to have a possibility of a quantitative assessment of the quality of computed data. For the survey of today's general ideas and results in this direction, we refer to [41]. In the case of the p and h-p versions there is relatively easy way for the quality assessment of any data of the interest by changing the degrees and by an extrapolation procedure. This approach is very effective because it indicates reliably the errors of any computed data of interest, the energy norm, value of stresses, stress intensity factor, etc. See e.g. [1], [8], [36], [58], [61].

9. THE COMPUTATIONAL AND ENGINEERING EXPERIENCE

Because of the developed commercial code PROBE and research code STRIPE an extensive experience is already available in the research and industrial use. For the industrial experience we refer e.g. to [27] where numerical results are presented. For the research one we refer e.g. to [1], [3], [8],

[9], [10], [18], [19], [20], [25], [36], [48], [55], [57], [63]. The experience shows that the p and h-p versions of the finite element method has many practical advantages for the engineering computations for linear elliptic problems.

10. RELATION TO SOME OTHER METHODS

The ideas of the p and h-p versions are related for example to the various methods used in fluid dynamics as the spectral method and its variation. We refer e.g. to [43] and references given there. A commercial fluid dynamics code written by T. Patera is closely related to the idea of h-p version of the finite element method.

The h-p version of course generates various finite element approaches and in principle encompasses various diverse approaches, see e.g. [29]. The h-p version can be naturally also used for solving integral equations, boundary element method, etc. See e.g. [47], [51], [67].

11. SUMMARY

The p and h-p version of the finite element method is a new development which gives new possibilities, theoretical and practical for the finite element method. It is today reasonably well understood in the case of elliptic equations, both theoretically and practically. The aim of the present paper was to give a brief survey of various aspects of the p and h-p versions of the finite element method for solving elliptic linear problems and provide the comprehensive references. Nevertheless, many theoretical and practical aspects of the method for other problems, linear and nonlinear are still to be resolved as well as the problems of implementation for three dimensional problems.

In addition we refer to the references [2], [3], [4], [37], [38], [42], [44], [52], [58], [60], [62], [65], [66] which are directly related to the subject discussed.

REFERENCES

- [1] Andersson, B., Error estimation and adaptivity in FE-analysis of structures-Research program, Rep. FFAH-85-03. The Aeronautical Research Institute of Sweden, Structures Dept., Box 11021, 16111 Bromma, Sweden.
- [2] Andersson, B., Falk, U., Jarlas, R., Self-adaptive FE-analysis of solid structures, Rep. FFA TM 1986-27. The Aeronautical Research Institute of Sweden, Structures Dept., Box 11021, 16111 Bromma, Sweden.
- [3] Andersson, B., Falk, U., Finite element analysis of three-dimensional structures using adaptive p-extensions, Rep. FFA TN 1986-57. The Aeronautical Research Institute of Sweden, Structures Dept. Box 11021, 16111 Bromma, Sweden.
- [4] Babuška, I., The p and h-p versions of the finite element methods. The state of the Art. To appear in Proc. of the Workshop on Finite Element Method, Nasa Langley, ed. R. Voigt, Springer 1987 (Tech. Note BN-1056, 1986, Institute for Physical Science and Technology, University of Maryland, College Park, MD 20742).
- [5] Babuška, I., Are High Degree Elements Preferable? Some Aspects of the h and h-p versions of the Finite Element Method in Numerical Techniques for Engineering Analysis and Design. Proceedings of the International Conference on Numerical Methods in Engineering: Theory and Applications. Numeta 87, Swansea, 6-10 July 1987. G. N. Pande & J. Middleton eds. Martinus Nijhoff Publishers, Dordrecht/Boston/Lancaster, 1987, paper S1.
- [6] Babuška, I., Dorr, M. R., Error Estimates for the Combined h and p-Version of the Finite Element Method, Numer. Math. 37 (1981), 257-277.
- [7] Babuška, I., Guo. B., Regularity of the solution of elliptic problems with piecewise analytic data, Part I: Boundary value problem for linear elliptic equations of second order. To appear in SIAM J. Math. Anal., 1987 (Tech. Note BN-1047, 1986, Institute for Physical Science and Technology, University of Maryland, College Park, MD 20742).
- [8] Babuška, I., Guo, B., The h-p version of the Finite Element Method for Domains with Curved Boundaries. To appear in SIAM J. Numer. Anal., 1987 (Tech Note BN-1057, 1986, Institute for Physical Science and Technology, University of Maryland, College Park, MD 20742).

- [9] Babuška, I., Guo, B., The h-p version of the finite element method for problems with nonhomogeneous essential boundary conditions. To appear.
- [10] Babuška, I., Guo, B., Numerical treatment of the eigenvalue problem by the h-p version of the finite element method. To appear.
- [11] Babuška, I., Guo, B., Regularity of the solution of elliptic problems with piecewise analytic data. Part II. Boundary value problem for system of equations of second order. To appear.
- [12] Babuška, I., Guo, B., Suri, M., Implementation of nonhomogeneous Dirichlet boundary conditions in the p-version of finite element method. To appear.
- [13] Babuška, I., Gui, W., Szabó, B. A., Performance of the h-p and h, p and h-p versions of the finite element method in Research in Structures and Dynamics, (R. J. Hayduk, A. K. Noor, eds) NASA Conference Publication 2325 (1984), 73-94.
- [14] Babuška, I., Oh, Hae-Soo, Pollution problems for the p-version and the h-p version of finite element method. To appear in Communications in Applied Numerical Methods, 1987 (Tech. Note BN-1054, 1986, Institute for Physical Science and Technology, University of Maryland, College Park, MD 20742).
- [15] Babuška, I., Oh, Hae-Soo, The p and h-p versions of the finite element method for elliptic problems on infinite domains. To appear.
- [16] Babuška, I., Osborn, J., Estimates for the errors in eigenvalue and eigenvector approximation by Galerkin methods, with particular attention to the case of multiple eigenvalues. To appear in SIAM J. Num. Anal. 1987 (Tech. Note BN-1056, 1986, Institute for Physical Science and Technology, University of Maryland, College Park, MD 20742).
- [17] Babuška, I., Osborn, J., Eigenvalue Problems. To appear in Handbook of Numerical Analysis, ed. P. G. Ciarlet, J. L. Lions, North-Holland, Amsterdam [Tech Not BN-1066, 1987, Institute for Physical Science and Technology, University of Maryland, College Park, MD 20742).
- [18] Babuška, I., Rank, E., An expert-system-like feedback approach in the h-p version of the finite element method. To appear in Finite Elements in Analysis and Design, 1987 (Tech. Note BN-1048, 1986, Institute for Physical Science and Technology, University of Maryland,

College Park, MD 20742).

- [19] Babuška, I., Scapolla, T., Computational aspects of the h, p and h-p versions of the finite element method. Advances in Computer Methods in Partial Differential Equations-VI, R. Vichnevetsky and R. S. Stepleman eds. Int'l Association for Mathematics and Computer Simulation, 1987, 233-240.
- [20] Babuška, I., Scapolla, T., Benchmark Computation and Finite Element Performance Evaluation for a Rhombic Plate Problem. To appear.
- [21] Babuška, I., Suri, M., The optimal convergence rate of the p-version of the finite element method. To appear in SIAM J. Numer. Anal., 1987 (Tech Note BN-1045, 1986, Institute for Physical Science and Technology, University of Maryland, College Park, MD 20742).
- [22] Babuška, I., Suri, M., The h-p version of the finite element method with quasi uniform meshes. To appear in Math. Modeling Numer. Anal. (RAIRO), 1987 (Tech. Note BN-1046, 1986, Institute for Physical Science and Technology, University of Maryland, College Park, MD 20742).
- [23] Babuška, I., Suri, M., The treatment of nonhomogeneous Dirichlet boundary condition by the p-version of the finite element method, Tech. Note BN-1063, 1987, Institute for Physical Science and Technology, University of Maryland, College Park, MD 20742).
- [24] Babuška, I., Suri, M., The p-version of the finite method for constrained boundary conditions (Tech. Note BN-1064, 1987, Institute for Physical Science and Technology, University of Maryland, College Park, MD 20742).
- [25] Babuška, I., Szabó, B. A., Rates of convergence of the finite element method, Internat. J. Numer. Methods Engrg. 18 (1982), 323-341.
- [26] Babuška, I., Szabó B. A., Katz, I. N., The p-version of the finite element method. SIAM J. Numer. Anal. 18 (1981), 512-545.
- [27] Barnhart, M. A., Eisenmann, J. R., Analysis of a Stiffened Plate Detail Using p-Version and h-Version Finite Element Techniques. Presented at the First World Congress on Computational Mechanics, Sept. 22-26, 1986. The University of Texas at Austin.
- [28] Bathe, K. J., Brezzi, F., Studies of finite element procedures - the INF-SUP condition, equivalent forms and applications in Reliability of Methods for Engineering Analysis, J. J. Bathe, J. Owen, eds.

- Proceedings of the International Conference at Swansea, July 1986, Pineridge Press, Swansea, U. K., 1986, 197-220.
- [29] Delves, L. M., Phillips, C., A fast implementation of the global element method, J. Inst. Math. Appl. 25 (1980), 177-197.
 - [30] Dorr, M. R., The approximation theory for the p-version of the finite element method, SIAM J. Numer. Anal. 21 (1984), 1181-1.
 - [31] Dorr, M. R., The approximation of solutions of elliptic boundary value problems via the p-version of the finite element method, SIAM J. Numer. Anal. 23 (1986), 58-77.
 - [32] Dunavant, D. A., Szabó B. A., A-posteriori error indicators for the p-version of the finite element method, Internat. J. Numer. Methods Engrg. 10 (1983), 1851-1870.
 - [33] Frederiksson, B. and Mackerle, I., Structural Mechanics: Finite Element computer Programs, new up-to-date forth edition, Advanced Engineering Corp., Linkoping, Sweden, 1984.
 - [34] Gui, W., Babuška, I., The h, p and h-p versions of the finite element method in 1 dimension. Part 1: The error analysis of the p-version. Part 2: The error analysis of the h- and h-p versions. Part 3: The adaptive h-p version, Numerische Mathematik, 48 (1986), 577-612, 613-657, 619-683.
 - [35] Guo, B., The h-p version of the finite element method for elliptic equations of order $2m$. To appear in Numerische Mathematik.
 - [36] Guo, G., Babuška, I., The h-p version of the finite element method. Part 1: The basic approximation results. Part 2: General results and applications, Computational Mechanics 1 (1986), 21-41, 203-226.
 - [37] Guo, B., Babuška, I., The theory and practice of the h-p version of finite element method. Advances in computer methods in partial differential equations VI, R. Vichnevetsky and R. S. Stepleman, eds. Int'l Association for Mathematics and Computer Simulation (1987), 241-247.
 - [38] Izadpanah, K., Computation of the Stress Components of the Finite Element Method, Ph.D. Dissertation, Washington University, St. Louis, 1984.
 - [39] Katz, I. N., Wang, E. W., The p-version of the finite element method for problems requiring C^1 -continuity, SIAM J. Numer. Anal. 22 (1985), 1082-1106.

- [40] Noor, A. K., Books and monographs on finite element technology, Finite Elements in Analysis and Design, 1 (1985), 101-111.
- [41] Noor, A. K., Babuška, I., Quality assessment and control of finite element solutions, Finite Elements in Analysis and Design 3 (1987), 1-26.
- [42] Pammer, Z., Improved Structural Analysis Based on the p-Extended Finite Element Method. Presented at the Fifth World Congress, Oct. 2-9, Salzburg, Austria.
- [43] Patera, A. T., Advances and Future Directions of Research on Spectral Methods. Invited paper at ASME Winter Annual Meeting, Symposium on Future Directions in Computational Mechanics, 1986.
- [44] Peano, A. G., Walker, J. W., Modeling of Solid Continua by the p-Version of the Finite Element Method. Proc. 3rd World Congress on Finite Element Methods, Beverly Hills, California, 1981.
- [45] Rank, E., Babuška, I., An expert system for the optimal mesh design in the h-p version of the finite element method. To appear in Int. J. for Num. Meth. in Eng., 1987 (Tech. Note BN-1053, 1986, Institute for Physical Science and Technology, University of Maryland, College Park, MD 20742).
- [46] Rank, E., An Adaptive hp-Version in the Finite Element Method, in Numerical Techniques for Engineering Analysis and Design. Proceedings of the International Conference on Numerical Methods in Engineering: Theory and Applications. Numeta 87, Swansea, 6-10 July 1987. G. N. Pande & J. Middleton eds. Martinus Nijhoff Publishers, Dordrecht/Boston/Lancaster, 1987, paper S14.
- [47] Rank, E., Adaptive Boundary Element Methods, Rep. Nr 87/3, 1987, Fachgebiet Elektronisches Rechnen im Konstruktiven Ingenieurbau Technische Hochschule, München, Arcisstrasse 21, D-8000 - München 2.
- [48] Schiermeier, J. E., Szabó B. A., Optimal Design of a Torque Arm: A Case Study, Dep WU/CCM-86/2 Center for Computational Mechanics, 1986, Washington University, St. Louis, MO 63130.
- [49] Scott, L. R., Vogelius, M., Norm estimates for a maximal inverse of the divergence operator in spaces of piecewise polynomials, Math. Modelling and Num. Anal. (RAIRO) 19 (1985), 111-143.
- [50] Scott, L. R., Vogelius, M., Conforming Finite Element Methods for Incompressible and Nearly Incompressible Continua. Lectures in

- Applied Mathematics 22 (1985), 221-244.
- [51] Stephan, E. P., Suri, M., On the convergence of the p-version of the boundary element Galerkin method. To appear in Math. Comp.
 - [52] Suri, M. The p-Version of the Finite Element Method for Elliptic Problems in Computer Methods in partial Differential Equations VI, R. Vichnevetsky, R. S. Stepleman, eds. Int'l Association for Mathematical and Computer Simulation, 1987, 85-91.
 - [53] Suri, M. The p-Version of the Finite Element Method of elliptic equations of order 2k. To appear.
 - [54] Szabó, B. A., Probe: Theoretical Manual, 1985, NOETIC Tech, 7980 Clayton Road, Suite 205, St. Louis, MO 63117.
 - [55] Szabó, B. A., Mesh design for the p-version of the finite element method, Computer Methods in Applied Mathematics and Engineering 55 (1986), 181-197.
 - [56] Szabó, B. A., Implementation of a finite element software system with h- and p-extension capabilities, Finite Elements in Analysis and Design 2 (1986), 177-194.
 - [57] Szabó, B. A., On the Errors of Idealization and Discretization in Finite Element Analysis Advances in Computer Methods in Partial Differential Equations-VI, R. Vichnevetsky and R. S. Stepleman, eds. Int'l Association for Mathematics and Computer Simulation, 1987, 70-74.
 - [58] Szabó, B. A., Estimation and Control of Error Based on P-convergence in Accuracy Estimates and Adaptive Refinements in Finite Element computations (I. Babuška, J. Gago, E. R. de A. Oliveira, D. V. Zienkiwicz, eds.) John Wiley & Sons, 61-70.
 - [59] Szabó, B. A., On Geometric Idealization in Finite Element Computations, Tech. Note WU/CCM-87/4, Center for Computational Mechanics, Washington University, St. Louis, MO 63130. Presented at the Workshop on Special Topics in Computational Mechanics, Dallas, TX, April 13-14, 1987.
 - [60] Szabó, B. A., Computation of stress field parameters in areas of steep stress gradients. Comm. in Appl. Numer. Methods 2 (1986), 133-137.
 - [61] Szabó, B. A. and Babuška, I., Computation of the Amplitude of Stress Singular Terms for Cracks and Reentrant Corners, Rep. WU/CCM-86/1 Center for Computational Mechanics, Washington University, St. Louis,

MO 63130.

- [62] Szabó, B. A. and Peano, A. G., Hierarchic Finite Elements, Rep WU/CCM-83/1 Center for Computational Mechanics, Washington University, St. Louis, MO 63130.
- [63] Vasilopoulos, D., Treatment of Geometric Singularities with the p-Version of Finite Element Method. Ph.D. Dissertation, Washington University, St. Louis, MO, 1984.
- [64] Vogelius, M., An analysis of the p-version of the finite element method for nearly incompressible materials-uniformly valid, optimal estimates, Numer. Math. 41 (1983), 39-53.
- [65] Wang, D. W., Katz, I. N., Szabó, B. A., h and h-versions finite element analysis of a rhombic plate, Internat. J. Numer. Methods Engrg. 20 (1984), 1399-1405.
- [66] Wang, D. w. Katz, I. N., Szabó B. A., Implementation of C^1 triangular elements based on the p-version of the finite element method, Comp. & Structures 10 (1984), 381-392.
- [67] Wendland, W., Splines versus trigonometric polynomials. h-versus p-version in 2D boundary integral methods. Preprint Nr. 925, 1985, TH Darmstadt Fachbereich Mathematik.

This work was supported in part by the Office of Naval Research under Contract N00014-85-K-0169.

The Laboratory for Numerical analysis is an integral part of the Institute for Physical Science and Technology of the University of Maryland, under the general administration of the Director, Institute for Physical Science and Technology. It has the following goals:

- o To conduct research in the mathematical theory and computational implementation of numerical analysis and related topics, with emphasis on the numerical treatment of linear and nonlinear differential equations and problems in linear and nonlinear algebra.
- o To help bridge gaps between computational directions in engineering, physics, etc., and those in the mathematical community.
- o To provide a limited consulting service in all areas of numerical mathematics to the University as a whole, and also to government agencies and industries in the State of Maryland and the Washington Metropolitan area.
- o To assist with the education of numerical analysts, especially at the postdoctoral level, in conjunction with the Interdisciplinary Applied Mathematics Program and the programs of the Mathematics and Computer Science Departments. This includes active collaboration with government agencies such as the National Bureau of Standards.
- o To be an international center of study and research for foreign students in numerical mathematics who are supported by foreign governments or exchange agencies (Fulbright, etc.)

Further information may be obtained from Professor I. Babuška, Chairman, Laboratory for Numerical Analysis, Institute for Physical Science and Technology, University of Maryland, College Park, Maryland 20742.

END

DATE

FILMED

FEB.

1988