

AD-A187 451

ON AN OVERDETERMINED NEUMANN PROBLEM(U) MARYLAND UNIV
COLLEGE PARK C A BERNSTEIN 30 JUL 87 AFOSR-TR-87-1442
AFOSR-87-0073


1/1

UNCLASSIFIED

F/G 12/2

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

ITATION PAGE

DTIC FILE COPY

Form Approved
OMB No. 0704-0188

1a. REPORT AD-A187 451			1b. RESTRICTIVE MARKINGS		
2a. SECUR			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.		
2b. DECLASSIFICATION/CONTINUITY					
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR. 87-1442		
6a. NAME OF PERFORMING ORGANIZATION		6b. OFFICE SYMBOL (if applicable)		7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code)				7b. ADDRESS (City, State, and ZIP Code)	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (if applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-87-CX073	
8c. ADDRESS (City, State, and ZIP Code)				10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO.		PROJECT NO.	TASK NO.
					WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification)					
12. PERSONAL AUTHOR(S)					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) July 30, 1987	
15. PAGE COUNT					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
DTIC ELECTE S NOV 16 1987 D H					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT ☐ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT. ☐ DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION		
22a. NAME OF RESPONSIBLE INDIVIDUAL			22b. TELEPHONE (Include Area Code)		22c. OFFICE SYMBOL

AFOSR-TR. 87-1442



Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

On an overdetermined Neumann problem

Carlos A. Berenstein*

This note is a summary of recent results obtained jointly with Paul Yang. The author would like to express his gratitude to Professor S. Coen for the invitation to deliver this lecture at the Università di Bologna.

Several questions in harmonic analysis, partial differential equations and applied mathematics lead to the question of characterizing domains for which overdetermined boundary value problems have solutions. Given the existence of an excellent bibliography in [1] I will not attempt to trace the history of the problem (D) and (N) introduced below, except to say that their origins go back to the treatise [2] of Lord Rayleigh.

Let Ω be an open relatively compact subset of real analytic Riemannian manifold M . Assume further that $\partial\Omega$ is connected and

*Partially supported by the National Science Foundation and AFOSR-URI grant number AFOSR-87-0073

10-14 348

of Lipschitz class. What can we say about Ω if any of the following problems (D) or (N) has an eigenvalue $\alpha > 0$?

$$(D) \quad \begin{cases} \Delta u + \alpha u = 0 & \text{in } \Omega \\ u = 0 \text{ and } \frac{\partial u}{\partial n} = 1 & \text{on } \partial\Omega \\ \text{(overdetermined Dirichlet problem)} \end{cases}$$

$$(D) \quad \begin{cases} \Delta u + \alpha u = 0 & \text{in } \Omega \\ u = 1 \text{ and } \frac{\partial u}{\partial n} = 1 & \text{on } \partial\Omega \\ \text{(overdetermined Neumann problem)} \end{cases}$$

It is remarkable that the existence of such an eigenvalue already implies that $\partial\Omega$ is a real analytic submanifold of M (at least if $\partial\Omega$ is of class $C^{2+\varepsilon}$; if $M = \mathbb{R}^n$ or $\mathbb{H}^n$ - real hyperbolic space - it is enough if $\partial\Omega$ is Lipschitz, this follows from work of Caffarelli [3]. It is quite possible that this last condition is always sufficient). In order to proceed any further we have to impose restrictions on both M and α . For instance, if $M = \mathbb{R}^n$ (resp. $\mathbb{H}^n$) and α in (D) is the first eigenvalue of Dirichlet problem, i.e. $u > 0$, then $\Omega = \text{ball}$ (resp. geodesic ball). This follows from a theorem of Serrin [4]. A few other cases are known [5], [6], [7] of the non-existence of eigenvalues if $M = \mathbb{R}^n$, $\Omega \neq \text{ball}$. On the otherhand, if $M = \mathbb{R}^n$ (resp. $\mathbb{H}^n$) and $\Omega = \text{ball}$ (resp. geodesic ball) then one can see that, considering the radial eigenfunctions of the usual Dirichlet or Neumann problems in the role of u , there are infinitely many eigenvalues for (D) and for (N). Sometime ago I have proved in $\mathbb{R}^2$ that the converse was true [5], later I obtained the same result with P. Yang in $\mathbb{H}^2$ [8]. We have now:

Theorem [9] Let $M = \mathbb{R}^n$ (resp. $\mathbb{H}^n$), the existence of infinitely many eigenvalues for either of the problems (D) or (N) characterizes the balls (resp. geodesic balls) among all the

relatively compact domains Ω with connected Lipschitz boundary.

Note that in the Poincaré model $\mathbb{H}^n =$ unit ball of $\mathbb{R}^n$ and the geodesic balls are then euclidean balls.

It would seem to be natural to jump to the conclusion that this theorem should remain true in all M . The following example shows that one needs some caution.

Example Let $u = u(x) = x_1^2 - x_2^2 + \dots + x_{2n-1}^2 - x_{2n}^2$ defined in $\mathbb{R}^{2n}$, denote by v its restriction to $M = S^{2n-1}$. Denote by L the Laplace operator in $\mathbb{R}^{2n}$, Δ the Laplace Beltrami operator in M . It is well known they are related by

$$(1) \quad L = \frac{\partial}{\partial r^2} + \frac{2n-1}{r} \frac{\partial}{\partial r} + \frac{1}{r^2} \Delta$$

We also have

$$\frac{\partial u}{\partial r} = \nabla u \cdot \frac{x}{r} = \frac{2u}{r}, \quad r = |x|,$$

where the last identity holds by Euler's formula, u being homogeneous of degree 2. Therefore

$$(2) \quad \frac{\partial^2 u}{\partial r^2} = \frac{2}{r} \frac{\partial u}{\partial r} - \frac{2}{r^2} u = \frac{2u}{r^2}.$$

It follows that

$$\frac{2u}{r^2} + \frac{(2n-1)}{r^2} 2u + \frac{1}{r^2} \Delta u = Lu = 0,$$

and the function v satisfies $\Delta v + 4n v = 0$ on $\{r=1\} = M$.

Consider now the set $\Omega =$ connected component containing

$(1, 0, \dots, 0)$ of $\{x \in M: u(x) > 0\}$. Then on $\partial\Omega$ we have $v(x) = 0$

and therefore $\alpha = 4n$ is the first eigenvalue of Dirichlet

problem for Ω . We claim now that $\frac{\partial v}{\partial n}$ constant on $\partial\Omega$. Note

that the (exterior) normal derivative is an operator tangent to

the sphere M . First, observe that (2) implies

$$(3) \quad \nabla u(x) \perp x \quad \text{if } x \in \partial Q.$$

Since x is the normal vector to M in $\mathbb{R}^{2n}$ we have

$\nabla u(x) \in T_x M$ for $x \in \partial Q$. Therefore we have

$$(4) \quad \nabla u(x) = \frac{\partial v}{\partial \vec{n}}(x) \cdot \vec{n}(x) \quad \text{if } x \in \partial Q,$$

where $\vec{n}$ represents the unit exterior normal to Q in M . We only need to verify that $|\nabla u(x)| = \text{constant}$ when $x \in \partial Q$. But

$$\nabla u(x) = 2(x_1, -x_2, \dots, x_{2n-1}, -x_{2n})$$

and

$$(5) \quad |\nabla u|^2 = 4|x|^2 = 4 \quad \text{if } x \in M,$$

which says that v satisfies:

$$(6) \quad \begin{cases} \Delta v + 4n v = 0 & \text{in } Q \\ v > 0 & \text{in } Q \\ v = 0 & \text{on } \partial Q \\ \frac{\partial v}{\partial \vec{n}} = -2 & \text{on } \partial Q. \end{cases}$$

Note that topologically $\partial Q = S^{n-1} \times S^{n-1}$ which shows that Serrin's theorem fails on M . We want to show now that there are infinitely many solutions for both (D) and (N) in Q .

Let f be a twice differentiable function of a single real variable and define

$$\varphi(x) := f(u(x)).$$

Using again identity (2) we have

$$L\varphi = f''(u)|\nabla u|^2 + f'(u)Lu = f''(u)4|x|^2 = 4f''(u) \quad \text{on } M.$$

$$\frac{\partial \varphi}{\partial r} = f'(u) \frac{\partial u}{\partial r} = 2f'(u) \frac{u}{r} = 2f'(u)u \quad \text{on } M.$$

$$\begin{aligned} \frac{\partial^2 \varphi}{\partial r^2} &= 4f''(u) \left(\frac{u}{r} \right)^2 + \frac{2}{r} f'(u) \frac{\partial u}{\partial r} - \frac{2}{r^2} f'(u)u = \\ &= \frac{4}{r^2} f''(u)u^2 + \frac{2}{r} f'(u)u = 4f''(u)u^2 + 2f'(u)u \quad \text{on } M. \end{aligned}$$

Hence, on M ,

$$L\varphi = 4f''(u) = 4f''(u)u^2 + 4nf'(u)u + \Delta\varphi,$$

and

$$(7) \quad \Delta\varphi + \alpha\varphi = 4f''(u)(1-u^2) - 4nf'(u)u + \alpha f(u).$$

Therefore the equation $\Delta\varphi + \alpha\varphi = 0$ in Ω becomes the ordinary differential equation

$$(8) \quad 4(1-t^2)f''(t) - 4ntf'(t) + \alpha f(t) = 0, \quad 0 \leq t \leq 1$$

This equation has a regular singular point at end point $t = 1$.

Each eigenvalue α and eigenfunction of (8) satisfying

$$(9) \quad f(1) \text{ bounded, } f(0) = 0$$

provides an eigenvalue for (D) since

$$\nabla\varphi = f'(u)\nabla u$$

hence again $\nabla\varphi \cdot x = 0$ on $\partial\Omega$ and to check whether $\frac{\partial\varphi}{\partial n} = \text{constant}$ we only need to compute $|\nabla\varphi|^2$ on $\partial\Omega$. But

$$|\nabla\varphi|^2 = (f'(u))^2 |\nabla u|^2 = 4(f'(u))^2,$$

which shows

$$(10) \quad \frac{\partial\varphi}{\partial n} = \pm 2f'(0) \quad \text{on } \partial\Omega.$$

(This is different from zero since the eigenfunctions of (8) - (9) satisfy $f'(0) \neq 0$). This same computation shows that the eigenfunctions of (8) satisfying

$$(11) \quad f(1) \text{ bounded, } f'(0) = 0,$$

will provide eigenfunctions $\varphi(x)$ for (N). This time $\frac{\partial\varphi}{\partial n} = 0$ on $\partial\Omega$ and $\varphi = f(0) \neq 0$ on $\partial\Omega$. Since it is a well known theorem of the theory of ordinary differential equations that (8)-(9) and (8)-(10) have infinitely many eigenvalues, the domain Ω has infinitely many eigenvalues for (D) and (N). Ω is not even topologically a geodesic ball in M .

On this note we leave the reader to reflect on these beautiful questions.

References:

1. P. Berard, Lectures on the spectral properties of the Laplace operator, Brazil, 1985.
2. Lord Rayleigh, The Theory of Sound, MacMillan, 1877.
3. L. Cafarelli, The regularity of free boundaries in higher dimensions, Acta.Math. 139 (1977), 155-184.
4. J. Serrin, A Symmetry problem in potential theory, Arch.Rat. Mech.Anal. 43 (1971), 304-318.
5. C. Berenstein, An inverse spectral theorem and its relation to the Pompeiu problem, J.Anal.Math. 37 (1980), 128-144.
6. P. Aviles, Symmetry theorems related to Pompeiu's problem, Amer.J.Math. 108 (1986), 1023-1036.
7. L. Brown and J-P. Kahane, A note on Pompeiu problem for convex domains, Math.Anal.259 (182), 107-110.
8. C. Berenstein and P. Yang, An overdetermined Neumann problem in the unit disk, Advances in Math. 44(1982), 1-17.
9. C. Berenstein and P. Yang, An inverse Neumann problem, preprint 86-13 of the Systems Research Center, University of Maryland, to appear in J. Reine Angew.Math.
10. C. Berenstein, Autovalores del Laplacians y Geometria, ELAM., Caracas, 1984.

Approved for public release;
distribution unlimited.

Author's address:

Department of Mathematics
and Systems Research Center,
University of Maryland
College Park, MD 20742.
USA.

1986 RELEASE UNDER E.O. 14176
U.S. GOVERNMENT PRINTING OFFICE: 1980-12
GPO, Washington, D.C. 20540
Classification Division

END

FEB.

1988

DTic