

AD-A109 353


SUPERCONDUCTORS WITH STRUCTURED SURFACES FIELDS AND
CURRENTS(U) STATE UNIV OF NEW YORK AT BUFFALO DEPT OF
CHEMISTRY Z C MU ET AL. DEC 87 UBUFFALO/DC/87/TR-62
N00014-86-K-0043 F/G 28/3

1/1

UNCLASSIFIED

NL





OFFICE OF NAVAL RESEARCH
Contract N00014-86-K-0043
TECHNICAL REPORT No. 62

Superconductors with Structured Surfaces: Fields and Currents

by

Z. C. Wu, Daniel A. Jelski and Thomas F. George

Prepared for Publication

in

Laser and Particle-Beam Chemical Processing for Microelectronics
Edited by D. J. Ehrlich, G. S. Higashi and M. M. Oprysko
Materials Research Society Symposium Proceedings, Volume 101

Departments of Chemistry and Physics
State University of New York at Buffalo
Buffalo, New York 14260

December 1987

DTIC
ELECTE
DEC 30 1987
S D

Reproduction in whole or in part is permitted for any purpose of the
United States Government.

This document has been approved for public release and sale;
its distribution is unlimited.

87 12 21 129 3

AD-A189 353

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

AD-A18935-3

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/87/TR-62			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260			7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract N00014-86-K-0043		
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
					WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Superconductors with Structured Surfaces: Fields and Currents					
12. PERSONAL AUTHOR(S) Z. C. Wu, Daniel A. Jelski and Thomas F. George					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) December 1987	
				15. PAGE COUNT 7	
16. SUPPLEMENTARY NOTATION Prepared for publication in <u>Laser and Particle-Beam Chemical Processing for Microelectronics</u> , edited by D. J. Ehrlich, G. S. Higashi and M. M. Oprysko, Materials Research Society Symposium Proceedings, Volume 101					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	SUPERCONDUCTOR.		
			STRUCTURED SURFACE		
			SUPERCURRENT		
			EXTERNAL MAGNETIC FIELD		
			CRITICAL NUCLEATION FIELD		
			GINZBURG-LANDAU THEORY		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
This paper discusses the behavior of currents and fields along a structured superconductor. First the effect of surface structure on supercurrents is investigated. Then the effect of structure on the critical nucleation field is discussed in two cases, one with the magnetic field parallel to the ripples and the other with the field parallel to the grating wavenumber. In the first case, it is found that the critical field is reduced as a function of grating height, whereas in the latter case it is increased. Finally, the relevance of this work for laser-induced chemistry above a superconducting surface is discussed. The Ginzburg-Landau model is used throughout.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT ☒ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT ☐ DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

DD Form 1473, JUN 86

Previous editions are obsolete

SECURITY CLASSIFICATION OF THIS PAGE

UNCLASSIFIED

SUPERCONDUCTORS WITH STRUCTURED SURFACES: FIELDS AND CURRENTS

Z.C. WU, DANIEL A. JELSKI AND THOMAS F. GEORGE

Departments of Physics & Astronomy and Chemistry, 239 Fronczak Hall, State University of New York at Buffalo, Buffalo, New York 14260



ABSTRACT

This paper discusses the behavior of currents and fields along a structured superconductor. First the effect of surface structure on supercurrents is investigated. Then the effect of structure on the critical nucleation field is discussed in two cases, one with the magnetic field parallel to the ripples and the other with the field parallel to the grating wavenumber. In the first case, it is found that the critical field is reduced as a function of grating height, whereas in the latter case it is increased. Finally, the relevance of this work for laser-induced chemistry above a superconducting surface is discussed. The Ginzburg-Landau model is used throughout.

A-1

INTRODUCTION

The enormous recent interest in superconductivity [1] is no longer worthy of remark. In this paper, we shall not discuss the fundamental theory behind the phenomenon, nor shall we consider the specific structure of any given material. Instead, we shall investigate a generic superconducting material along the surface of which is inscribed a sinusoidal grating. We are ultimately interested in investigating the optical properties of thin films, which we suppose have microscopic roughness on the surface. While we are not yet able to treat the time-dependent problem satisfactorily, we shall discuss the response of the rippled surface to external fields. We wish to use the Ginzburg-Landau theory [2] to analyze the effect of the grating amplitude on the current, the superelectron density and, most importantly, on the critical nucleation field. The reason for doing this is to get a better idea of how a thin-film superconductor will respond to external fields. While we are at present restricted to static fields (the Ginzburg-Landau theory is not time dependent), we suppose the result to be valid in the low-frequency domain. We hope, eventually, to develop an explicit time-dependent model.

The Ginzburg-Landau (GL) model is very appropriate here. Its validity is largely independent of the microscopic causes of superconductivity, the microscopic theory being used only to evaluate necessary parameters. Further, it is an expansion in terms of the order parameter, the superelectron density, which may be expected to go to zero at the critical temperature. Hence the limiting condition is not what the critical temperature is, nor what the microscopic causes are, but simply that the temperature be close to T_c . Thus we expect our current results to apply equally to the new high- T_c ceramic materials as to more traditional substances.

In the next section, we shall discuss the effect of the grating amplitude on the supercurrent. Then we shall consider the critical nucleation field for two cases: the magnetic field parallel to the wavenumber (x-direction) and parallel to the

WU, JELSKI, GEORGE

1

6

ripples (y-direction). We shall find that the critical field is reduced in the first case, while it is increased in the second. Finally, we shall consider some implications of this work for optics and chemistry above superconducting surfaces.

ELECTRON DENSITY AND SUPERCURRENTS

The behavior of the GL equation near a flat surface has been discussed by de Gennes [3]. We have extended this result to allow for a rippled surface [4] by solving a two-dimensional elliptic equation rather than the one-dimensional harmonic equation. A more detailed discussion appears in that reference. A problem arises in choosing the proper boundary conditions. The appropriate boundary condition for the GL equation at a surface is

$$\left(\frac{\hbar}{2} \vec{\nabla} - \frac{e^*}{c} \vec{A} \right) \psi \Big|_n = \frac{1}{b} \psi \quad (1)$$

This condition can be satisfied in at least two ways. We can impose Dirichlet boundary conditions, where we set $b = 0$, which implies $\psi = 0$, or we can impose von Neumann boundary conditions, where $b = \infty$. The electron density function is, of course, strongly dependent on which boundary conditions one chooses. Figures 1 and 2 illustrate the differences. Figure 1 shows the effect of the Dirichlet conditions at the ripples. Here the density is sharply reduced in the "fingers". Conversely, the density calculated from imposing von Neumann conditions is shown in Fig. 2. Here the derivative of the wavefunction is zero across the boundary. Note that the greatest electron density is just below the ripples.




Figure 1
The electron density near a rippled surface given Dirichlet boundary conditions. See Ref. 4 for details. The z-coordinate denotes the normal distance from the surface into the interior of the material.




Figure 2
The electron density near a rippled surface given von Neumann boundary conditions. See Ref. 6 for details.

Let us now restrict our attention to the Dirichlet problem, and this for the following reason. We are using the linearized GL equation to solve the problem. The linear equation is valid as long as ψ is much smaller than ψ_∞ , where $|\psi_\infty|^2$ is the bulk value of the superelectron density. There are at least two ways in which this approximation can be made valid. One is when we are near the surface and the Dirichlet conditions hold, and the other is for an external field very near the critical field. Since we are presently considering currents, and not fields, we are thus restricted by the linear approximation to the Dirichlet condition. DeGennes has shown from an analysis based on the microscopic theory (BCS) that superconductors in zero field obey the Dirichlet conditions. In subsequent sections we shall consider the high field case where the von Neumann conditions are appropriate.

It has been found [4] that ripples increase the current path length. The ripples create vortices in the current flow. However, it has also been discovered that the ripples provide essentially the same Meissner-effect shielding as a "bulk" surface would. Hence, if the goal is to maximize current flow, then as flat a surface as possible is desirable. Conversely, if the purpose is to provide a constant current impervious to small changes in the external field, then the ripples are desirable.

CRITICAL NUCLEATION FIELD PARALLEL TO GRATING WAVENUMBER

It is interesting to calculate the critical field at which superconductivity can nucleate near a surface in a decreasing magnetic field, H_{C3} . At this critical field, the superelectron density must be infinitesimal and the linearized version of the GL equations becomes accurate. The analogous nucleation problem for the bulk material, or near a flat surface, has been solved by deGennes and Saint-James [3,5]. It is found that H_{C3} for a flat surface is $1.695 \cdot H_{C2}$, where H_{C2} is the critical field for the bulk.

We have investigated the critical nucleation field for a superconducting surface with a sinusoidal grating [6]. In this case, two mutually perpendicular directions along the surface must be distinguished. In this section, we shall consider the field to be parallel to the grating wavenumber, which we shall consider to be the x-direction. Then translational symmetry along this axis disappears, and the GL equation becomes two-dimensional. We use the von Neumann boundary conditions for the nucleation problem since we are interested in the superelectron density at the surface, precluded under the Dirichlet conditions. The problem is to find the maximum field under which the GL equation has a bound solution, i.e., the wavefunction must decay exponentially into the interior of the material. This equation must be solved numerically. There is a further parameter, namely the supercurrent density in the y-direction (parallel to the ripple), and this is chosen variationally to minimize the eigenvalue, which corresponds to maximum field.

Under these circumstances, it is found that the critical field decreases as a function of grating amplitude (Fig. 3). In the limit of infinitely large amplitude, it is found that the nucleation field approaches that of the bulk limit, which is to be expected from an analysis of a series of parallel thin films.




Figure 3
Nucleation critical field vs. grating amplitude for a field oriented parallel to the grating wavenumber. See Ref. 6 for details. The unit for the field is $2\pi m^* c |\alpha| / \hbar e$.

CRITICAL NUCLEATION FIELD PARALLEL TO THE RIPPLE

Another interesting case we now consider is the nucleation field in the ripple direction (y-direction). The gauge for the vector potential is

$$\vec{A} = (H_y z, 0, 0) .$$

The GL equation can be written as

$$-\frac{\partial^2 f(x,z)}{\partial x^2} - \frac{\partial^2 f(x,z)}{\partial z^2} + \frac{4\pi i \hbar^2 y z}{\phi_0} \frac{\partial f(x,z)}{\partial x^2} + \left(\frac{2\pi \hbar^2 y z}{\phi_0} \right)^2 f(x,z) = -\frac{2m^* \alpha}{\hbar^2} f(x,z) , \quad (2)$$

where

$$\phi_0 = \frac{\hbar c}{2e} ,$$

and we have used the ansatz

$$\psi = e^{ik_y y} f(x,z) ,$$

due to translational symmetry along the y-axis. Equation (2) must be supplemented by the von Neumann boundary conditions at the surface, given by Eq. (1) with $b = \infty$. The critical field must be the maximum H_y at which one can find a bound solution, i.e., the wavefunction must decay exponentially into the bulk. There we set the boundary condition so that the magnitude of the wave-function must be constant at constant z . The phase of the wavefunction, at this bottom boundary, is a variational parameter, chosen to maximize the nucleation field. Unlike the previous case, our calculation shows that the critical field increases as a function of grating amplitude (Fig. 4). Here we have used units such that $H_y / \phi_0 = 1$. It is expected that the value of the critical field will reach the value for superconducting films, with a thickness of the same order as the period of the structure. It is noted that if the thickness of a film is much smaller than the GL coherence length, the critical field must be much greater than the bulk value.




Figure 4
Nucleation critical field vs. grating amplitude for a field oriented parallel to the ripples. The units are the same as for Fig. 3.

CONCLUSIONS

We have considered the behavior of a Ginzburg-Landau superconductor as a function of grating amplitude. We have discussed the two types of problems for which the linearized GL theory is relevant. These are the current density when Dirichlet conditions apply, and the critical nucleation field when von Neumann conditions apply. In subsequent work, we hope to remove the linear condition (i.e. find a quick way of solving the nonlinear equation), and we would also like to remove the condition of static fields.

We have found that if the magnetic field is oriented in a direction parallel to the ripples, then the nucleation field is increased. Conversely, if the field is oriented in the direction of the wavenumber, then the nucleation field is decreased. It follows that if an incident laser of frequency well below the gap frequency is p-polarized with respect to the surface, the intensity of the laser could be stronger than would otherwise be possible, and still not destroy the existence of the superconducting phase along the surface. Similarly, if the laser is s-polarized, the surface nucleation field reduces to that of the bulk in the large amplitude limit.

This research forms the first step in elucidating the behavior of a superconducting surface under laser irradiation. In order to finish this project, it will be necessary to consider the frequency dependence of the GL equations, and also to consider the relation between the superconducting electrons, the normal electrons, and the resulting dielectric constant of the material. Given this information, it will be very interesting to calculate such observables as the reflectivity of a superconducting grating, to consider the existence or non-existence of plasmon waves along the surface, or to evaluate the response of a molecular dipole above the surface.

We close with a few words about the validity of GL theory when applied to the new high T_c superconductors. It appears that the Pippard coherence length of the ceramics is on the order of 4Å, which is much too short to make the present calculation worthwhile. However, the GL coherence length is a function of temperature and increases as the temperature approaches the critical value, although as the temperature comes very near the critical value, fluctuation effects will dominate and invalidate our calculation. Thus we suppose that for higher temperatures, above liquid nitrogen, some of the calculated results here should be observable. In particular, the qualitative variation of the behavior of the critical field depending on orientation should be measurable.

ACKNOWLEDGMENTS

This research was supported by the Office of Naval Research and the Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract F49620-86-C-0009. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon.

REFERENCES

1. J.G. Bednorz and K.A. Müller, Z. Phys. B 64, 189 (1986).
2. V.L. Ginzburg and L.D. Landau, Zh. Eksperim. i Teor. Fiz. 20, 1064 (1950); L.P. Gor'kov, Zh. Eksperim. i Teor. Fiz., 36, 1918 (1959); M. Tinkham, Introduction to Superconductivity (McGraw-Hill, New York, 1975), Chapt. 5.
3. P.G. deGennes, Superconductivity of Metals and Alloys (W.A. Benjamin, New York, 1966), Chapt. 6.
4. D.A. Jelski, Z.C. Wu and T.F. George, J. Chem. Phys., submitted.
5. D. Saint-James and P.G. deGennes, Phys. Lett. 7, 306 (1983).
6. Z.C. Wu, D.A. Jelski and T.F. George, Phys. Rev. B, submitted.

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. C. B. Harris
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. H. Weaver
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. F. Kutzler
Department of Chemistry
Box 5055
Tennessee Technological University
Cookeville, Tennessee 38501

Dr. A. Reisman
Microelectronics Center of North Carolina
Research Triangle Park, North Carolina
27709

Dr. D. DiLella
Chemistry Department
George Washington University
Washington D.C. 20052

Dr. M. Grunze
Laboratory for Surface Science and
Technology
University of Maine
Orono, Maine 04469

Dr. R. Reeves
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. J. Butler
Naval Research Laboratory
Code 6115
Washington D.C. 20375-5000

Dr. Steven M. George
Stanford University
Department of Chemistry
Stanford, CA 94305

Dr. L. Interante
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Mark Johnson
Yale University
Department of Chemistry
New Haven, CT 06511-8118

Dr. Irvin Heard
Chemistry and Physics Department
Lincoln University
Lincoln University, Pennsylvania 19352

Dr. W. Knauer
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. K.J. Klaubunde
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. Murday
Naval Research Laboratory
Code 6170
Washington, D.C. 20375-5000

Dr. J. B. Hudson
Materials Division
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Theodore E. Madey
Surface Chemistry Section
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

Dr. J. E. Demuth
IBM Corporation
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. M. G. Lagally
Department of Metallurgical
and Mining Engineering
University of Wisconsin
Madison, Wisconsin 53706

Dr. R. P. Van Duyne
Chemistry Department
Northwestern University
Evanston, Illinois 60637

Dr. J. M. White
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. D. E. Harrison
Department of Physics
Naval Postgraduate School
Monterey, California 93940

Dr. R. L. Park
Director, Center of Materials
Research
University of Maryland
College Park, Maryland 20742

Dr. W. T. Peria
Electrical Engineering Department
University of Minnesota
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson
Department of Metallurgy and
Materials Science
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. S. Sibener
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Arnold Green
Quantum Surface Dynamics Branch
Code 3817
Naval Weapons Center
China Lake, California 93555

Dr. A. Wold
Department of Chemistry
Brown University
Providence, Rhode Island 02912

Dr. S. L. Bernasek
Department of Chemistry
Princeton University
Princeton, New Jersey 08544

Dr. W. Kohn
Department of Physics
University of California, San Diego
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Richard Colton
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Dan Pierce
National Bureau of Standards
Optical Physics Division
Washington, D.C. 20234

Dr. R. Stanley Williams
Department of Chemistry
University of California
Los Angeles, California 90024

Dr. R. P. Messmer
Materials Characterization Lab.
General Electric Company
Schenectady, New York 22217

Dr. Robert Gomer
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Ronald Lee
R301
Naval Surface Weapons Center
White Oak
Silver Spring, Maryland 20910

Dr. Paul Schoen
Code 6190
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. John T. Yates
Department of Chemistry
University of Pittsburgh
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene
Code 5230
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. L. Kesmodel
Department of Physics
Indiana University
Bloomington, Indiana 47403

Dr. K. C. Janda
University of Pittsburgh
Chemistry Building
Pittsburg, PA 15260

Dr. E. A. Irene
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Adam Heller
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton SO9 5NH
UNITED KINGDOM

Dr. H. Tachikawa
Chemistry Department
Jackson State University
Jackson, Mississippi 39217

Dr. John W. Wilkins
Cornell University
Laboratory of Atomic and
Solid State Physics
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis
Department of Physics
University of California
Irvine, California 92664

Dr. D. Ramaker
Chemistry Department
George Washington University
Washington, D.C. 20052

Dr. J. C. Hemminger
Chemistry Department
University of California
Irvine, California 92717

Dr. T. F. George
Chemistry Department
University of Rochester
Rochester, New York 14627

Dr. G. Rubloff
IBM
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. Horia Metiu
Chemistry Department
University of California
Santa Barbara, California 93106

Dr. W. Goddard
Department of Chemistry and Chemical
Engineering
California Institute of Technology
Pasadena, California 91125

Dr. P. Hansma
Department of Physics
University of California
Santa Barbara, California 93106

Dr. J. Baldeschwieler
Department of Chemistry and
Chemical Engineering
California Institute of Technology
Pasadena, California 91125

Dr. J. T. Keiser
Department of Chemistry
University of Richmond
Richmond, Virginia 23173

Dr. R. W. Plummer
Department of Physics
University of Pennsylvania
Philadelphia, Pennsylvania 19104

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 44106

Dr. N. Winograd
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. Roald Hoffmann
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. A. Steckl
Department of Electrical and
Systems Engineering
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. G.H. Morrison
Department of Chemistry
Cornell University
Ithaca, New York 14853

END

DATE

FILMED

MARCH

1988

DTIC