

AD-A191 252

ENHANCED OXIDATION AND SOLVOLYSIS REACTIONS IN

1/1

CHEMICALLY INERT MICROWAVE (U) ECOLE POLYTECHNIQUE

FEDERALE DE LAUSANNE (SWITZERLAND) A N BRAS 24 OCT 87

UNCLASSIFIED


DAJ449-85-C-0828

P/C 7/4

ML



END
PAGE
001
OF
001



MICROCOPY RESOLUTION TEST CHART
 NATIONAL BUREAU OF STANDARDS-1963-A

AD-A191 252

5038-CH-01.

②

ENHANCED OXIDATION AND SOLVOLYSIS REACTIONS
IN CHEMICALLY INERT MICROHETEROGENEOUS SYSTEMS

PD Dr. André M. BRAUN
Ecole Polytechnique Fédérale de Lausanne

THE 11 0022

DAJA45-85-C-0026, P00001

Fifth Interim Report
January - September 1987

DTIC
ELECTE
S FEB 18 1988 D
E

The research reported in this document has been made possible through the support and sponsorship of the U.S. Government through its European Research Office of the U.S. Army. ~~The report is intended only for the internal management use of the Contractor and the U.S. Government.~~

88 2 11 071 1



SUMMARY

In the second year of contract, the following parts of the original project have been accomplished:

1. Model calculations of the kinetics of the decomposition of H_2O_2 in H_2O and fitting of the model of sequence and rates of reactions to the experimental results.
 2. Synthesis of perfluoro 2-methyl-2-propanol
 3. Formulation of microemulsions containing totally perfluorinated components:
 - sodium perfluorocaprylate
 - perfluoro n-hexane
 - perfluoro 2-methyl-2-pentanol or perfluoro t-butanol
 - water
- 











SHORT TERM PROJECTS

The projects for the third year of contract have already been transmitted (1). The work will be done in close collaboration with Dres. M.-T. Maurette and E. Oliveros (Université Paul Sabatier, Toulouse).

MECHANISM OF H_2O_2 DECOMPOSITION. MODEL CALCULATIONS AND KINETIC EXPERIMENTS

After having accomplished an extensive investigation on the kinetics of H_2O_2 decomposition in H_2O and in perhydrogenated four components microemulsions (s.Final Report, 1986) we began in this second year of contract to study some sequences of reactions which may be of importance for the observed decomposition of H_2O_2 and which could be fitted to the experimental kinetic results in using rate constants independently determined.

The general scheme of reactions comprises:



The rate and equilibrium constants have been taken as determined and published independently by different authors³⁻⁹:

$$K_1 = 2.4 \times 10^{-12} \text{ M}^3$$

$$k_2 = 8.6 \times 10^{-6} \text{ M}^{-1} \text{ s}^{-1} \quad 3$$

$$k_3 = 4.7 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1} \quad 4$$

$$K_3 = 1.49 \times 10^{-5} \text{ M}^4$$

$$k_4 = 8.5 \times 10^7 \text{ M}^{-1} \text{ s}^{-1} \quad 5$$

$$k_5 = 6 \text{ M}^{-1} \text{ s}^{-1} \quad 6$$

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



$$k_6 = 1 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1} \text{ }^6$$

$$k_7 = 8.5 \times 10^5 \text{ M}^{-1} \text{ s}^{-1} \text{ }^7$$

$$k_8 = 2.65 \times 10^7 \text{ M}^{-1} \text{ s}^{-1} \text{ }^8$$

$$k_9 = 5.5 \times 10^9 \text{ M}^{-1} \text{ s}^{-1} \text{ }^9$$

$$k_{10} = 1.3 \times 10^{11} \text{ M}^{-1} \text{ s}^{-1} \text{ }^{10}$$


$$K_{10} = 10^{-14} \text{ M}$$


The fitting to our experimental results (o) is shown in figures 1 and 2. The model fits in fact excellently yielding only slight changes for k_1 and k_5 , as well as a major variation on k_3 which has to be taken as a first order rate constant ($7.0 \times 10^5 \text{ s}^{-1}$).

A complete fitting requires some experiments with times of reaction between 700 and 1000 h in order to confirm the acceleration of the reaction, as predicted by the model (figures 1 and 2). These experiments are currently undertaken in an appropriately stabilized equipment. The results already obtained confirm qualitatively the shape of the calculated curves.

SYNTHESIS OF PERFLUORINATED 2-METHYL-2-PENTANOL

Perfluoro 2-methyl-2-pentanol has been prepared using and optimizing the procedure of K.V. Scherer, Jr. et al.¹¹:





The synthesis is currently undertaken on a level of 50 g of substrate in order to prepare sufficient material for the planned kinetic experiments.

FORMULATION OF NEW MICROEMULSIONS CONTAINING THREE PERFLUORINATED COMPONENTS AND WATER

Following the procedure already published¹², three new pseudo-ternary phase diagrams have been established concerning the following formulations:

- sodium perfluorocaprylate/perfluoro 2-methyl-2-pentanol (1/1), perfluoro n-hexane, water (figure 3)
- sodium perfluorocaprylate/perfluoro 2-methyl-2-pentanol (1/2), perfluoro n-hexane, water (figure 4)
- sodium perfluorocaprylate/perfluoro t-butanol (1/2), perfluoro n-hexane, water (figure 5)

CONCLUSION

The results of the work made possible through this contract provide the basis for

- the confirmation of the observed carboxylate catalyzed decomposition of H₂O₂ in perfluorinated microemulsions,
- the determination of OH. radical recombination in a chemically inert environment,

- the application of perfluorinated microheterogeneous media for oxidation reactions and efficiency determinations with respect to aqueous and perhydrogenated microheterogeneous media.

REFERENCES

- 1 A.M. Braun, DAJA45-85-C-0026, Project 3rd year (October 1987-September 1988)
- 2 A.M. Braun, DAJA45-85-C-0026, Final Report, 1986
- 3 S. Rinaudo, thesis, Université de Haute Alsace, Mulhouse, 1985
- 4 I.B. Afanas'ev, Russ.Chem.Rev. 48, 977 (1979)
- 5 D.E. Adams, E.M. Fielden, B.D. Michael, "The chemistry of peroxy radicals and its implications to radiation biology" in "Fast processes in radiation chemistry and biology", M. Simic, ed., Wiley, New York, 1975, p. 162
- 6 D.T. Sawyer, M.J. Gribble, Tetrahedron 35, 1471 (1979)
- 7 A.A. Frimer, "Organic reactions involving the superoxide anion", in "The Chemistry of Functional Groups, Peroxides", S. Patai, ed., Wiley, New York, 1983, chap. 14, p. 439
- 8 C. Walling, R.E. Partch, T. Weil, Proc.Natl.Acad. Sci.USA, 72, 140 (1975)
- 9 J. Rabani, S.O. Nielsen, J.Phys.Chem. 73, 3736 (1969)
- 10 Z. Haniotis, Ciba-Geigy AG, Basel, private communication
- 11 K.V. Scherer, Jr., T.F. Terranova, D.D. Lawson, J.Org.Chem. 46, 2379 (1981)
- 12 E. Oliveros, M.-T. Maurette, A.M. Braun, Helv.Chim.Acta 66, 1183 (1983)


FINANCIAL STATEMENT


As of August 30, 1987, SFR 57534.30 have been paid in salaries and chemicals leaving an uncovered balance of SFR 34766.50 since the last payment due June 30, 1987 has not yet been credited.

Lausanne, October 24, 1987



André M. Braun








Figure 3. - Pseudo ternary phase diagram of a mixture of sodium perfluorocaprylate/perfluoro 2-methyl-2-pentanol (1/1), perfluoro n-hexane and water




Figure 4. - Pseudo ternary phase diagram of a mixture of sodium perfluorocaprylate/perfluoro 2-methyl-2-pentanol (1/2), perfluoro n-hexane and water




Figure 5. - Pseudo ternary phase diagram of a mixture of sodium perfluorocaprylate/perfluoro t-butanol (1/2), perfluoro n-hexane and water

DATE
FILMED
58