

AD-A192 660


EFFECTS OF AIR POLLUTION ON HUMAN EXERCISE PERFORMANCE
(U) ARMY RESEARCH INST OF ENVIRONMENTAL MEDICINE NATICK
MA P N FRYKMAN FEB 88


1/1

UNCLASSIFIED

F/G 24/1

NL





MICROCOPY RESOLUTION TEST CHART
NBS 1963-A

1

SECURITY CLASSIFICATION

AD-A192 660

MENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. RI Unclassified		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY SELECTED MAR 21 1988		3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE		4. PERFORMING ORGANIZATION REPORT NUMBER RA	
4. PERFORMING ORGANIZATION REPORT NUMBER		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION US Army Research Institute of Environmental Medicine	6b. OFFICE SYMBOL (If applicable) SGRD-UE-PH	7a. NAME OF MONITORING ORGANIZATION Same as 6a.	
6c. ADDRESS (City, State, and ZIP Code) Kansas St. Natick, MA 01760-5007		7b. ADDRESS (City, State, and ZIP Code)	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code)		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO. 62787A	PROJECT NO. 3M161102BS15
		TASK NO.	WORK UNIT ACCESSION NO. 8450082101 045
11. TITLE (Include Security Classification) Effects of Air Pollution on Human Exercise Performance			
12. PERSONAL AUTHOR(S) Peter N. Frykman			
13a. TYPE OF REPORT Review Article	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, Month, Day) February 1988	15. PAGE COUNT 21
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
		ozone, nitrogen dioxide, peroxyacetyl nitrate, aerosols, sulfur dioxide, carbon monoxide, strength, performance.	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)			
<p>The pollutants commonly experienced in cities of the United States are: carbon monoxide, ozone, peroxyacetyl nitrate, aerosols, sulfur dioxide, and nitrogen dioxide. Only carbon monoxide has been show to reduce exercise performance. The investigations which evaluated the impact of other pollutants on performance, may not have been sensitive enough to detect the small performance decrements caused. Suggested ways to avoid performance decrements are included. <i>Keywords:</i></p>			
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION UNCLAS	
22a. NAME OF RESPONSIBLE INDIVIDUAL Peter N. Frykman		22b. TELEPHONE (Include Area Code) 617-651-4888	22c. OFFICE SYMBOL SGRD-UE-PH

EFFECTS OF AIR POLLUTION ON HUMAN EXERCISE PERFORMANCE

Peter N. Frykman

**Exercise Physiology Division
US Army Research Institute
of
Environmental Medicine
Natick, MA 01760**

88 3 18 055

Introduction and historical perspective

Most of us would like to live, work and exercise in a pristine environment. Sports such as cross country running, skiing, orienteering and kayaking seem to be associated with unpolluted environmental conditions. Unfortunately, the realities of modern life are very often different from this ideal. Most Americans live in or near a metropolitan area. These cities, with their attendant industry, automotive traffic and higher population densities, are large scale pollution sources. It is evident that living in a city can routinely bring us into contact with considerable amounts of air pollution.

Air pollution is not strictly an urban problem. The smog generated by agricultural burning and mining operations can generate staggering amounts of pollution. Natural disasters such as forest fires, dust storms and volcanic eruptions often result in large amounts of airborne contaminants.

In recent history there have been a succession of air pollution catastrophies. A good example of this is the 1952 London episode. At that time in London, coal and coke burning provided the energy for industry and home heating. England's consumption of these fuels resulted in 4.7 million tons of the pollutant sulfur dioxide(SO₂) being discharged into the air over southern England each year(16). In the first week of December 1952, the weather conditions in London left a thick fog trapped beneath a thermal inversion layer. The air was stagnant and the smoke, soot, and sulfur dioxide from the burning of coal and coke built up in the atmosphere. Visibility was so drastically impaired that the public buses were forced to drive with their lights on during the day. The level of SO₂ in the air averaged .7 ppm which is

tion For	
RA&I	<input checked="" type="checkbox"/>
B	<input type="checkbox"/>
nced	<input type="checkbox"/>
cation	

tion/
Availability Codes

Dist	Avail and/or Special
A-1	



5 times the US primary national average air quality standards. The concentration of SO₂ topped out at 1.3 ppm.

Twelve hours after the fog set in and the smog began concentrating, the hospitals in London began admitting extraordinarily large numbers of patients for respiratory problems. The symptoms were a dry cough, sore throat, nasal discharge and sudden bouts of vomiting. Those individuals with a previous history of respiratory trouble were the most seriously

Insert Figure 1 here.

effected. Figure 1 shows the relationship between the increase in atmospheric SO₂ and the death rate. The British Committee on Air Pollution estimates that a total of 4000 extra deaths, over the normal number, occurred from the second of December to the ninth of December 1952. Unfortunately, this episode was not an isolated case. In the winter of 1956 another smog bank covered London, resulting in over 1000 excess deaths due to the exposure to sulfur dioxide and particulates.

The response to these large air pollution catastrophies has been the development of standards for allowable levels of atmospheric pollutants. In the United States, a number of different regional and national standards have been developed. Among these are the Pollutants Standards Index and the National Average Air Quality Standards.

Environmental conditions leading to smog build-up

Two essential factors are needed for a smog problem to develop: a source of pollution and an environmental trap. The pollution source can be exhaust from a vehicle or an industrial source such as a smelter or the dust ejected by a volcanic eruption. In general persistent, chronic

sources are more significant in their effects. None the less, it is hard to ignore the impact of a forest fire or Mt. St. Helens erupting.

Insert Figure 2 here.

An example of a terrain trap is the area around Denver, Colorado. Figure 2 shows that Denver sits at the bottom of a large basin, forming an ideal collecting trap for all of the industrial and automotive pollutants in that area.

Insert Figure 3 here.

Thermal inversion conditions often provide a ceiling that bars the escape of polluted air from a valley or basin. In normal atmospheric conditions (Figure 3a) the air over the earth gets colder with increasing altitude. Under these conditions, the pollutants which are relatively warm, rise through the layers of dense, cold air. Inversion conditions (Figure 3b) occur when a layer of warm air is interposed between layers of colder air. These warm layers can be very stable and can sometimes persist for days. The inversion layer forms a roof that traps the upward movement of airborne pollutants, and it then spreads horizontally beneath this inversion. These elements of a pollution source and an environmental trap are common to almost all situations where there is a build-up of atmospheric pollution.

Pollutants and their effects on performance

Pollutants can be divided into primary and secondary pollutants. Primary pollutants are those that exert their effects as they come

directly from the source. Secondary pollutants are formed by the interaction of primary pollutants with each other, with other compounds or with ultraviolet light. The primary pollutants are: carbon monoxide, sulfur oxides, nitrogen oxides and primary particulates.

From the standpoint of coaches and athletes, carbon monoxide is the most important of the primary pollutants. The principle source of carbon monoxide is automotive exhaust. Carbon monoxide exerts its effect by binding to and blocking the oxygen binding sites on hemoglobin in the red blood cells. Hemoglobin has an affinity for carbon monoxide that is 230 times greater than its affinity for oxygen. This high affinity means that, once the carbon monoxide is bound to hemoglobin, forming carboxyhemoglobin (COHb), it is very difficult to detach it from the hemoglobin molecule(6). By blocking oxygen from attaching to the binding sites on hemoglobin, the ability of blood to carry oxygen is reduced.

Another effect of carbon monoxide binding to hemoglobin is that it causes the remaining binding sites on that particular hemoglobin molecule to develop a higher affinity for oxygen. This makes it more difficult for the hemoglobin to release its' oxygen at the tissues that

Insert Figure 4 here.

need it. Figure 4 demonstrates the results of increased COHb concentrations in the blood. The vertical axis shows the percent saturation of the blood by oxygen, the horizontal axis shows the partial pressure of oxygen in the blood. In blood with increased levels of COHb, the oxygen dissociation curve is shifted to the left. This is an indication of the hemoglobin's elevated affinity for oxygen, which makes

tissue oxygen extraction more difficult. This means that increased levels of carbon monoxide in the blood compromise both the transport of oxygen in the blood, and the extraction of oxygen to the tissues.

The immediate impact this has on exercise performance is that as the concentration of COHb in the blood increases there is a decrement in

Insert Figure 5 here.

maximum oxygen consumption (Figure 5). In 1975 Horvath et al.(9) noted a concentration threshold at 4.3% CO below which there were no decrements in maximum oxygen consumption.

Other effects of increased levels of carboxyhemoglobin are: decreases in peak V_e during maximal exercise(18), decreases in maximal exercise time(3,13), decreases in exercise time before the onset of angina(2) and an increase in submaximal heart rate(18,4)

Another of the primary pollutants are the sulfur oxides. These compounds are the result of incomplete combustion of fossil fuels. Approximately 98% of all atmospheric sulfur is initially sulfur dioxide(11). The effect of sulfur dioxide is to transiently increase the resistance to flow in the upper respiratory tract. This effect is most prevalent in asthmatics but doesn't significantly decrease submaximal exercise performance in normally healthy individuals. There have been no studies at this time that have examined the effects of SO_2 on maximal exercise(12).

Of the pollutants in the nitrogen oxide family, nitrogen dioxide is known to be potentially harmful to man and is the only one that has been extensively studied. The sources of nitrogen dioxide are high

temperature combustion processes, with motor vehicle exhaust accounting for about 40% of the total emissions(11). There have been only a limited number of studies concerning nitrogen dioxide's effects on submaximal performance and there have been no studies addressing the effects of NO₂ on maximal exercise. The studies of submaximal exercise have shown nitrogen dioxide to be a mild upper respiratory tract irritant, but no significant effects on pulmonary function have been found(12).

The inhaling of particulates from soot, cigarette smoke, or dust can serve as respiratory tract irritants causing bronchoconstriction(17). The effects of these irritants on exercise have not been studied to any extent in humans. Particles of different sizes exert their impact at different locations in the respiratory tract. The larger particles are filtered out by the nose and mucous of the upper respiratory tract. The smallest particles find their way down to the alveoli where their action occurs. The body's defends itself against particulate pollutants by changing the depth and pattern of breathing, and by increased mucociliary transport(7,17).

Secondary pollutants are important because they evolve from the interaction of other pollutants, water, salts, and ultraviolet light. The secondary pollutants include ozone, peroxyacetyl nitrate, and various aerosols. The most important of these is ozone. Ozone is formed by the interaction of oxygen, nitrogen dioxide, hydrocarbons, and ultraviolet light. The production of ozone is closely tied to sunlight, and consequently, is highest just after midday(11).

Ozone is a potent airway irritant that can cause reflex bronchoconstriction at high concentrations. Studies of submaximal exercise with exposure to ozone show that the problems encountered are

more associated with discomfort in breathing than in the physiological impairment of the exercise process. It appears that there is no decrement in submaximal exercise performance associated with ozone exposure(12).

With respect to the effects of ozone on maximal exercise very few studies have been completed. Savin and Adams(15) found no change in exercise capacity or cycling $\dot{V}O_2\text{max}$, associated with ozone concentrations. Horvath et al.(8) exposed subjects to different concentrations of ozone. They found no significant changes in treadmill $\dot{V}O_2\text{max}$, HRmax or maximum performance time associated with the different ozone concentrations. The evidence regarding ozone is so limited that it is difficult to draw conclusions.

The primary source of atmospheric peroxyacetyl nitrate is automobile exhaust. This compound is a potent eye and nose irritant and is often responsible for the burning eye sensation that accompanies exposure to smog. Again, there are only a limited number of studies available concerning either submaximal or maximal exercise. The studies that have been completed cite blurred vision, eye irritation and eye fatigue as prominent symptoms of exposure(5). None of the studies have demonstrated clear evidence of respiratory, metabolic, or thermoregulatory distress with peroxyacetyl nitrate exposure during submaximal exercise(12). Studies by Raven et al.(13,14) found no decrement in $\dot{V}O_2\text{max}$ during peroxyacetyl nitrate exposure of various concentrations. These authors speculate that the maximum concentration of peroxyacetyl nitrate used may have been below the threshold at which physiologically meaningful changes may have occurred.

Aerosols are formed by the interaction of various acids and salts. The most commonly studied aerosols are ammonium sulfate, ammonium bisulfate, and ammonium nitrate. These compounds are eye and upper respiratory tract irritants. While they may cause discomfort during either submaximal or maximal exercise, they do not appear to be conclusively linked to exercise performance decrements(7).

In summary, only carbon monoxide shows a clear decremental effect on maximal exercise performance. Other compounds (i.e. ozone, SO₂, NO₂ and peroxyacetyl nitrate) cause irritation of the eyes and respiratory tract, but there is little or no evidence that they cause decrements in either maximal or submaximal exercise performance. There has been a genuine lack of studies dealing with pollutant effects on exercise capacity and performance, especially those that have adequate sample sizes to detect small changes in performance. While it is not clear that some pollutants have significant effects on performance, it makes sense to treat these compounds as potential problems rather than ignore them until it is too late.

Training in smoggy conditions

The key issues to be considered in order to minimize the impact of smog on exercise training can be summed up in four words: TIMING, LOCATION, INTENSITY, DURATION.

Timing

It is important to consider the timing of workouts relative to smog build up. Some pollutants like carbon monoxide (Figure 6a) show rapid increases in concentration that follow the times of peak automotive traffic. Although there is some seasonal change in the overall levels of carbon monoxide, the main changes occur during the course of each day(11).

Insert Figure 6 here.

Other pollutants (i.e. ozone) depend on heat and ultraviolet radiation from the sun for their formation. Figure 6b demonstrates a strong build up of ozone in the early afternoon. The seasonal occurrence of ozone is well known, and summer ozone problems are clearly linked increases in the duration and intensity of solar radiation. In general, the best times of day to avoid smog are the early morning or the late afternoon(11).

Location

This may seem obvious, but athletes should make a conscious effort to avoid exercising near a pollution source. To avoid the exhaust of passing vehicles, run on the upwind side of the road. If forced to stop at an intersection stay away from the exhaust pipes of the stopped vehicles. If necessary stand well away from the intersection to avoid the higher concentration of pollutants that occur in that area. Don't be obsessed by trying to avoid every last bit of pollution, just use your common sense. Also, be aware that air pollution can occur in indoor settings as well as. A classic example of this is a case in which a group of young ice hockey players suffered CO intoxication at an indoor rink. The source of the CO was a gasoline powered ice resurfacing machine(1).

Intensity and Duration

The concept of effective dose is important to understanding the effect of intensity on exercise performance in the smog. The effective dose is the product of the concentration of the pollutant, the volume of

air breathed per minute (which is a reflection of exercise intensity) and the duration of the exercise(10). While all three of these factors affect the pollutant dose during a workout only the duration and the intensity (ventilation volume) can be controlled by the athlete. .

[POLLUTANT CONC.] X (VENTILATION) X (DURATION) = EFFECTIVE DOSE

(A) [1] X (6 liters) X (480 min) = 2880

(B) [1] X (120 liters) X (30 min) = 3600

Equation (A) demonstrates the pollutant dose for a sedentary person having an eight hour exposure to a pollutant of a concentration of 1 and with a minute ventilation of 6 liters. This gives an effective pollutant dose of 2880 units. Equation (B) shows the results of this person exercising for only 30 minutes in the same concentration of pollutants, at a ventilation level of 120 liters per minute they would receive an effective dose of 3600 units or 125% of the sedentary 8 hour dose, in just 30 minutes. It is easy to see how exercise intensity effects the magnitude of a pollutant dose.

Athletes can effectively alter the "intensity" of their workouts by reducing their running speed or by dropping the number of repetitions and sets in their weight workouts. Use the smoggy days for technique workouts rather than for the intensive body building workouts that would demand high ventilation rates. Also weight workouts with high loads (1RM), low repetitions and long rest periods between sets would be appropriate for smoggy day training.

It is also important to limit the duration of exposure to the polluted air. By cutting back on the overall length of exposure and limiting the intensity of the exercise the effective dose of the pollutants can be minimized.

Competing in smoggy conditions

Important international, intercollegiate or interscholastic competitions are typically held in or near large cities. Five of the last seven summer Olympic games were held in cities with histories of high levels of air pollution. Unfortunately the athletes cannot control the atmospheric conditions in which they compete they are stuck with the conditions at hand.

For some sports, the physiological decrements caused by smog exposure may not be substantial. But, for those sports with high metabolic demands, the impact of air pollution can be significant. At the highest levels of competition the difference between victory and defeat is often a tiny fraction of the overall performance. These differences of performance are much smaller than those discernible in most of the air pollution studies conducted to date. Unfortunately, the sample sizes needed to detect such small changes in performance are far larger than those found with the current studies. Studies of a more epidemiological nature are needed to more fully assess the impact of smog on exercise performance, particularly high level athletic performance.

What then are the strategies to help an athlete deal with competing in the smog? First, minimize exposure to pollutants en route to the competition. This seems obvious, but we have all been caught in a traffic jam with the windows of the car open. Another thing to avoid is riding with someone smoking cigars or cigarettes in the car. Secondary cigarette smoke is loaded with carbon monoxide, nicotine and particulates, all of which have detrimental effects on health and performance(11).

Secondly, arrive at the site of the event early. This is most important in sports with high metabolic demands. This is to allow the maximum time to rid oneself of carbon monoxide picked up while travelling. The time needed to washout 50% the CO from the blood ranges from 3 to 4 hours(11).

When you get to the competition site minimize physical activity. By keeping activity low an athlete may cut his/her dose of pollutants prior to the event. The intensity of warm ups should be as low as possible.

Last but not least, avoid pollution sources while competing. Stay away from vehicles on road race courses. Running on the upwind side of a busy road can be a useful strategy for coping with automotive pollutants.

Summary:

First, pay heed to the weather and terrain conditions in the areas where you or people in your charge will be exercising. Evidence shows definite performance decrements only from carbon monoxide. It is quite possible that the studies of other pollutants (i.e. NO₂, SO₂, ozone, peroxyacetyl nitrate, etc.) were not able to detect the small performance decrements that may have been caused by those agents.

Guidelines for training in polluted conditions


- (1) Minimize the impact of air pollution on their training by cutting the intensity and duration of their workouts. This can be done by:
 - (A) Reducing the exercise rate (running, cycling or rowing cadence).
 - (B) Dropping the number of reps and sets in weight workouts.
- (2) Use these times for technique oriented workouts.
- (3) Avoid weight workouts (high repetition, moderate weight, low inter-set rest) or other workouts (intervals, race pace or overpace work) that would demand high ventilation rates.

Strategies for competing in polluted conditions.

- (1) Avoid exposure to pollutants traveling to the event site.
- (2) Arrive at the site of the event early to allow the athletes body to rid itself of as much CO as possible
- (3) Minimize pre-event exercise (including warm-up activity) to limit exposure to polluted air.
- (4) Avoid pollution sources during the event.

Bibliography

1. American Medical Association. (1984) Carbon monoxide intoxication associated with use of a resurfacing machine at an ice-skating rink. *JAMA*, 251:1016.
2. Anderson, E.W., R.J. Andelman, J.M. Strauch, N.J. Fortuin and J.H. Knelson (1973) Effect of a low-level carbon monoxide exposure on onset and duration of angina pectoris: a study in ten patients with ischemic heart disease. *Ann Intern Med*, 79:46-50.
3. Drinkwater, B.L., P.B. Raven, S.M. Horvath, J.A. Gliner, R.O. Ruhling, N.W. Bolduan and S. Taguchi (1974) Air pollution, exercise and heat stress. *Arch Environ Health*, 28:177-181.
4. Ekholm, B. and R. Huot (1972) Response to submaximal and maximal exercise at different levels of carboxyhemoglobin. *Acta Physiol Scand*, 86:474-482.
5. Gliner, J.A., S.M. Horvath, B.L. Drinkwater and J.C. Sutton (1975) Man's physiologic response to thermal and pollutant stress. *J Appl Physiol*, 39:628-632.
6. Guyton, A.C. (1986) *Textbook of Medical Physiology*. 7th edition, Philadelphia, W.B. Saunders.
7. Horvath, S.M. (1981) The Impact of Air Quality in Exercise Performance. pp 265-296 In: *Exercise and Sports Sciences Reviews*, D.I. Miller (Ed.). Philadelphia: The Franklin Institute Press.
8. Horvath, S.M., J.A. Gliner, and J.A. Matsen-Twisdale (1979) Pulmonary function and maximum exercise responses following acute ozone exposure. *Aviat Space Environ Med*, 50:901-905.
9. Horvath, S.M., P.B. Raven, T.E. Dahms and D.J. Gray (1975) Maximal aerobic capacity at different levels of carboxyhemoglobin. *J Appl Physiol*, 38:300-303.
10. Larson, D.C. (1985) Toxic vapor exposure and aerobic exercise. *Physician and Sports Med*, 13:76-83.
11. McCafferty, W.B. (1981) *Air Pollution and Athletic Performance*. Charles C. Thomas Publishers, Springfield, Illinois.
12. Pandolf, K. (In Press) Air Quality and Human Performance, ch 16. In: *Human Performance Physiology and Environmental Medicine at Terrestrial Extremes*, Pandolf K. Sawka M. and Gonzalez R. (Eds) Benchmark Press Inc., Indianapolis, Indiana.
13. Raven, P.B., B.L. Drinkwater, S.M. Horvath, R.O. Ruhling, J.A. Gliner, J.C. Sutton and N.W. Bolduan (1974) Age, smoking habits, heat stress, and their interactive effects with carbon monoxide and peroxyacetylnitrate on man's aerobic power. *Int J Biometeor*, 18:222-232.
14. Raven, P.B., B.L. Drinkwater, R.O. Ruhling, N.W. Bolduan S. Taguchi, J.A. Gliner, and S.M. Horvath. (1974) Effect of carbon monoxide and peroxyacetyl nitrate on man's maximal aerobic capacity. *J Appl Physiol*, 36:288-293.
15. Savin, W.M. and W.C. Adams. (1979) Effects of ozone inhalation on work performance and $\dot{V}O_{2max}$. *J Appl Physiol*, 46:309-314.
16. Shaheen, Esber (1974) *Environmental Pollution: Awareness and Control*, pp 27-64, Engineering Technology Incorporated, Mahomet, Illinois.
17. Widdicombe, J.G., D.C. Kent and J. A. Nadel. (1962) Mechanism of bronchoconstriction during inhalation of dust. *J Appl Physiol*, 17:613-616.
18. Vogel, J.A. and M.A. Gleser. (1972) Effect of carbon monoxide on oxygen transport during exercise. *J Appl Physiol*, 32:234-239.



Days of December, 1952

—●— Excess Deaths —+— SO2

Fig 2




Fig 3




Fig 4







Fig 4

Fig. 6





% CARBOXYHEMOGLOBIN

END

DATE

FILMED

6-1988

DTIC