

AD-A192 713

A HONG-ZAKAI TYPE THEOREM FOR CERTAIN DISCONTINUOUS
SEMI-MARTINGALES. (U) BROWN UNIV PROVIDENCE RI LEFSCHETZ
CENTER FOR DYNAMICAL SYSTE. G FERREYRA JAN 88

1/1


UNCLASSIFIED

LCD5/CCS-88-1 AFOSR-TR-88-0388

F/G 12/1

NL

END
DATE
12/1/88
G. FERREYRA
DTIC



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

②

AD-A192 713

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR :R- 88- U 388	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER DTIC FILE COPY
4. TITLE (and Subtitle) A Wong-Zakai Type Theorem for Certain Discontinuous Semimartingales		5. TYPE OF REPORT & PERIOD COVERED <i>Reprint</i>
7. AUTHOR(s) Guillermo Ferreyra		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Lefschetz Center for Dynamical Systems Division of Applied Mathematics Brown University, Providence, RI 02912		8. CONTRACT OR GRANT NUMBER(s) AFOSR-85-0315
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research Bolling Air Force Base Washington, DC 20332 <i>nm</i>		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F, 2304/A1
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) <i>Same 95 11</i>		12. REPORT DATE January 1988
		13. NUMBER OF PAGES 12
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release: distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) DTIC ELECTE S MAY 02 1988 D <i>OE</i>		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) INCLUDED		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

AFOSR-TR- 88 - 0388


A WONG-ZAKAI TYPE THEOREM FOR
CERTAIN DISCONTINUOUS SEMIMARTINGALES,

by

Guillermo Ferreyra

January 1988

LCDS/CCS #88-1



Lefschetz Center for Dynamical Systems
and
Center for Control Sciences

88 502 142

Division of Applied Mathematics

Brown University Providence RI 02912

**A WONG-ZAKAI TYPE THEOREM FOR
CERTAIN DISCONTINUOUS SEMIMARTINGALES,**

by

Guillermo Ferreyra

January 1988

LCDS/CCS #88-1

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



**A WONG-ZAKAI TYPE THEOREM FOR CERTAIN
DISCONTINUOUS SEMIMARTINGALES**

by

Guillermo Ferreyra

**Lefschetz Center for Dynamical Systems
Division of Applied Mathematics
Brown University
Providence, RI 02912**

July 1987

On leave from the Department of Mathematics, Louisiana State University,
Baton Rouge, LA 70803.

Research partially supported by the AFOSR under contract #AFOSR-85-0315, the
ARO under contract #DAAG29-84-K-0082 and #DAAL03-86-K-0171, and by the
Louisiana Education Quality Support Fund under contract #86-LBR-016-04.

Abstract

Solutions of stochastic differential equations having differentials of bounded variation processes on the right hand side can be defined by means of Lebesgue-Stieltjes integrals or by continuous extension of Stieltjes integrals. Both solutions are compared here and formulas that extend the Wong-Zakai theorem are obtained.

Key words and phrases: stochastic equations, Stratonovich integration, Wong-Zakai Theorem.

§1. Introduction

Solutions of stochastic differential equations having differentials of bounded variation processes on the right hand side are usually defined by means of pathwise Lebesgue-Stieltjes integrals (see, for example, Meyer [2]). Partially motivated by stochastic control problems we have introduced a different idea of solution in Ferreyra [1]. With this notion, the change of variables formula for solutions of stochastic differential equations driven by certain semimartingales is not complicated by the jumps of the processes involved. In fact, our interpretation is an extension of the concept of solution of stochastic differential equations in Stratonovich sense, together with the usual change of variables formula. Moreover, robustness in the driving process is built into this definition of solution. Hence results complementary to those considering approximation of driving martingales such as in Protter [4] and Picard [3] follow.

In this paper we obtain formulas relating both definitions of solutions. These formulas are in the vein of the theorems of Wong-Zakai [5].

We introduce the necessary notation and definitions in Section 2 along with some material from Ferreyra [1]. We explain the goal of this paper once more at the end of Section 2. In Section 3 a simple example is presented. Finally, Section 4 gives several formulas of the Wong-Zakai type.

§2. Notation, prerequisites and hypotheses

Let $(\Omega, \mathcal{F}, P)$ be a complete probability space together with an increasing family of sub σ -algebras $\mathcal{F}_t$, $0 \leq t \leq T < \infty$, such that $\mathcal{F}_0$ contains all P -null elements of $\mathcal{F}$, and $\mathcal{F}_t$, $0 \leq t \leq T$, is right-continuous, that is, $\mathcal{F}_t = \mathcal{F}_{t+} = \bigcap_{t < s \leq T} \mathcal{F}_s$.

for all t .

Two processes $X(t)$, and $Y(t)$, $0 \leq t \leq T$, are identified if they are *indistinguishable*, i.e., for almost all $\omega \in \Omega$ the equality $X(t, \omega) = Y(t, \omega)$ holds for all t . A process with paths which are continuous on the right (respectively, left) and have limits on the left (respectively, right) will be called *corlol* (respectively, *collor*). Other authors use the French versions *cadlag* and *caglad* respectively. We assume that all martingales (but not all processes) are *corlol*. If $X(t)$, $0 \leq t \leq T$ is a *corlol* or a *cadlag* process, then $\Delta X(t)$ denotes the jump $X(t^+) - X(t^-)$ at t . Two σ -fields on $[0, T] \times \Omega$ are of importance to us. The *optional* σ -field Σ_0 which is generated by the family of all adapted *corlol* processes and the *predictable* σ -field Σ_p which is generated by the adapted *collor* processes. A process $X(t)$, $0 \leq t \leq T$, is said to be *optional* (respectively, *predictable*) if it is Σ_0 -measurable (respectively, Σ_p -measurable). A process is said to be of *bounded variation* if it is adapted, *corlol*, and it has paths of bounded variation. A process of bounded variation $A(t)$, $0 \leq t \leq T$, is said to be of *integrable variation* if $E \int_0^T |dA(s)| < \infty$. It is assumed that $A(t) = 0$, $t < 0$. If $A(t)$, $0 \leq t \leq T$ is of integrable variation and $H(t)$, $0 \leq t \leq T$, is an optional process such that $E \int_0^\infty |H(s)| |dA(s)| < \infty$, then the *stochastic integral* $I(t) = \int_0^t H(s) dA(s)$ is well defined as a pathwise Lebesgue-Stieltjes integral (cf. Meyer [2], p.258). The process $I(t)$, $0 \leq t \leq T$, turns out to be adapted, continuous on the right and of integrable variation. An adapted process $M(t)$, $0 \leq t \leq T$, vanishing at zero is called a *local martingale* if there exist stopping times $T_n \uparrow T$ such that the stopped process $M^{T_n}(t) = M(t \wedge T_n)$ are uniformly integrable martingales. An adapted process $Z(t)$ is a *semimartingale* if it admits a decomposition of the

form $Z(t) = Z(0) + M(t) + A(t)$, where $M(t)$ is a local martingale vanishing at zero, and $A(t)$ is a process of bounded variation vanishing at zero. If $Z(t)$ is a semimartingale, $Z^c(t)$ will denote its *continuous part*. If $M(t)$ is a square integrable martingale we let $\langle M, M \rangle(t)$ denote the unique *increasing predictable process* such that $\langle M, M \rangle(0) = M^2(0)$ and $M^2(t) - \langle M, M \rangle(t)$ is a martingale. If $M(t)$ and $N(t)$ are both square integrable martingales, then $\langle M, N \rangle(t) = \frac{1}{2}(\langle M + N, M + N \rangle(t) - \langle M, M \rangle(t) - \langle N, N \rangle(t))$ is the unique *predictable process of integrable variation* such that $\langle M, N \rangle(0) = M(0)N(0)$ and $M(t)N(t) - \langle M, N \rangle(t)$ is a martingale. We are ready now to state Ito's rule for semimartingales.

Theorem (Meyer [2], p. 301): Let $Z(t)$ be an $\mathbb{R}^n$ -valued process such that each one of its components $Z^i(t)$, $i = 1, \dots, n$, is a semimartingale. Let $F \in C^2(\mathbb{R}^n)$. Then $F(Z(t))$ is a semimartingale and

$$\begin{aligned}
 (1) \quad F(Z(t)) &= F(Z(0)) + \sum_{i=1}^n \int_0^t \frac{\partial F}{\partial x_i} (Z(s^-)) dZ^i(s) \\
 &+ \frac{1}{2} \sum_{i,j=1}^n \int_0^t \frac{\partial^2 F}{\partial x_i \partial x_j} (Z(s^-)) d\langle Z^{ic}, Z^{jc} \rangle(s) \\
 &+ \sum_{0 \leq s \leq t} (F(Z(s)) - F(Z(s^-)) - \sum_{i=1}^n \frac{\partial F}{\partial x_i} (Z(s^-)) \Delta Z^i(s)).
 \end{aligned}$$

Next, we describe results obtained in Ferreyra [1]. The meaning of the expression

$$(2) \quad dX(t) = f(X(t)) \circ du(t) + g(X(t))dt + \sum_{v=1}^J \sigma_v(X(t)) \circ dW^v(t)$$

will be given below when we define what is meant by solving (2) with initial condition

$$(3) \quad X(0) = X.$$

Let the process $(W^1(t), \dots, W^J(t))$, $0 \leq t \leq T$, be a given J -dimensional Brownian motion. Assume that $u(t)$, $0 \leq t \leq T$, belongs to the set U of real valued, uniformly bounded, collor processes. Assume, without loss of generality, that all processes $v \in U$ satisfy $v(0) = 0$. Assume that $u(t) = 0$, for $t < 0$. The unknown process $X(t)$, $0 \leq t \leq T$, has values in $\mathbb{R}^n$. Furthermore, assume

$$(H1) \quad E|X|^p < \infty, \text{ for some } p > 2,$$

$$(H2) \quad f \in C^3(\mathbb{R}^n), f_x \in C_b^2(\mathbb{R}^n),$$

$$(H3) \quad g \in C_b^1(\mathbb{R}^n), \text{ and}$$

$$(H4) \quad \sigma_v \in C_b^2(\mathbb{R}^n), v = 1, \dots, J.$$

Let $\mathcal{Z}$ denote the set of adapted real valued processes $v(t)$, $0 \leq t \leq T$, having Lipschitz paths with a uniform Lipschitz constant. If $u(t) = v(t) \in \mathcal{Z}$, then $X(t)$ is said to solve (2) if

$$(4) \quad dX(t) = f(X(t)) \frac{dv}{dt}(t)dt + g(X(t))dt + \sum_{v=1}^J \sigma_v(X(t)) \circ dW^v(t)$$

in Stratonovich sense. It is well known that for such $v(t) \in \mathcal{Z}$, the problem (3) - (4) has a unique solution. This concept of solution is extended to allow all $u \in U$ as follows.

Definition 1: An $\mathbb{R}^n$ -valued process $X(t)$, $0 \leq t \leq T$, is said to be a solution of (2) - (3) if there exists a map $\Gamma : [0, T] \times U \times \Omega \rightarrow \mathbb{R}^n$ such that

(D1) for each $v \in U$, $\Gamma(t, v)$, $0 \leq t \leq T$, is collor,

- (D2) if $v \in \mathfrak{Z}$, then the process $\Gamma(t, v)$ solves (4) in Stratonovich sense,
 (D3) if $v \in U$ and $\{v_j\}$ is a uniformly bounded sequence of elements in U such that for every t , $0 \leq t \leq T$, $v_j(t) \rightarrow v(t)$, a.s., then for each t ,

$$E|\Gamma(t, v_j) - \Gamma(t, v)|^2 \rightarrow 0, \text{ as } j \rightarrow \infty,$$

 (D4) $\Gamma(t, u) = X(t)$, $0 \leq t \leq T$, and
 (D5) for all $v \in U$, $\Gamma(0, v) = X$.

The extension from $\mathfrak{Z}$ to U is aided by the following.

Lemma 1 (Ferreyra [1]): Let $u \in U$, and let $u_j(t) = j \int_{t-1/j}^t u(s) ds$. Then $\{u_j\}$ is a uniformly bounded sequence of elements in $\mathfrak{Z}$ such that for each t , $0 \leq t \leq T$, $u_j(t) \rightarrow u(t)$, a.s. as $j \rightarrow \infty$.

Given $u \in U$, the problem (2) - (3) is shown to have a unique solution in Ferreyra [1]. We sketch here the proof of existence. Let $F : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ be the flow of f , that is, the solution of $\frac{\partial F}{\partial s}(s, x) = f(F(s, x))$, $F(0, x) = x$. Let $\bar{g}(s, x) = \left[\frac{\partial F}{\partial x}(s, x) \right]^{-1} g(F(s, x))$, and $\bar{\sigma}_v(s, x) = \left[\frac{\partial F}{\partial x}(s, x) \right]^{-1} \sigma_v(F(s, x))$. Then Γ is defined by

$$(5) \quad \Gamma(t, v) = F(v(t), Y(t)),$$

where $Y(t)$ is the process with continuous paths satisfying

$$(6) \quad dY(t) = \bar{g}(v(t), Y(t))dt + \sum_{v=1}^J \bar{\sigma}_v(v(t), Y(t)) \circ dW^v(t),$$

$$(7) \quad Y(0) = X.$$

The proof of (D2) follows easily by application of the rule for change of variables for Stratonovich integrals. The proof of (D3) is a little more complicated. Basically, it involves estimates for F , its partial derivatives of

first order, and estimates for $Y(t)$. The reader is referred to Ferreyra [1] for more details.

Finally, we restate the goal of this paper. We intend to find an expression for $X(t)$, the solution of (2) - (3), in terms of integrals of Lebesgue-Stieltjes and Itô type such as those appearing in (1).

§3. A simple example

To clarify the relation between (1) and (2) we consider the following deterministic example. Let $T = 3$, $u^- = 1_{(1,3]} + 1_{(2,3]}$, and $u^+ = 1_{[1,3]} + 1_{[2,3]}$, where for $A \subset [0,3]$, 1_A denotes the characteristic function of the set A . Consider the 2-dimensional system of the type of (2) - (3)

$$\begin{aligned} dX^1(t) &= 1 \circ du^-(t), & X^1(0) &= 0, \\ dX^2(t) &= \psi(X^1(t)) \circ du^-(t), & X^2(0) &= 0, \end{aligned}$$

where $\psi(x) = 3x^2$.

The solution of this system is computed using Definition 1 as follows. Let $u_j(t) = j(t-1 + 1/j)1_{(1-1/j,1]}(t) + j(t-2 + 1/j)1_{(2-1/j,2]}(t) + u^-(t)$. Then solve by standard methods

$$\begin{aligned} dX_j^1(t) &= \frac{du_j}{dt}(t) dt, & X_j^1(0) &= 0, \\ dX_j^2(t) &= \psi(X_j^1(t)) \frac{du_j}{dt}(t) dt, & X_j^2(0) &= 0. \end{aligned}$$

Finally, take the limit as $j \rightarrow \infty$. Hence

$$\begin{aligned} X_j^1(t) &= u_j(t), & X_j^2(t) &= \int_0^t \psi(u_j(s)) du_j(s) = [u_j(t)]^3, \\ X^1(t) &= u(t), & X^2(t) &= [u^-(t)]^3. \end{aligned}$$

On the other hand the 2-dimensional system of the type of (1) - (3)

$$dY^1(t) = du^+(t), \quad Y^1(0) = 0$$

$$dY^2(t) = \psi(Y^1(t^-))du^+(t), \quad Y^2(0) = 0$$

is solved by means of Lebesgue-Stieltjes integrals as

$$Y^1(t) = u^+(t),$$

$$Y^2(t) = \int_0^t \psi(u^+(s^-))du^+(s) = \int_0^t 3[u^+(s^-)]^2 du^+(s).$$

Thus $X^2(t)$ jumps 1 unit at $t = 1$ and 7 units at $t = 2$, while $Y^2(t)$ jumps 0 units at $t = 1$ and 3 units at $t = 2$. The difference between $X^2(t)$ and $Y^2(t)$ can be deduced from (1). In fact,

$$X^2(t) - Y^2(t) = \sum_{0 < s \leq t} \left\{ [u^+(s)]^3 - [u^+(s^-)]^3 - 3[u^+(s^-)]^2 \Delta u^+(s) \right\} - \Delta X^2(t)$$

$$= \sum_{0 < s \leq t} \left\{ X^2(s^+) - X^2(s) - 3[X^2(s)]^2 \Delta u^-(s) \right\} - 3[X^2(t)]^2 \Delta u^-(t).$$

We will prove below several generalizations of this formula.

§4. Stochastic integrals and stochastic differential equations

In the previous section we found a formula relating $X^2(t) = \lim_{j \rightarrow \infty} \int_0^t \psi(u_j(s))du_j(s)$ and the Lebesgue-Stieltjes integral $Y^2(t) = \int_0^t \psi(u^+(s^-))du^+(s)$. More general situations are treated here.

We assume throughout this section that $u(t)$, $0 \leq t \leq T$, is a fixed process in U such that it has paths of bounded variation. Hence the process $u(t^+)$, $0 \leq t < T$, is well defined and it is of bounded variation. Let $u_j(t)$, $j = 1, 2, \dots$, $0 \leq t \leq T$, be a sequence of processes in $\mathcal{Z}$ approximating $u(t)$ in the sense of Lemma 1, that is, for each $0 \leq t \leq T$, $u_j(t)$ converges to $u(t)$, a.s. as $j \rightarrow \infty$.

Theorem 1: Let $\varphi = \varphi(t, x)$ be a real valued function in $C^1([0, T] \times \mathbb{R})$. Then, for each $0 \leq t \leq T$,

$$\begin{aligned} \lim_{j \rightarrow \infty} \int_0^t \varphi(s, u_j(s)) du_j(s) &= \psi(0, u(0^+)) + \int_{0^+}^t \varphi(s, u(s)) du(s^+) \\ &+ \sum_{0 < s < t} [\psi(s, u(s^+)) - \psi(s, u(s)) - \varphi(s, u(s)) \Delta u(s)] - \varphi(t, u(t)) \Delta u(t), \end{aligned}$$

where $\psi(t, x) = \int_0^x \varphi(t, \xi) d\xi$.

Proof: Let ψ be defined as above. Then, by the calculus for Riemann-Stieltjes integrals

$$\psi(t, u_j(t)) = \int_0^t \varphi(s, u_j(s)) du_j(s) + \int_0^t \frac{\partial \psi}{\partial t}(s, u_j(s)) ds.$$

Since for each t , $0 \leq t \leq T$, $u_j(t) \rightarrow u(t)$, a.s. as $j \rightarrow \infty$, then our hypotheses on φ and the Dominated Convergence Theorem imply

$$(8) \quad \psi(t, u(t)) = \lim_{j \rightarrow \infty} \int_0^t \varphi(s, u_j(s)) du_j(s) + \int_0^t \frac{\partial \psi}{\partial t}(s, u(s)) ds.$$

But (1) with $n = 2$, $Z^1(t) = t$, $Z^2(t) = u(t^+)$ imply

$$\begin{aligned} (9) \quad \psi(t, u(t^+)) &= \psi(0, u(0^+)) + \int_{0^+}^t \varphi(s, u(s)) du(s^+) + \int_0^t \frac{\partial \psi}{\partial t}(s, u(s)) ds \\ &+ \sum_{0 < s < t} [\psi(s, u(s^+)) - \psi(s, u(s)) - \varphi(s, u(s)) \Delta u(s)]. \end{aligned}$$

Comparison of (8) and (9) prove the desired formula.

The above theorem is further generalized as follows. Let $\alpha, \beta_v \in C([0, T])$, $v = 1, \dots, J$, and define

$$S_j(t) = \int_0^t \alpha(s) du_j(s) + \sum_{v=1}^J \int_0^t \beta_v(s) \circ dW^v(s).$$

Here α and β_ν are deterministic functions, the first integral is pathwise of Riemann-Stieltjes type and the integrals with respect to W^ν are of Stratonovich (equal to Itô in this case) type. Let $\varphi = \varphi(t, x) \in C^2([0, T] \times \mathbb{R})$ and consider the stochastic integral

$$\int_0^t \varphi(s, S_j(s)) \circ dS_j(s) = : \int_0^t \varphi(s, S_j(s)) \alpha(s) du_j(s) + \sum_{\nu=1}^J \int_0^t \varphi(s, S_j(s)) \beta_\nu(s) \circ dW^\nu(s).$$

Finally, define the (collor) process

$$S(t) = \int_0^t \alpha(s) du(s) + \sum_{\nu=1}^J \int_0^t \beta_\nu(s) dW^\nu(s),$$

where the first integral is of Riemann-Stieltjes type and the other J integrals are of Itô (equal to Stratonovich in this case) type.

Theorem 2: Let $\psi(t, x) = \int_0^x \varphi(t, \xi) d\xi$. Then, for each $0 \leq t \leq T$, the following relation holds.

$$\lim_{j \rightarrow \infty} \int_0^t \varphi(s, S_j(s)) \circ dS_j(s) = \psi(0, S(0^+)) + \int_{0^+}^t \varphi(s, S(s)) dS(s^+) + \frac{1}{2} \sum_{\nu=1}^J \int_0^t \frac{\partial \varphi}{\partial x}(s, S(s)) \beta_\nu^2(s) ds + \sum_{0 \leq s < t} [\psi(s, S(s^+)) - \psi(s, S(s)) - \varphi(s, S(s)) \Delta S(s)] - \varphi(t, S(t)) \Delta S(t).$$

Proof: By the calculus for Stratonovich integrals

$$\psi(t, S_j(t)) = \int_0^t \varphi(s, S_j(s)) \circ dS_j(s) + \int_0^t \frac{\partial \psi}{\partial t}(s, S_j(s)) ds.$$

Theorem 1 implies that

$$\lim_{j \rightarrow \infty} \int_0^t \alpha(s) du_j(s) = \alpha(0)u(0^+) + \int_{0^+}^t \alpha(s) du(s^+) - \alpha(t)\Delta u(t) = \int_0^t \alpha(s) du(s).$$

Then, for each $0 \leq t \leq T$, $S_j(t) \rightarrow S(t)$, a.s. as $j \rightarrow \infty$. Hence the Dominated Convergence Theorem implies

$$(10) \quad \psi(t, S(t)) = \lim_{j \rightarrow \infty} \int_0^t \varphi(s, S_j(s)) \circ dS_j(s) + \int_0^t \frac{\partial \psi}{\partial t}(s, S(s)) ds.$$

On the other hand, (1) with $n = 2$ and $Z^1(t) = t$, $Z^2(t) = S(t^+)$ give

$$(11) \quad \begin{aligned} \psi(t, S(t^+)) &= \psi(0, S(0^+)) + \int_{0^+}^t \varphi(s, S(s)) dS(s^+) + \frac{1}{2} \sum_{v=1}^J \int_0^t \frac{\partial \varphi}{\partial x}(s, S(s)) \beta_v^2(s) ds \\ &\quad + \int_0^t \frac{\partial \psi}{\partial t}(s, S(s)) ds + \sum_{0 < s \leq t} [\psi(s, S(s^+)) - \psi(s, S(s)) - \varphi(s, S(s)) \Delta S(s)]. \end{aligned}$$

Putting (10) and (11) together we obtain our result.

Theorem 3: Let $X(t)$, $0 \leq t \leq T$, be the solution of (2) - (3) in the sense of Definition 1. Then

$$(12) \quad \begin{aligned} X(t) &= X + \int_{0^+}^t f(X(s)) du(s^+) + \int_0^t g(X(s)) ds + \sum_{v=1}^J \int_0^t \sigma_v(X(s)) dW^v(s) \\ &\quad + \frac{1}{2} \sum_{i=1}^n \sum_{v=1}^J \int_0^t \frac{\partial \sigma_v}{\partial x_i}(X(s)) \sigma_v^i(X(s)) ds + \sum_{0 \leq s < t} \Delta X(s) - \sum_{0 < s \leq t} f(X(s)) \Delta u(s). \end{aligned}$$

Proof: The solution of (2) - (3) can be expressed, according to (5)- (7), as

$$(13) \quad X(t) = F(u(t), Y(t)),$$

where F is the flow of f and $Y(t)$, $0 \leq t \leq T$, is the continuous semimartingale solution of $Y(0) = X$, $dY(t) = \bar{g}(u(t), Y(t))dt + \sum_{v=1}^J \bar{\sigma}_v(u(t), Y(t)) \circ dW^v(t)$. As indicated at the beginning of this section, the process $u(t^+)$, $0 \leq t < T$, is of bounded variation. Then, it follows from (1)

that

$$\begin{aligned} X(t^+) &= F(u(0^+), Y(0)) + \int_{0^+}^t \frac{\partial F}{\partial s}(u(s), Y(s)) du(s^+) + \sum_{i=1}^n \int_0^t \frac{\partial F}{\partial x_i}(u(s), Y(s)) dY^i(s) \\ &+ \frac{1}{2} \sum_{i,j=1}^n \int_0^t \frac{\partial^2 F}{\partial x_i \partial x_j}(u(s), Y(s)) d\langle Y^i, Y^j \rangle(s) \\ &+ \sum_{0 < s \leq t} [F(u(s^+), Y(s)) - F(u(s), Y(s)) - \frac{\partial F}{\partial s}(u(s), Y(s)) \Delta u(s)]. \end{aligned}$$

The following equalities are used to replace the various terms in the above equation. It is easy to see that

$$\begin{aligned} F(u(s^+), Y(s)) - F(u(s), Y(s)) &= \Delta X(s), \\ F(u(0^+), Y(0)) &= X + \Delta X(0), \\ X(t^+) &= X(t) + \Delta X(t), \\ \frac{\partial F}{\partial s}(u(s), Y(s)) &= f(F(u(s), Y(s))) = f(X(s)), \\ \frac{\partial F}{\partial x}(u(s), Y(s)) \bar{g}(u(s), Y(s)) &= g(X(s)), \\ \frac{\partial F}{\partial x}(u(s), Y(s)) \bar{\sigma}_\nu(u(s), Y(s)) &= \sigma_\nu(X(s)), \end{aligned}$$

and

$$\begin{aligned} \sum_{i=1}^n \frac{\partial F}{\partial x_i}(u(s), Y(s)) \sum_{\nu=1}^J \sum_{j=1}^n \frac{\partial \bar{\sigma}_\nu^i}{\partial x_j}(u(s), Y(s)) \bar{\sigma}_\nu(u(s), Y(s)) &= \\ &= - \sum_{i,j=1}^n \frac{\partial^2 F}{\partial x_i \partial x_j}(u(s), X(s)) d\langle Y^i, Y^j \rangle(s) + \sum_{\nu=1}^J \sum_{i=1}^n \frac{\partial \sigma_\nu}{\partial x_i}(X(s)) \sigma_\nu^i(X(s)). \end{aligned}$$

In the last equality we used the identity obtained from differentiation of $F_x(s, x)F_x(s, x)^{-1} = I$ with respect to x . The proof of (12) is then concluded.

Acknowledgement: The author wishes to thank P. Dupuis for his valuable comments.

References

- [1] G. Ferreyra, Approximation of stochastic equations driven by predictable processes, submitted for publication.
- [2] P.A. Meyer, Un cours sur les integrales stochastiques, Seminaire de Probabilites X, Lecture Notes in Math. 551, Springer-Verlag, NY (1976) 245-400.
- [3] J. Picard, Approximation of stochastic differential equations and application of the stochastic calculus of variations to the rate of convergence, preprint.
- [4] P. Protter, Approximation of solutions of stochastic differential equations driven by semimartingales, Ann. of Probability 13 (1985), pp. 716-743.
- [5] E. Wong and M. Zakai, On the convergence of ordinary integrals to stochastic integrals, Ann. Math. Stat. 36 (1965) pp. 1560-1564.

END

DATE

FILMED

6-1988

DTIC