

AD-A192 841

A NOTE ON VECTOR BIMEASURES(U) NORTH CAROLINA UNIV AT
CHAPEL HILL CENTER FOR STOCHASTIC PROCESSES C HOUDRE
NOV 87 TR-214 AFOSR-TR-88-0349 F49620-85-C-0144


1/1

UNCLASSIFIED

F/G 12/3

NL





MICROCOPY RESOLUTION TEST CHART
NBS 1963-A

(2)


AD-A192 841

REPORT DOCUMENTATION PAGE

2a. SECURITY CLASSIFICATION AUTHORITY			1b. RESTRICTIVE MARKINGS FILE COPY		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited.		
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. 214			5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 88 - 0349		
6a. NAME OF PERFORMING ORGANIZATION University of North Carolina		6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION AFOSR/NM		
6c. ADDRESS (City, State, and ZIP Code) Statistics Dept. 321-A Phillips Hall 039-A Chapel Hill, NC 27514			7b. ADDRESS (City, State, and ZIP Code) AFOSR/NM Bldg 410 Bolling AFB DC 20332-6448		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION AFOSR		8b. OFFICE SYMBOL (if applicable) NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR No. F49620 85C 0144.		
8c. ADDRESS (City, State, and ZIP Code) AFOSR/NM Bldg 410 Bolling AFB DC 20332-6448			10. SOURCE OF FUNDING NUMBERS		WORK UNIT ACCESSION NO.
			PROGRAM ELEMENT NO. 61102F	PROJECT NO. 2304	TASK NO. AS
11. TITLE (Include Security Classification) A note on vector bimeasures					
12. PERSONAL AUTHOR(S) Houdre, Christian					
13a. TYPE OF REPORT preprint		13b. TIME COVERED FROM 9/87 TO 8/88		14. DATE OF REPORT (Year, Month, Day) November 1987	
15. PAGE COUNT 6					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Key Words & Phrases: N/A		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) A Fubini type theorem is obtained for vector bimeasure integrals.					
DTIC ELECTE APR 04 1988 S D					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT ☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT. ☐ DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified/unlimited		
22a. NAME OF RESPONSIBLE INDIVIDUAL Maj. Brian Woodruff			22b. TELEPHONE (Include Area Code) (202) 767-5026		22c. OFFICE SYMBOL NM

CENTER FOR STOCHASTIC PROCESSES

Department of Statistics
University of North Carolina
Chapel Hill, North Carolina



A NOTE ON VECTOR BIMEASURES

by

Christian Houdré

Technical Report No. 214

November 1987

A NOTE ON VECTOR BIMEASURES

Christian Houdré*
Center for Stochastic Processes
Department of Statistics
University of North Carolina at
Chapel Hill

Accession For	
NTIS CRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Abstract A Fubini type theorem is obtained for vector bimeasure integrals.

AMS (1980) subject classification: Primary 28B05; Secondary 60G12

*Supported in part by AFOSR Grant No. F49620 85 C 0144.

Morse and Transue [6-8] initiated the development of a theory of integration with respect to a bimeasure which was subsequently studied by Thomas [10]. For these authors, bimeasures are continuous bilinear functionals on $C_c(E_1) \times C_c(E_2)$, where $C_c(E_i)$, $i=1,2$, are the usual spaces of continuous functions with compact support on the locally compact Hausdorff spaces E_i , $i=1,2$. More recently, motivated by the problem of finding a Fourier representation for the covariance of a second order process, this theory has been expanded by Niemi [9] and Chang and Rao [2]. Both the bilinear functional and the set function approaches have now been studied as well as the Banach valued case developed by Ylinen [11].

In the works mentioned above the authors consistently impose, in their definition of integrability, a Fubini type condition which cannot usually be bypassed. The purpose of this note is to show that under a suitable restriction of the definition of integrability, the Fubini-type requirement becomes obsolete.

Let X be a Banach space over $F = \mathbb{R}$ or $\mathbb{C}$ and let $(E, \mathcal{M})$ be a measurable space. A vector measure is a σ -additive set function $\mu: \mathcal{M} \rightarrow X$. Integration of functions $f: E \rightarrow F$ with respect to vector measures is taken in the Bartle-Dunford and Schwartz [1] sense, the reader being referred to Dunford and Schwartz [3 IV 10] for the properties of this vector integral.

Let $(E_1, \mathcal{M}_1)$ and $(E_2, \mathcal{M}_2)$ be two measurable spaces. A vector bimeasure (bimeasure when $X = F$) is a separately σ -additive set function $\beta: \mathcal{M}_1 \times \mathcal{M}_2 \rightarrow X$, i.e., $\beta(\cdot, B)$ and $\beta(A, \cdot)$ are vector measures for all $A \in \mathcal{M}_1$, $B \in \mathcal{M}_2$.

The proof of our result as well as our definition of integrability will rely on the following two lemmas. The first one is classical and can be

found in [3 p. 323] while the second is in [11].

Lemma 1 Let $f: E \rightarrow F$ be μ -integrable. Then, the set function $\nu(A) = \int_A f d\mu$, $A \in \mathcal{M}$ is a vector measure.

Lemma 2 Let $f: E_1 \rightarrow F$ be $\beta(\cdot, B)$ -integrable for all $B \in \mathcal{M}_2$. Then the set functions ${}_f\beta(A, \cdot): \mathcal{M}_2 \rightarrow X$, $B \rightarrow {}_f\beta(A, \cdot)(B) = \int_A f d\beta(\cdot, B)$ are vector measures for all $A \in \mathcal{M}_1$.

In the above and for $g: E_2 \rightarrow F$ the vector measures $\beta_g(\cdot, B)$ can be obtained in a completely symmetrical way.

We can now define integrability.

Definition 3 A pair of functions (f, g) , $f: E_1 \rightarrow F$, $g: E_2 \rightarrow F$ is said to be integrable with respect to the vector bimeasure $\beta: \mathcal{M}_1 \times \mathcal{M}_2 \rightarrow X$ (β -integrable for short) if the following two conditions hold.

- (i) f is $\beta(\cdot, B)$ -integrable for all $B \in \mathcal{M}_2$ and g is $\beta(A, \cdot)$ -integrable for all $A \in \mathcal{M}_1$.
- (ii) f is $\beta_g(\cdot, B)$ -integrable for all $B \in \mathcal{M}_2$ and g is ${}_f\beta(A, \cdot)$ -integrable for all $A \in \mathcal{M}_1$.

Remark 4 For $X = F$ our definition of integrability is stronger than that of Morse and Transue. For these authors, (f, g) is integrable if in (i) and (ii) A and B are replaced by E_1 and E_2 and if in addition

$$\int_{E_1} f d\beta_g(\cdot, E_2) = \int_{E_2} g d{}_f\beta(E_1, \cdot). \quad (1)$$

It is also more restrictive than the strong integral of Niemi or the

β -integral of Ylinen. For both of them, a pair (f, g) is integrable if in (ii) A and B are respectively replaced by E_1 and E_2 and if in addition (1) is satisfied.

However, our definition is weaker than the strict β -integral of Chang and Rao (there is no Borel assumption on f and g or the additional Fubini condition).

As already mentioned, with other definitions of integrability, (1) cannot be bypassed (see [8], [11]). However, with Definition 3, this condition will always hold.

Theorem 5 Let the pair (f, g) be β -integrable, then

$$\int_A f d\beta_g(\cdot, B) = \int_B g d_f\beta(A, \cdot), \quad \forall A \in \mathcal{M}_1, B \in \mathcal{M}_2. \quad (2)$$

The common value in (2) can thus be denoted by $\int_A \int_B f g d\beta$.

Proof Let (f, g) be β -integrable. If both f and g are simple functions, then (2) is trivial. Let f and g be bounded (f and g are measurable since integrable in the Bartle-Dunford and Schwartz sense). Then, f and g are uniform limits of simple functions and by the dominated convergence theorem for vector measures (see [3 p. 328]) (2) is again true.

Let f be bounded and let $B_n = \{y \in B \mid |n| \leq |g| < n+1\}$.

$$\begin{aligned} \text{Then,} \quad \int_B g d_f\beta(A, \cdot) &= \sum_{n=0}^{\infty} \int_{B_n} g d_f\beta(A, \cdot) && (\text{Lemma 1}) \\ &= \sum_{n=0}^{\infty} \int_A f d\beta_g(\cdot, B_n) && (g \text{ is bounded on } B_n) \end{aligned}$$

$$= \int_A f d\beta_g(\cdot, B) \quad (\text{Lemma 2}).$$

If f is not bounded, then $A = \bigcup_{n=0}^{\infty} A_n$ with f bounded on each A_n . Hence,

$$\begin{aligned} \int_A f d\beta_g(\cdot, B) &= \sum_{n=0}^{\infty} \int_{A_n} f d\beta_g(\cdot, B) \quad (\text{Lemma 1}) \\ &= \sum_{n=0}^{\infty} \int_B g d_f \beta(A_n, \cdot) \quad (f \text{ is bounded on } A_n) \\ &= \int_B g d_f \beta(A, \cdot) \quad (\text{Lemma 2}). \end{aligned}$$

and the result is obtained.

Remark 6 Let $X : \mathbb{R} \rightarrow L^2(\Omega, \mathfrak{G}, P)$ be a continuous V -bounded process, i.e.,

$$X_t = \int_{\mathbb{R}} e^{itx} d\mu(x), \quad t \in \mathbb{R}, \text{ for some vector measure } \mu : \mathfrak{G}(\mathbb{R}) \rightarrow L^2(\Omega, \mathfrak{G}, P).$$

($\mathfrak{G}(\mathbb{R})$ is the Borel σ -algebra of $\mathbb{R}$). Then $\beta(A, B) = E\mu(A)\overline{\mu(B)}$, $A, B \in \mathfrak{G}(\mathbb{R})$, is a bimeasure and $f : \mathbb{R} \rightarrow \mathbb{C}$ is μ -integral if and only if $(f, \overline{f})$ is β -integrable (in the sense of Definition 3). Furthermore, $EX_t \overline{X_s} = \int_{\mathbb{R}} \int_{\mathbb{R}} e^{itx} e^{-isy} d\beta(x, y)$, $t, s \in \mathbb{R}$.

In view of Theorem 5 as well as the above statements, Definition 3 appears to provide (at least in a stochastic framework) the appropriate conditions for bimeasure integration. In fact, such an analysis can be extended to matrix bimeasure, as shown in Houdré [4]. The reader is also referred to Kluváněk [5] for illuminating remarks on bimeasures.

Acknowledgements Professors P. E. Caines and J. C. Taylor of McGill University are gratefully acknowledged for their supervision and support throughout the author's doctoral studies, upon which this work is based. The author is also grateful to Professor E. Thomas for his correspondence on bimeasures.

References

- [1] G. Bartle, N. Dunford and J. T. Schwartz, "Weak compactness and vector measures". *Canad. J. Math.* 7, (1955), pp. 289-305.
- [2] D. K. Chang and M. M. Rao, "Bimeasures and sampling theorems for weakly harmonizable processes". *Stochastic Anal. Appl.* 1, (1983), pp. 21-55.
- [3] N. Dunford and J. T. Schwartz, *Linear Operators, Part I: General Theory*, Interscience, New York, 1958.
- [4] C. Houdré, *Non-Stationary Processes, System Theory and Prediction*. Ph.D. Thesis, McGill University, Montréal, Québec, 1987.
- [5] I. Kluvánek, "Remarks on bimeasures". *Proc. A.M.S.*, 81, (1981), pp. 233-239.
- [6] M. Morse and W. Transue, "Integral representations of bilinear functionals". *Proc. Nat. Acad. Sci. U.S.A.*, 35, (1949), pp. 136-143.
- [7] M. Morse and W. Transue, "C-bimeasures Λ and their superior integrals Λ^* ". *Rend. Circ. Matem. Palermo*, (2) 4, (1955), pp. 270-300.
- [8] M. Morse and W. Transue, "C-bimeasures Λ and their integral extensions". *Ann. Math.* 64, (1956), pp. 480-504.
- [9] H. Niemi, "Stochastic processes as Fourier transforms of stochastic measures". *Ann. Acad. Sci. Fenn.*, Ser. AI, 591, (1975), pp. 1-47.
- [10] E. Thomas, "L'intégration par rapport à une mesure de Radon vectorielle". *Ann. Inst. Fourier (Grenoble)* 20:2, (1970), pp. 55-191.
- [11] K. Ylisen, "On vector bimeasures". *Ann. Acad. Pura Appl.* (4) 117, (1978), pp. 115-138.

END

DATE

FILMED

DTIC

July 88