

AD-A193 584

SIMULATING DISCOUNTED COSTS(U) STANFORD UNIV CA DEPT OF 1/1
OPERATIONS RESEARCH B L FOX ET AL. AUG 87 TR-23
DAGC29-84-K-0030

UNCLASSIFIED

F/G 5/3

ML

END

DATE

784



AD-A193 584

SIMULATING DISCOUNTED COSTS

by

Bennett L. Fox

and

Peter W. Glynn

TECHNICAL REPORT NO. 23

August 1987

Prepared under the Auspices

of

U.S. Army Research Contract*
DAAG29-84-K-0030

Approved for public release: distribution unlimited.

Reproduction in whole or in part is permitted for any
purpose of the United States government.

DEPARTMENT OF OPERATIONS RESEARCH
STANFORD UNIVERSITY
STANFORD, CALIFORNIA

DTIC
ELECTE

APR 12 1988

CE
E

*This research was also partially supported under
National Science Foundation Grants DCR 85-09668
and ECS-8404809 and the Natural Sciences and
Engineering Research Council of Canada.

SIMULATING DISCOUNTED COSTS

by

Bennett L. Fox¹

and

Peter W. Glynn²

ABSTRACT

We numerically estimate, via Monte Carlo simulation, the expected infinite-horizon discounted cost d of running a stochastic system; we permit the discount rate to be state dependent. By exploiting various types of stochastic structure, we beat the naive strategy of estimating a finite-horizon approximation to d . Efficient estimators are obtained for systems which are semi-Markov and/or regenerative. These estimators are then ranked with respect to asymptotic variance as a function of computer-time budget and discount rate.

KEYWORDS: Discounted costs, simulation, semi-Markov process, regenerative process.

Accession For	
DTIC	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Avail and/or	
Dist	Special

A-1

DTIC
COPY
SELECTED

¹ Department of Computer Science and Operations Research, University of Montreal. Supported by the Natural Sciences and Engineering Research Council of Canada.

² Department of Industrial Engineering, University of Wisconsin-Madison. Supported by the National Science Foundation under Grant ECS-8404809 and the U.S. Army under Contract No. DAAG29-84-K-0030.

INTRODUCTION

In many settings, discounted costs arise naturally. This paper describes simulation methodologies for estimation of expected discounted costs associated with systems that exhibit stochastic fluctuations. Such techniques are important for numerical computation of discounted costs for stochastic processes in which conventional numerical methods either fail to apply or are inefficient. Examples of such processes include non-Markov processes or infinite state space Markov chains. The discussion given here of simulation algorithms for the discounted cost problem also merits interest to the extent that it provides an excellent vehicle for illustrating several sophisticated "variance reduction" methods for stochastic simulation. These techniques are more accurately called efficiency increase techniques, as we shall see.

To formulate the estimation problem mathematically, we let $X = \{X(t) : t \geq 0\}$ be a stochastic process taking values in a state space S . Suppose that f, g are two real-valued functions defined on S , in which $f(x)$ represents the cost of running X in state x and $g(x)$ corresponds to the (positive) discount rate in state x . Then

$$D = \int_0^\infty \exp(-V(s)) f(X(s)) ds \quad (1)$$

is the infinite-horizon discounted cost, where $V(s) = \int_0^s g(X(u)) du$. Our goal in this paper is to construct Monte Carlo simulation algorithms for numerically evaluating $d = ED$.

We now describe the layout of the rest of this paper. Section 2 develops a naive estimator for d based on truncation of the infinite-horizon integral, and studies its relevant theory. In Section 3, an estimator based on randomizing the truncation point is developed, and it is shown that for large computational budgets, this estimator beats the naive truncation estimator of Section 2. Section 4 shows how to exploit semi-Markov process structure to improve the efficiency of the randomized estimator of Section 3 by "conditioning out" the holding times. In Section 5, an estimator which makes use of regenerative structure is explored, whereas Section 6 studies an estimator which utilizes both semi-Markov and regenerative structure to obtain efficiency. Finally, in Section 7, we compare the asymptotic variances of the above five estimators when the discount rate is small; a small discount rate is natural in many economic settings. General conclusions on the choice of estimator can also be found there. Unless otherwise indicated, all proofs are deferred to the Appendix to make the paper easier to read.

The table below summarizes features of the five estimators that we consider.

	δ_1	δ_2	δ_3	δ_4	δ_5
truncates	Yes	No	No	No	No
"conditions out" holding times	No	No	Yes	No	Yes
uses regenerative structure	No	No	No	Yes	Yes
becomes less efficient as discount rate decreases	No	Yes	Yes	No	No
variance per run	see below	δ_3 beats δ_2		δ_5 beats δ_4	
overall efficiency	always loses for large enough computer budgets	δ_2, δ_3 always lose to δ_4, δ_5 for small enough discount rate		depends on expected work per run as well as variance per run; see text	

Efficiency of a simulation algorithm depends on both the time required to generate observations and their variance. In the setting of continuous-time Markov chains, δ_3 and δ_5 are more cheaply simulated than δ_2 and δ_4 ; since they also reduce variance, they are clear winners for this class of processes. For small discount rates, δ_5 is always more efficient than δ_3 ; for moderate to large rates, no universal conclusion is possible. If the process simulated is non-regenerative, the discount rate is small, and the computer-time budget is modest, then δ_1 probably beats the other estimates. Our mathematical analysis in the following sections supports these assertions.

2. A TRUNCATION ALGORITHM

Naive Monte Carlo simulation, based directly on (1), is impractical since generating the r.v. D generally requires an infinite amount of computation time. A straightforward alternative truncates the r.v. D at some finite time horizon β , yielding the quantity

$$D(\beta) = \int_0^\beta \exp(-V(s)) f(X(s)) ds.$$

Given a computational budget t , it is clear that the truncation point β should increase with t . (Observe that a sample mean estimator based on $D(\beta)$ with β fixed converges to $ED(\beta)$, which is in general not equal to d .)

As a consequence, we need to define a sampling plan $\{\beta(t) : t > 0\}$, in which $\beta(t)$ corresponds to the truncation point associated with computer budget t . Assuming that the time required to generate a replicate of $D(\beta)$ is $c_1\beta$ ($c_1 > 0$), we find that the number $n(t)$ of runs completed with budget t is $\lfloor t/c_1\beta(t) \rfloor$. Given t , our estimator for d will then be

$$\delta_1(t) = \begin{cases} \frac{1}{n(t)}(D_1(\beta(t)) + D_2(\beta(t)) + \cdots + D_{n(t)}(\beta(t))); & n(t) \geq 1 \\ 0; & n(t) = 0 \end{cases} \quad (2)$$

where $\{D_n(\cdot) : n \geq 1\}$ is a sequence of i.i.d. replicates of $D(\cdot)$.

We now investigate the choice of sampling plan which optimizes the behavior of the estimator $\delta_1(t)$. Given the exponential character of discounting, it seems reasonable to expect that the bias of $\delta_1(t)$ behaves like $a \exp(-c\beta(t))$ for some constants a, c . This expectation can be justified when X is a regenerative stochastic process; see Glynn and Whitt (1987). In any case, it then follows that the mean square error (MSE) of $\delta_1(t)$ is given approximately by

$$E(\delta_1(t) - d)^2 \approx c_1(\text{var } D) \frac{\beta(t)}{t} + a^2 \exp(-2c\beta(t))$$

The choice of $\beta(t)$ which minimizes the above MSE expression is

$$\beta^*(t) = \lambda_0^* + \lambda_1^* \log t$$

where $\lambda_0^* = -\log(c_1 \text{var } D / 2a^2c) / 2c$, $\lambda_1^* = 1/2c$.

Theorem 1 below shows that the above approximations can be justified rigorously; its proof relies heavily on the general theory of replication estimators of the form (2), as described in Fox and Glynn (1947). The following (reasonable) assumptions will be needed:

H1. f, g are strictly positive functions on S .

H2. $0 < \text{var } D < \infty$

H3. $b(\beta) \equiv d - ED(\beta) \sim ae^{-c\beta}$ as $\beta \rightarrow \infty$ for some constants a, c .

We require that f be strictly positive merely to simplify the technical statements of the theorems presented in this paper. It is not necessary and can be replaced by suitable (cumbersome) absolute integrability hypotheses on f .

Theorem 1. Assume H1-H3 and suppose β^* is defined by (3). Then:

i) If $\beta(t) = \beta^*(t)$, then

$$\sqrt{\frac{t}{\log t}}(\delta_1(t) - d) \Rightarrow \left(\frac{c_1 \text{var } D}{2c}\right)^{1/2} N(0, 1).$$

ii) If $\beta(t)/\beta^*(t) \rightarrow \kappa > 1$ as $t \rightarrow \infty$, then

$$\sqrt{\frac{t}{\log t}}(\delta_1(t) - d) \Rightarrow \left(\frac{\kappa c_1 \text{var } D}{2c}\right)^{1/2} N(0, 1).$$

iii) If $\beta(t)/\beta^*(t) \rightarrow \kappa < 1$ as $t \rightarrow \infty$, then

$$\sqrt{\frac{t}{\log t}}|\delta_1(t) - d| \Rightarrow \infty.$$

Note that part iii) forces the selection $\beta(t) \geq \beta^*(t)$ for large enough t . On the other hand, if the constant κ appearing in ii) is strictly greater than one, the variance of the limiting normal r.v. is greater than that obtained when $\beta = \beta^*$. We conclude that Theorem 1 shows that the asymptotically optimal choice of sampling plan is $\beta = \beta^*$.

Implementing this choice requires determining a and c . (A glance at the proof of Theorem 1 shows that in fact c is the crucial parameter, in the sense that if $\beta(t) = \log t/2c + \xi$, then convergence result i) always ensues, regardless of the choice of ξ). Theorem 1 indicates that if one is to guess a choice for c , it is better to underestimate c than overestimate it. In particular, suppose that one uses $\beta(t) = \log t/2c' + \xi$ with $c' < c$. Then, $\delta_1(t)$ will satisfy relation ii) with $\kappa = c/c'$; on the other hand, if $c' > c$, $\delta_1(t)$ has the poor convergence structure associated with iii). An underestimate c' of c is always available when $\Lambda = \inf\{g(x) : x \in S\} > 0$, namely $c' = \Lambda$.

Theorem 1 also shows that even if β is chosen optimally, the best possible rate of convergence is $O(\sqrt{\log t/t})$. This is unsatisfactory in comparison to the canonical rate of $O(1/\sqrt{t})$ typical of Monte Carlo simulation. Thus, the straightforward truncation approach of this section appears inefficient for large computational budgets t , and the investigation of alternative algorithms is warranted. Heuristic adjustments to $\beta^*(t)$ may be appropriate when the computer-time budget is only moderate.

3. A RANDOMIZED ESTIMATOR

A principal difficulty in estimating d is that the naive Monte Carlo estimator based on replicates of D is inadmissible: it requires infinite time to simulate a single observation of D . Hence, it is clear that one must carefully consider the computational effort required per observation in order to properly assess the efficiency of an estimator. Hammersley and Handscomb (1964) proposed evaluating the efficiency of a Monte Carlo procedure via the formula

$$\text{Efficiency} = (\text{Time})^{-1} \cdot (\text{Variance})^{-1} \quad (3)$$

where Time = expected computation time per observation

Variance = variance per observation.

Glynn and Whitt (1986) rigorously justify this criterion. Thus, the efficiency of an estimator may be improved by reducing computation time per observation and/or reducing variance. An important implication of this observation is that the efficiency of an estimator may be improved by increasing the variance per observation provided that the computational time required per observation is appropriately decreased. The estimator proposed in this section has precisely this property. Specifically, the variance per observation is greater than that of D , but the observations can be generated in finite time.

Suppose R is an exponential r.v. with mean one, which is independent of X . Set

$$\hat{D}(1) = \int_0^{V^{-1}(R)} f(X(t)) dt$$

where $V^{-1}(\cdot) = \inf\{t > 0 : V(t) \geq \cdot\}$; we call $\hat{D}(1)$ a randomized estimator since it involves adding additional randomness to the probability space. Note that $\hat{D}(1)$ requires simulating X only up to time $V^{-1}(R)$, and can therefore be generated in finite time if:

$$\text{H4. } V(\infty) \equiv \int_0^\infty g(X(t)) dt = \infty \text{ a.s.}$$

When $g \equiv \alpha$, then $EV^{-1}(R) = 1/\alpha$ and $\text{var } V^{-1}(R) = 1/\alpha^2$. Thus, both the expected work per run and the variance per run are (generally significantly) affected by α .

The following proposition shows that efficient estimation of d can be based on $\hat{D}(1)$, since $E\hat{D}(1) = d$.

Proposition 1. Assume H1, H2, H4. Then,

$$D = E\{\hat{D}(1)|X\}, \text{ so that } E\hat{D}(1) = d.$$

Standard properties of conditional expectation guarantee that $\text{var } D \leq \text{var } \tilde{D}(1)$, so that the variance per observation is increased by using an estimator based on $\tilde{D}(1)$. To analyze the efficiency of $\tilde{D}(1)$, we will obtain a central limit theorem (CLT) for the corresponding estimator. Let $\{(\tilde{D}_n(1), V_n^{-1}(R_n)) : n \geq 1\}$ be a sequence of i.i.d. copies of $(\tilde{D}(1), V^{-1}(R))$. Given t units of computation time, the number of observations generated is

$$N_1(t) = \max\{n \geq 0 : c_1(V_1^{-1}(R_1) + \dots + V_n^{-1}(R_n)) \leq t\}$$

(disregarding overhead for generating the R_i 's) and the estimator $\delta_2(t)$ available after t units of computational effort have been expended is

$$\delta_2(t) = \begin{cases} \frac{1}{N_1(t)}(\tilde{D}_1(1) + \dots + \tilde{D}_{N_1(t)}(1)); & N_1(t) \geq 1 \\ 0; & N_1(t) = 0. \end{cases} \quad (4)$$

Theorem 2 shows that $\delta_1(t)$ converges at rate $O(t^{-1/2})$; it can also be used, in a straightforward way, to obtain confidence intervals for d .

Theorem 2. Assume H1, H4, and $E\tilde{D}(1)^2 < \infty$. Then,

$$t^{1/2}(\delta_2(t) - d) \Rightarrow (c_1 \text{var } \tilde{D}(1) \cdot EV^{-1}(r))^{1/2} N(0, 1)$$

Furthermore, $\text{var } \tilde{D}(1) = 2 \int_0^\infty \int_0^t E\{\exp(-V(t))f(X(s))f(X(t))\}dsdt - d^2$ and $EV^{-1}(R) = \int_0^\infty E \exp(-V(t))dt$.

This theorem confirms the efficiency criterion specified by (3), in the sense that the asymptotic variance of the limiting normal r.v. is precisely the reciprocal of the efficiency given by (3). In the next three sections, we will describe estimation algorithms that will increase the efficiency of $\delta_2(t)$ by reducing the variance of $\tilde{D}(1)$ without increasing the average amount of time required to generate an observation; the improved efficiency will be obtained by utilizing special stochastic structure in X .

4. DISCRETE-TIME CONVERSION FOR SEMI-MARKOV PROCESSES

In this section, we construct an efficient estimator for d which exploits semi-Markov process (SMP) type structure; the idea is to eliminate some of the variance in $\tilde{D}(1)$ by conditioning on the embedded discrete-time process which describes the sequence of states visited by X . This "discrete-time conversion" is similar in spirit to the estimator discussed in Fox and Glynn (1986) for estimation of steady-state quantities associated with SMP's. It "undoes" some of the variance increase due to randomization.

Specifically, we assume in this section the existence of a discrete-time process $Y = \{Y_n : n \geq 0\}$ taking values in S and a strictly increasing sequence of random times $\{S_n : n \geq 0\}$ such that

H5. i) $X(t) = \sum_{n=0}^{\infty} Y_n I(S_n \leq t < S_{n+1})$ where $S_0 = 0$.

ii) $\{\beta_n : n \geq 0\}$ is conditionally independent given Y , where $\beta_n = S_{n+1} - S_n$.

iii) $P\{\beta_n \in dt | Y\} = F(Y_n, Y_{n+1}, dt)$ for some family of distributions F on $S \times S$.

H5 generalizes the notion of SMP, since we do not require here that Y be a Markov chain.

To apply discrete-time conversion, we let $N(t) = \max\{n \geq 0 : S_n \leq t\}$ be the number of transitions of X by time t , and set $M = N(V^{-1}(r))$. From H5, we get

$$\begin{aligned} \tilde{D}(1) &= \sum_{j=0}^{M-1} \int_{S_j}^{S_{j+1}} f(X(t)) dt + \int_{S_M}^{V^{-1}(R)} f(X(t)) dt \\ &= \sum_{j=0}^{M-1} f(Y_j) \beta_j + f(Y_M)(V^{-1}(R) - S_M) \end{aligned}$$

The "discrete-time" estimator of this section is based on $\tilde{D}(2) = E\{\tilde{D}(1) | Y, M\}$, that is

$$\begin{aligned} \tilde{D}(2) &= \sum_{j=0}^{M-1} f(Y_j) E\{\beta_j | Y, M\} \\ &\quad + f(Y_M) E\{V^{-1}(r) - S_M | Y, M\}. \end{aligned} \tag{5}$$

Let $\varphi(x, y, \lambda)$ be the Laplace transform of the distribution $F(x, y, dt)$ defined by

$$\varphi(x, y, \lambda) = \int_{[0, \infty)} e^{-\lambda t} F(x, y, dt).$$

It is easy to show, using a dominated convergence argument, that the derivative $\varphi'(x, y, \lambda)$ with respect to λ exists for all positive arguments λ . Let $\{(\varphi_k, \varphi'_k) : k \geq 0\}$ be the sequence defined by

$$\varphi_k = \varphi(Y_k, Y_{k+1}, g(Y_k))$$

$$\varphi'_k = \varphi'(Y_k, Y_{k+1}, g(Y_k)).$$

With this notation in hand, the next proposition calculates the conditional expectations appearing in (5), as well as the conditional distribution of M given Y .

Proposition 2. Assume H1, H4, and $E\tilde{D}(1) < \infty$. Then, for $k < m$

$$\begin{aligned} P\{M = m | Y\} &= \prod_{j=0}^{m-1} \varphi_j (1 - \varphi_m) \\ E\{\beta_k | Y, M = m\} &= -\varphi'_k / \varphi_k \\ E\{V^{-1}(R) - S_M | Y, M\} &= \frac{(1 + g(Y_M)\varphi'_M - \varphi_M)}{g(Y_M)(1 - \varphi_M)}. \end{aligned}$$

As a consequence of Proposition 2, we find that

$$\tilde{D}(2) = - \sum_{j=0}^{M-1} f(Y_j) \varphi'_j / \varphi_j + f(Y_M) \frac{(1 + g(Y_M) \varphi'_M - \varphi_M)}{g(Y_M)(1 - \varphi_M)} \quad (6)$$

Formula (6) shows that we get $\tilde{D}(2)$ by generating Y up to time M , where M is generated by using the conditional distribution given in Proposition 2. The following algorithm can be used to produce r.v.'s with the distribution of $\tilde{D}(2)$; its validity follows immediately from (6), noting that M is generated by "inversion".

Algorithm A:

1. Generate a random variate U , uniform on $(0, 1)$.

2. Generate Y_0 .

3. Set $m \leftarrow 0, \Lambda \leftarrow 1, \Gamma \leftarrow 0$.

Comment: now $\Lambda = P\{M \geq 0|Y\}$.

4. Generate Y_{m+1} .

5. Set $\Lambda \leftarrow \Lambda \varphi_m$.

Comment: now $\Lambda = P\{M \geq m+1|Y\}$

6. If $U > \Lambda$, then

i) Set $D \leftarrow f(Y_m) \frac{(1 + g(Y_m) \varphi'_m - \varphi_m)}{g(Y_m)(1 - \varphi_m)} - \Gamma$.

Comment: now $M = m$

ii) exit

7. Else,

i) set $\Gamma \leftarrow \Gamma + f(Y_m) \varphi'_m / \varphi_m$

ii) set $m \leftarrow m + 1$

iii) go to step 4.

An estimator $\delta_3(t)$ based on a sequence $\{(\tilde{D}_n(2), M_n) : n \geq 1\}$ of i.i.d. replicates of $\tilde{D}(2)$ can be constructed analogously to $\delta_2(t)$ (see (4)). The estimator $\delta_3(t)$ so defined is a sample mean of $N_2(t)$ observations of $\tilde{D}(2)$, where $N_2(t)$ is the number of observations generated in t units of computer time. To a first approximation, $N_2(t) = \max\{n \geq 0 : c_2(M_1 + \dots + M_n) \leq t\}$ where c_2 is the computer time required to increment m by one in Algorithm A. (This disregards the set-up time to generate M , and the fact that the effort required to execute steps 4 to 7 of Algorithm A depends on the random states occupied at times m and $m+1$.)

The following CLT describes the behavior of the estimator $\delta_3(t)$, and can be used to construct confidence intervals for d .

Theorem 3. Assume H1, H4, and $E\tilde{D}(2)^2 < \infty$. Then,

$$t^{1/2}(\delta_3(t) - d) \Rightarrow (c_2 EM \cdot \text{var } \tilde{D}(2))^{1/2} \cdot N(0, 1)$$

as $t \rightarrow \infty$.

The proof of this result follows immediately from Section 5 of Glynn and Whitt (1986). Since $\tilde{D}(2) = E\{\tilde{D}(1)|Y, M\}$, it follows by the principle of conditional Monte Carlo (see Bratley, Fox, and Schrage (1983)) that $\text{var } \tilde{D}(2) \leq \text{var } \tilde{D}(1)$. Thus, the estimator $\delta_3(t)$ is obtained from $\delta_2(t)$ by reducing the variance per observation. However, as Theorems 2 and 3 point out, an efficiency increase is obtained only if $(c_2 EM)/(c_1 EV^{-1}(R)) \leq \text{var } \tilde{D}(1)/\text{var } \tilde{D}(2)$.

To fully understand this condition, note that $\text{var } \tilde{D}(1)/\text{var } \tilde{D}(2)$ reflects the degree to which randomness in $\tilde{D}(1)$ is due to the holding times β_j , as opposed to the embedded sequence Y . On the other hand, the ratio $c_2 EM/c_1 EV^{-1}(R)$ describes the complexity of generating a $\tilde{D}(2)$ observation relative to a $\tilde{D}(1)$ variate. Observe that both types of observations require generating Y up to time M ; the difference is that $\tilde{D}(1)$ additionally requires generating the holding times β_j , while $\tilde{D}(2)$ involves the Laplace transform quantities φ_j and φ'_j . If the $F(x, y, dt)$'s are distributions having Laplace transforms that are easily numerically evaluated (as is the case with gamma r.v.'s, for example), then the (possible) increase in effort involved in passing from $\tilde{D}(1)$ to $\tilde{D}(2)$ should be modest; in these circumstances, $\delta_3(t)$ is more efficient than $\delta_2(t)$. For a more detailed comparison of "discrete-time" estimators with their "continuous-time" analogs, see Section 2 of Fox and Glynn (1986).

5. ESTIMATION FOR REGENERATIVE PROCESSES

We assume now that X is a (possibly) delayed regenerative process with regeneration times $0 \leq T_0 < T_1 < \dots$ (If X is non-delayed, set $T_0 = 0$.); thus, we do not require in this section that X satisfy the semi-Markov hypothesis H5. The independence of regenerative cycles implies that

$$d = EA(0) + EC(0)EK(0) \tag{7}$$

where

$$\begin{aligned} A(i) &= \int_0^{T_i} \exp\left(-\int_0^t g(X(T_{i-1} + s))ds\right) f(X(T_{i-1} + t))dt \\ C(i) &= \exp\left(-\int_0^{T_i} g(X(T_{i-1} + t))dt\right) \dots \dots \dots \\ K(i) &= \int_0^\infty \exp\left(-\int_0^t g(X(T_{i-1} + s))ds\right) f(X(T_i + t))dt \end{aligned}$$

and $\tau_i = T_i - T_{i-1}$. A similar analysis of $EK(0)$ shows that

$$EK(0) = EA(1) + EC(1)EK(1).$$

But $K(1)$ has the same distribution as $K(0)$ by the regenerative property, so $EK(0) = EK(1)$. We conclude that $EK(0) = EA(1) \cdot (1 - EC(1))^{-1}$. Substituting into (7) yields

$$d = EA(0) + EC(0)EA(1) \cdot (1 - EC(1))^{-1}. \quad (8)$$

Equation (8) suggests that d can be estimated by simulating regenerative cycles. Since each regenerative cycle can be generated in finite time, independently of g , we will avoid the problems inherent in trying to generate D explicitly, or, when the discount rate is small, in randomizing as in Sections 3 and 4. (See also Section 7.) In the discounting context, it is important to allow the possibility that X is a delayed regenerative process (as opposed to steady-state simulation). For example, if one is asked to compute the discounted cost for a Markov chain initiated with a distribution concentrated on more than one point, this generalization would be required.

Since (8) involves two different types of cycles (delayed and non-delayed), it offers the possibility to stratify the computation effort so as to maximize the efficiency of the resulting estimators. Given a computational budget t , we allocate a proportion p to generating pairs $(C(0), A(0))$ and a proportion $q = 1 - p$ to simulating the pairs $(C(1), A(1))$ from the non-delayed cycle. An estimator $\delta_4(t)$ is then obtained by substituting the resulting sample means in (8).

To be precise, let $\{(C_n(i), A_n(i), \tau_{ni}) : n \geq 1 (i = 0, 1)\}$ be two independent sequences of i.i.d. random vectors where $(C_n(i), A_n(i))$ shares the same distribution as $(C(i), A(i))$, and where τ_{ni} represents the length of the corresponding cycle used to obtain $(C_n(i), A_n(i))$. Thus, if we set $p_0 = p, p_1 = q$, then $N^i(t) = \max\{n \geq 0 : c_1(\tau_{1i} + \dots + \tau_{ni}) \leq p_i t\}$ is the number of type i cycles completed by time t . Put

$$(\bar{C}_t(i), \bar{A}_t(i)) = \begin{cases} \frac{1}{N^i(t)} ((C_1(i), A_1(i)) + \dots + (C_{N^i(t)}(i), A_{N^i(t)}(i))) & N^i(t) \geq 1 \\ 0 & N^i(t) = 0. \end{cases}$$

Then, the estimator $\delta_4(t)$ is given by

$$\delta_4(t) = \bar{A}_t(0) + \bar{C}_t(0)\bar{A}_t(1) \cdot (1 - \bar{C}_t(1))^{-1}$$

To analyze the behavior of this estimator, we derive a CLT for $\delta_4(t)$. (Again, this can also be used to produce confidence intervals for d .) We require that:

H6. $E\tau_i < \infty$ ($i = 0, 1$).

Theorem 4. Assume H1, H2, and H6. Then, for $0 < p < 1$,

$$t^{1/2}(\delta_4(t) - d) \Rightarrow (\sigma_0^2/p + \sigma_1^2/q)^{1/2} N(0, 1)$$

as $t \rightarrow \infty$, where

$$\begin{aligned}\sigma_0^2 &= c_1 \text{var}(A(0) + C(0) \cdot EK(1)) \cdot E\tau_0 \\ \sigma_1^2 &= c_1 \left(\frac{EC(0)}{1 - EC(1)} \right)^2 \text{var}(A(1) + C(1) \cdot EK(1)) \cdot E\tau_1.\end{aligned}$$

To optimize the performance of $\delta_4(t)$, we select p to minimize the asymptotic variance term $\sigma_0^2/p + \sigma_1^2/q$. It is easily verified that the minimizer is given by

$$p^* = \sigma_0(\sigma_0 + \sigma_1)^{-1}$$

(provided $\sigma_0 + \sigma_1 > 0$) where $\sigma_i = \sqrt{\sigma_i^2}$, in which case the corresponding variance is $(\sigma_0 + \sigma_1)^2$. To compare the efficiency of the estimator with the previous ones, in particular the truncation estimator, it is useful to relate the coefficients defining σ_0^2 and σ_1^2 to $\text{var } D$ appearing in Theorem 1.

Proposition 3. Assume H1 and H2. Then

$$\text{var } D = \text{var}(A(0) + C(0) \cdot EK(1)) + \frac{EC(0)^2}{1 - EC(1)^2} \text{var}(A(1) + C(1) \cdot EK(1))$$

To aid in comparison, note that $EC(0)^2 \geq (EC(0))^2$ and $E(1 - C(1)^2) \geq (1 - EC(1))^2$. (For the second inequality, $0 \leq C(1) \leq 1$ so $EC(1) \geq EC(1)^2$. Hence $1 - EC(1) \leq 1 - EC(1)^2$. But since $0 \leq EC(1) \leq 1$, $(1 - EC(1))^2 \leq 1 - EC(1)$.) In the non-delayed case where $C(0) = 1$, $A(0) = 0$, we choose $p = 1$ (obviously). Theorems 1 and 4 then suggest that

$$\text{var } \delta_1(t) / \text{var } \delta_4(t) \sim \frac{\log t \cdot (1 - EC(1))^2}{2bE\tau_1 \cdot (1 - EC(1)^2)}. \quad (9)$$

We conclude that if $t \gg \exp(2bE\tau_1)$, it is better to use $\delta_4(t)$.

6. ESTIMATION FOR REGENERATIVE SEMI-MARKOV PROCESSES

In this section, we illustrate how the methods of Section 4 and 5 can be combined to obtain an estimator $\delta_5(t)$ which exhibits the best features of $\delta_3(t)$ and $\delta_4(t)$. In particular, $\delta_5(t)$ exploits the regenerative structure of X while "filtering out" the variance in $\delta_4(t)$ due to holding time randomness; the latter property is achieved by using discrete-time conversion.

Returning to the set-up of Section 3, we now assume that the embedded sequence Y is regenerative. Thus, we require that Y possess regeneration times $0 \leq U_0 < U_1 < \dots$ and set

$U_{-1} = 0$, $\eta_i = U_i - U_{i-1}$. By the conditional independence of the β_j 's given Y , it follows that the random times $T_i = S_{U_i}$ are regeneration times for X . Hence, (8) is valid for the T_i 's; as an immediate consequence, we obtain the identity

$$d = E\tilde{A}(0) + E\tilde{C}(0)E\tilde{A}(1) \cdot (1 - E\tilde{C}(1))^{-1} \quad (10)$$

where $\tilde{A}(i) = E\{A(i)|Y\}$, $\tilde{C}(i) = E\{C(i)|Y\}$. To compute the conditional expectations appearing in (10), observe that

$$\begin{aligned} E\{C(i)|Y\} &= E\left\{\prod_{j=U_{i-1}}^{U_i-1} \exp(-g(Y_j)\beta_j)|Y\right\} \\ &= \prod_{j=U_{i-1}}^{U_i-1} \varphi_j \end{aligned}$$

and

$$\begin{aligned} E\{A(i)|Y\} &= E\left\{\prod_{j=U_{i-1}}^{U_i-1} \int_{S_j}^{S_{j+1}} \exp\left(-\int_0^t g(X(T_{i-1}+s))ds\right) f(Y_j)|Y\right\} \\ &= E\left\{\prod_{j=U_{i-1}}^{U_i-1} \prod_{k=0}^{j-1} \exp(-g(Y_k)\beta_k) \int_0^{\beta_j} \exp(-g(Y_j)t)dt f(Y_j)|Y\right\} \\ &= \prod_{j=U_{i-1}}^{U_i-1} \prod_{k=0}^{j-1} \varphi_k (1 - \varphi_j) \frac{f(Y_j)}{g(Y_j)}. \end{aligned}$$

Given the above formulas, it is straightforward to generate the pairs $(\tilde{C}(i), \tilde{A}(i))$ by simulating the sequence Y . As in Section 4, the computational effort may be assigned so that a fraction p_i of the total time t is delegated to generation of pairs $(\tilde{C}(i), \tilde{A}(i))$, $(i = 0, 1)$. An estimator $\delta_5(t)$ can then be constructed analogously to $\delta_4(t)$.

We can derive a CLT for $\delta_5(t)$ which describes its convergence and can be used for confidence interval estimation; the proof is analogous to that of Theorem 4 and is therefore omitted. The result (Theorem 5 below) assumes that the computational effort required to generate $(\tilde{C}(i), \tilde{A}(i))$ is $c_3\eta_i$. The constant c_3 reflects the difficulty of simulating the chain Y and numerically evaluating the φ_j 's (we do not assume that $c_2 = c_3$ since the discrete-time algorithm of Section 4 also involves numerical evaluation of the derivatives of the φ_j 's, which may be harder. For continuous-time Markov chains, however, both φ_j and φ_j' have simple closed forms.)

H7. $E\eta_i < \infty$ ($i = 0, 1$).

Theorem 5. Assume H1, H2, and H7. Then, for $0 < p < 1$,

$$t^{1/2}(\delta_5(t) - d) \Rightarrow (\tilde{\sigma}_0^2/p + \tilde{\sigma}_1^2/q)^{1/2} N(0, 1)$$

as $t \rightarrow \infty$, where

$$\begin{aligned}\tilde{\sigma}_0^2 &= c_3 \cdot \text{var}(E\{A(0) + C(0) \cdot EK(1)|Y\}) \cdot E\eta_0 \\ \tilde{\sigma}_1^2 &= c_3 \left(\frac{EC(0)}{1 - EC(1)} \right)^2 \text{var}(E\{A(1) + C(1) \cdot EK(1)|Y\}) \cdot E\eta_1.\end{aligned}$$

The principle of conditional Monte Carlo again guarantees that $\tilde{\sigma}_0^2 \leq \sigma_0^2$, $\tilde{\sigma}_1^2 \leq \sigma_1^2$; the amount of variance reduction depends on the extent to which the randomness of D is due to the holding times. As an immediate consequence, we find that if $c_1 E\tau_1 \approx c_3 E\eta_1$, $\delta_5(t)$ is more efficient than $\delta_4(t)$. The following proposition relates $\tilde{\sigma}_0^2$ and $\tilde{\sigma}_1^2$ to $\text{var}(E\{D|Y\})$; its proof is similar to Proposition 3 and is omitted.

Proposition 4. Assume H1 and H2. Then,

$$\text{var}(E\{D|Y\}) = \text{var}(\tilde{A}(0) + \tilde{C}(0) \cdot EK(1)) + \frac{E\tilde{C}(0)^2}{1 - E\tilde{C}(1)^2} \text{var}(\tilde{A}(1) + \tilde{C}(1) \cdot EK(1)).$$

This result can be used to compare the efficiency of $\delta_1(t)$ to $\delta_5(t)$ when Y is non-delayed. By arguing as in (9), we find that

$$\frac{\text{var } \delta_1(t)}{\text{var } \delta_5(t)} \sim \frac{\log t \cdot c_1 \text{var } D \cdot (1 - EC(1))^2}{2bE\eta_1 \cdot c_3 \cdot \text{var}(E\{D|Y\}) \cdot (1 - E\tilde{C}(1)^2)}.$$

Here, we find that if $t \gg \exp(2bc_3 E\eta_1 \text{var}(E\{D|Y\})/(\text{var } D \cdot c_1))$, $\delta_5(t)$ is more efficient than $\delta_1(t)$.

7. ANALYSIS OF EFFICIENCY FOR SMALL DISCOUNT RATES

In this section, we study the relative efficiencies for small discount rates of the methods considered above. Smallish discount rates arise naturally in many economic contexts (e.g. low inflation rate settings) and a considerable literature has developed on this topic. (See, for example, Veinott (1969) or Whitt (1972).)

To make our analysis precise, let

$$V_\alpha(t) = \alpha \int_0^t g(X(s)) ds$$

where g does not depend on α and set $d(\alpha) = ED_\alpha$, where

$$D_\alpha = \int_0^\infty \exp(-V_\alpha(t)) f(X(t)) dt.$$

We are interested in the efficiency of our five estimators for $D(\alpha)$ when α is small. Given Theorems 1 through 5, we examine the asymptotic behavior of the scaling constants appearing in front of the limiting normal r.v. These scaling constants determine the width

of the confidence interval associated with a given method, and consequently one wishes to choose estimators for which the scaling constants are as small as possible.

Our subsequent mathematical analysis requires:

H8. X is a (possibly) delayed regenerative process with regeneration times

$$0 \leq T_0 < T_1 < \dots$$

H9. $E(Y_i(f)^4 + Y_i(g)^4) < \infty$ ($i = 0, 1$), where

$$Y_i(f) = \int_{T_{i-1}}^{T_i} f(X(s)) dx$$

$$Y_i(g) = \int_{T_{i-1}}^{T_i} g(X(s)) dx$$

and $T_{-1} = 0$.

Although the results stated here require the regenerative structure for the proofs, it seems likely that the same asymptotic behavior holds for more general classes of processes. This belief is supported by some of the more general limit theorems appearing in Glynn and Whitt (1987).

To state the following theorem, we add an α -dependence to all the r.v.'s and constants appearing in Theorems 1 to 5. For example, $\tilde{D}_\alpha(1)$ is defined as

$$\int_0^{V_\alpha^{-1}(R)} f(X(t)) dt.$$

Theorem 6. Assume H1-H8. Then:

- (a) $d(\alpha) \sim \frac{1}{\alpha} \frac{r(f)}{r(g)}$
- (b) $\text{var } D_\alpha \sim \frac{1}{2\alpha} \frac{\sigma^2}{r(g)}$
- (c) $c(\alpha) \sim \alpha r(g)$
- (d) $\text{var } \tilde{D}_\alpha(1) \sim \frac{1}{\alpha^3} \frac{r^2(f)}{r^3(g)}$
- (e) $EV_\alpha^{-1}(R) \sim (\alpha r(g))^{-1}$
- (f) $\text{var } \tilde{D}_\alpha(2) \sim \frac{1}{\alpha^3} \frac{r^2(f)}{r^3(g)}$
- (g) $EM(\alpha) \sim (\alpha r(g))^{-1} \frac{E\eta_1}{E\tau_1}$
- (h) $(\sigma_0(\alpha) + \sigma_1(\alpha))^2 \sim \frac{\sigma_1^2}{\alpha^3} \frac{\sigma^2}{r(g)}$
- (i) $(\tilde{\sigma}_0(\alpha) + \tilde{\sigma}_1(\alpha))^2 \sim \frac{\sigma_1^2}{\alpha^3} \frac{\tilde{\sigma}^2}{r(g)} \frac{E\eta_1}{E\tau_1}$

as $\alpha \downarrow 0$, where $r(f) = EY_1(f)/E\tau_1$, $r(g) = EY_1(g)/E\tau_1$,

$$\sigma^2 = \frac{\text{var}(r(g)Y_1(f) - r(f)Y_1(g))}{E\tau_1} \cdot \frac{1}{r(g)^3}$$

$$\tilde{\sigma}^2 = \frac{\text{var}(E\{r(g)Y_1(f) - r(f)Y_1(g)|Y\})}{E\tau_1} \cdot \frac{1}{r(g)^3}.$$

Given a computation budget of (at least) moderate size t , the above theorem tells us that if the discount rate is small, then we can expect that

$$\begin{aligned}\text{var } \delta_1(t) &\approx \frac{\log t}{t} \cdot \frac{c_1}{\alpha^2} \cdot \frac{\sigma^2}{4r(g)} \\ \text{var } \delta_2(t) &\approx \frac{1}{t} \cdot \frac{c_1}{\alpha^3} \cdot \frac{r^2(f)}{r^3(g)} \\ \text{var } \delta_3(t) &\approx \frac{1}{t} \cdot \frac{c_2}{\alpha^3} \cdot \frac{r^2(f)}{r^3(g)} \cdot \frac{E\eta_1}{Er_1} \\ \text{var } \delta_4(t) &\approx \frac{1}{t} \cdot \frac{c_1}{\alpha^2} \cdot \frac{\sigma^2}{r(g)} \\ \text{var } \delta_5(t) &\approx \frac{1}{t} \cdot \frac{c_3}{\alpha^2} \cdot \frac{\tilde{\sigma}^2}{r(g)} \cdot \frac{E\eta_1}{Er_1}\end{aligned}$$

Assuming that $c_j E\eta_1 \leq c_1 Er_1$ for $j = 2, 3$ (i.e. the cost of simulating a regenerative cycle in discrete time is less than or equal to the cost of simulating a cycle in continuous time), the above analysis suggests that we can order (for small discount rates) the estimators in order of decreasing preference as follows: $\delta_5(t), \delta_4(t), \delta_1(t), \delta_3(t), \delta_2(t)$.

For larger discount rates, we would expect that the $\log t$ term in $\text{var } \delta_1(t)$ would dominate the additional factor of $1/\alpha$ appearing in $\text{var } \delta_2(t)$ and $\text{var } \delta_3(t)$. Thus, for larger discount rates, we recommend using the estimators in the order $\delta_5(t), \delta_4(t), \delta_3(t), \delta_2(t), \delta_1(t)$. This is a heuristic analysis, however, and for a given budget t , there may be some realignment in this order.

The above results also show that the discounting problem does not get harder as $\alpha \downarrow 0$, provided that we take advantage of regenerative structure. Suppose that we wish to construct a $100(1 - \delta)\%$ confidence interval for $d(\alpha)$ with half-width equal to $\epsilon\%$ of $d(\alpha)$. If the estimator $\delta_5(t)$ is used, the computational effort $t(\alpha)$ required for this relative width confidence interval is given approximately by

$$t(\alpha) \approx \frac{z^2(\delta) \tilde{\sigma}^2 c_3 \cdot E\eta_1 r(g)}{\epsilon^2 Er_1 r(f)^2}$$

where $z(\delta)$ solves $P\{N(0, 1) \leq z(\delta)\} = 1 - \delta/2$. Since the right-hand side does not depend on α , this shows that the discounting problem does not get harder, in a relative error sense, as the discount rate is driven to zero. This is in contrast to the problem of estimating steady-state queue-length in heavy-traffic, where the relative error problem does get harder as the traffic intensity increases to one (see Whitt (1987)).

APPENDIX

Proof of Theorem 1. First, observe that the positivity of f shows that the bias $b(\beta)$ is positive for all β . Furthermore, by H2, it is evident that $b(\beta)$ converges to zero as $\beta \rightarrow \infty$, and thus a and b must be positive finite constants.

We now apply the results of Fox and Glynn (1987) to obtain the theorem; it is easily checked that their hypotheses are in force. Their Proposition 1 states that $\beta(t) \rightarrow \infty$ as $t \rightarrow \infty$ is necessary for consistency of $\delta_1(t)$, while their Theorem 2 proves that

$$q(t)(\delta_1(t) - d) \Rightarrow N(0, 1) + \gamma \quad (A1)$$

as $t \rightarrow \infty$, where $q(t) = (t/c_1\beta(t) \text{ var } D(\beta(t)))^{1/2}$, and $\gamma = \lim_{t \rightarrow \infty} q(t)b(\beta(t))$. Since $\beta(t) \rightarrow \infty$, it is evident that

$$\text{var } D(\beta(t)) / \text{var } D \rightarrow 1 \quad (A2)$$

as $t \rightarrow \infty$. Furthermore,

$$t^{1/2}b(\beta(t)) = \frac{b(\beta(t))}{a \exp(-c\beta(t))} \cdot a \exp \left[-c(\beta(t) - \frac{1}{2c} \log t) \right]. \quad (A3)$$

Hence, if $\beta(t) = \beta^*(t)$, it follows that $t^{1/2}b(\beta(t))$ converges to a finite constant, so that $\gamma = 0$; part i) is then obtained by using (A1) and (A2). Similarly, for part ii), $t^{1/2}b(\beta(t)) \rightarrow 0$ so that $\gamma = 0$ and the result again follows immediately from (A1) and (A2). Finally for iii), (A3) shows that for t sufficiently large,

$$t^{1/2}b(\beta(t)) \geq \frac{a}{2} \exp \left[\frac{1}{4}(1 - \kappa) \log t \right] = \frac{a}{2} t^{(1-\kappa)/4}$$

so that $\gamma = \infty$, yielding the result.

Proof of Proposition 1. We can write $\tilde{D}(1)$ as

$$\begin{aligned} \tilde{D}(1) &= \int_0^\infty I(V^{-1}(R) > t) f(X(t)) dt \\ &= \int_0^\infty I(R > V(t)) f(X(t)) dt. \end{aligned}$$

The result then follows from Lemma 1 below, by noting that the independence of X and K proves that $E\{I(R > V(t))f(X(t))|X\} = f(X(t))P\{R > V(t)|X\} = f(X(t))\exp(-V(t))$.

Lemma 1. Let Z be a nonnegative process on a probability space $(\Omega, \mathcal{F}, P)$ such that $E \int_0^\infty Z(t) dt < \infty$. If $\mathcal{G}$ is a sub- σ -field of $\mathcal{F}$, then

$$E\left\{ \int_0^\infty Z(t) dt \middle| \mathcal{G} \right\} = \int_0^\infty E\{Z(t)|\mathcal{G}\} dt \quad \text{a.s.}$$

Proof of Lemma 1. We use the defining relation for conditional expectation, as given on p. 298 of Chung (1974). Note that $\int_0^\infty E\{Z(t)|\mathcal{G}\}dt$ is a $\mathcal{G}$ -measurable r.v. such that if $A \in \mathcal{G}$,

$$\begin{aligned} E\left(\int_0^\infty E\{Z(t)|\mathcal{G}\}dt \cdot I(A)\right) &= \int_0^\infty E(I(A)E\{Z(t)|\mathcal{G}\})dt \\ &= \int_0^\infty E(I(A)Z(t))dt \\ &= E\left(\int_0^\infty Z(t)dt \cdot I(A)\right); \end{aligned}$$

the first and third equalities use Fubini's theorem, whereas the second follows from the defining relation for $E\{Z(t)|\mathcal{G}\}$. We have therefore demonstrated that $\int_0^\infty E\{Z(t)|\mathcal{G}\}dt$ satisfies the defining relation for $E\{\int_0^\infty Z(t)dt|\mathcal{G}\}$, proving the result.

Proof of Theorem 2. The CLT for $\delta_2(t)$ follows immediately from Section 5 of Glynn and Whitt (1986). For the expression for $\text{var } \tilde{D}(1)$, note that

$$\begin{aligned} E\tilde{D}(1)^2 &= 2E\left\{\int_0^{V^{-1}(R)} \int_0^t f(X(s))f(X(t))dsdt\right\} \\ &= 2E\left\{\int_0^\infty \int_0^t I(R > V(t))f(X(s))f(X(t))dsdt\right\} \\ &= \int_0^\infty \int_0^t E\{I(R > V(t))f(X(s))f(X(t))\}dsdt. \end{aligned}$$

But $E\{I(R > V(t))f(X(s))f(X(t))|X\} = f(X(s))f(X(t)) \cdot P\{R > V(t)\} = f(X(s))f(X(t))\exp(-V(t))$, yielding the formula. A similar proof gives the expression for $EV^{-1}(R)$.

Proof of Proposition 2. For the first formula, note that

$$\begin{aligned} P\{M \geq m|Y\} &= P\{V^{-1}(R) > S_m|Y\} \\ &= P\{R > V(S_m)|Y\} \\ &= E\{P\{R > V(S_m)|X\}|Y\} \\ &= E\{\exp(-V(S_m))|Y\} \\ &= \prod_{j=0}^{m-1} E\{\exp(-g(Y_j)\beta_j)|Y\} \\ &= \prod_{j=0}^{m-1} \varphi(Y_j, Y_{j+1}, g(Y_j)), \end{aligned}$$

from which the result follows. For the second expression, observe that $E\{\beta_k|Y, M = m\} = E\{\beta_k I(M = m)|Y\}/P\{M = m|Y\}$. To analyze the numerator, note that for $k < m$,

$$\begin{aligned} E\{\beta_k I(M \geq m)|Y\} &= E\{E\{\beta_k I(R > V(S_m))|X\}|Y\} \\ &= E\{\beta_k \exp(-V(S_m))|Y\} \\ &= -\varphi'_k \cdot \prod_{\substack{j=0 \\ j \neq k}}^{m-1} \varphi_j \end{aligned}$$

so that $E\{\beta_k I(M = m) | Y\} = -\varphi'_k \cdot (1 - \varphi_m) \cdot \prod_{j=0}^{m-1} \varphi_j$. This, when combined with the first formula, yields the second identity.

The proof of the third formula follows a similar pattern. We write $E\{V^{-1}(R) - S_M | Y, M = m\} = E\{(V^{-1}(R) - S_M)I(M = m) | Y\} / P\{M = m | Y\}$; again, we handle the denominator using the first formula. For the numerator, we note that $E\{V^{-1}(R)I(M = m) | Y\}$ can be expressed as

$$\begin{aligned} & \int_0^\infty P\{V^{-1}(R) > t; M = m | Y\} dt \\ &= \int_0^\infty P\{R > V(t); R > V(S_m) | Y\} dt - \int_0^\infty P\{R > V(t); R > V(S_{m+1}) | Y\} dt \end{aligned} \quad (A4)$$

By conditioning on X , we find that $P\{R > \max(V(t), V(S_k)) | Y\} = E\{\exp(-\max(V(t), V(S_k))) | Y\}$. So by Lemma 1

$$\begin{aligned} \int_0^\infty P\{R > \max(V(t), V(S_k)) | Y\} dt &= E\left\{\int_0^\infty \exp(-\max(V(t), V(S_k))) dt\right\} \\ &= E\{S_k \exp(-V(S_k)) | Y\} + E\left\{\int_{S_k}^\infty 0 \exp(-V(t)) dt | Y\right\}. \end{aligned} \quad (A5)$$

On the other hand, $E\{S_m I(M = m) | Y\} = E\{S_m I(V(S_m) \leq R < V(S_{m+1})) | Y\} = E\{S_m (\exp(-V(S_m)) - \exp(-V(S_{m+1}))) | Y\}$. Combining this with (A4) and (A5) shows that the numerator equals

$$\begin{aligned} & E\left\{\exp(-V(S_m)) \left(\int_0^{\beta_m} \exp(-g(Y_m)t) dt - \beta_m \exp(-g(Y_m)\beta_m)\right) | Y\right\} \\ &= \prod_{i=0}^{m-1} \varphi_i \left(\frac{1}{g(Y_m)}(1 - \varphi_m) + \varphi'_m\right). \end{aligned}$$

Dividing by the denominator gives the third formula.

Proof of Theorem 4. Standard weak convergence arguments prove that

$$\delta_4(t) = \{\bar{A}_t(0) + \bar{C}_t(0)EK(1)\} + \left[\frac{EC(0)}{1 - EC(1)}(\bar{A}_t(1) - EK(1)(1 - \bar{C}_t(1)))\right] + o_p(t^{-1/2})$$

where $o_p(t^{-1/2})$ represents a process $\chi(t)$ such that $t^{1/2}\chi(t) \Rightarrow 0$. The random time-change results of Section 5 of Glynn and Whitt (1986) can now be applied to the bracketed terms above to obtain the result. (To show that the respective variances are finite, see the proof of Proposition 3.)

Proof of Proposition 3. The regenerative structure of X proves that ($\stackrel{D}{=}$ denotes equality in distribution)

$$D \stackrel{D}{=} A(0) + C(0)K(1)$$

where $(A(0), C(0))$ is independent of $K(1)$. Squaring both sides and taking expectations, we get

$$ED^2 = EA(0)^2 + 2EA(0)C(0)EK(1) + EC(0)^2EK(1)^2. \quad (A6)$$

Since all terms on the right-hand side are positive, we see that H2 implies the finiteness of all the quantities appearing there. We apply the same analysis to $K(1)$:

$$K(1) \stackrel{D}{=} A(1) + C(1)K(2).$$

Using the fact that $K(2) \stackrel{D}{=} K(1)$, we get $EK(1)^2 = (EA(1)^2 + 2EA(1)C(1) \cdot EK(1))(1 - EC(1)^2)^{-1}$. ($EC^2(1) < 1$ by H1). Substituting this into (A6) yields the result, after algebraic simplification.

Proof of Theorem 6. For a) we use (8) and let $\alpha \downarrow 0$. A Taylor expansion gives

$$\begin{aligned} A_\alpha(0) &= \int_0^{T_0} \exp(-\alpha V(t)) f(X(t)) dt \\ &= \int_0^{T_0} \left[1 - \alpha V(t) + \frac{\alpha^2}{2} V^2(t) e^{-\gamma V(t)} \right] f(X(t)) dt \end{aligned}$$

where $\gamma = \gamma(\alpha) \in (0, \alpha)$. Since $V^2(t) \exp(-\gamma(\alpha)V(t))f(X(t)) \leq V^2(t)f(X(t))$ uniformly in α and

$$\int_0^{T_0} V^2(t) f(X(t)) dt = Y_0(f)Y_0(g)^2$$

is integrable by H9, it follows from the dominated convergence theorem that

$$EA_\alpha(0) = EY_0(f) - \alpha E \left\{ \int_0^{T_0} V(t) f(X(t)) dt \right\} + \frac{\alpha^2}{2} E \left\{ \int_0^{T_0} V^2(t) f(X(t)) dt \right\} + o(\alpha^2). \quad (A7)$$

Similarly, one can show that

$$EC_\alpha(0) = 1 - \alpha EY_0(y) + \frac{\alpha^2}{2} EY_0(y)^2 + o(\alpha^2).$$

Corresponding expressions for $EA_\alpha(1)$ and $EC_\alpha(1)$ lead immediately to a). For b), h), i), we use Propositions 3 and 4 and arguments similar to the above. Relation c) can be found in Glynn and Whitt (1987).

For e), observe that

$$V_\alpha^{-1}(R) \stackrel{D}{=} L_\alpha(0) + I_\alpha(0)Q_\alpha(1) \quad (A8)$$

where $L_\alpha(0) = V_\alpha^{-1}(R) \wedge T_0$, $I_\alpha(0) = I(V_\alpha^{-1}(R) > T_0)$, $Q_\alpha(1) = V_\alpha^{-1}(r) - T_0$, and $(L_\alpha(0), I_\alpha(0))$ is independent of $Q_\alpha(1)$. Furthermore, on $\{V_\alpha^{-1}(R) > T_0\}$,

$$Q_\alpha(1) \stackrel{D}{=} L_\alpha(1) + I_\alpha(1)Q_\alpha(2) \quad (A9)$$

where $L_\alpha(1) = (V_\alpha^{-1}(R) \wedge T_1) - T_0$, $I_\alpha(1) = I(V_\alpha^{-1}(R) > T_1)$, $Q_\alpha(2) = V_\alpha^{-1}(R) - T_1$, $(L_\alpha(1), I_\alpha(1))$ is independent of $Q_\alpha(2)$, and the distribution of $Q_\alpha(i)$ conditional on $\{V_\alpha^{-1}(R) > T_i\}$ is independent of i ($i = 0, 1$). Taking expectations in (A8) and (A9) and using the independence leads to an expression similar to (8). One then expands the expectations in a manner similar to (A7) to obtain e). Results d), f), and g) are proved using decompositions analogous to (A8) and (A9), followed by Taylor expansions for small α .

REFERENCES

- BRATLEY, P., B. L. FOX, and L. E. SCHRAGE, *A Guide to Simulation*, Springer-Verlag, New York, 1983.
- CHUNG, K. L., *A Course in Probability Theory*, Academic Press, New York, 1974.
- FOX, B. L. and P. W. GLYNN, "Discrete-time Conversion for Simulating Semi-Markov Processes," *Operations Res. Letters* (1986), 191-196.
- FOX, B. L. and P. W. GLYNN, "Replication Schemes for Limiting Expectations," technical report, University of Montreal, 1987.
- GLYNN, P. W. and W. WHITT, "The Efficiency of Simulation Estimators," technical report, AT&T Bell Laboratories, Holmdel, New Jersey, 1986.
- GLYNN, P. W. and W. WHITT, "Limit Theorems for State-Dependent Discounted Costs," forthcoming technical report.
- HAMMERSLEY, J. M. and D. C. HANDSCOMB, *Monte Carlo Methods*, Chapman and Hall, New York, 1964.
- VEINOTT, JR., A. F., "Discrete Dynamic Programming with Sensitive Discount Optimality Criteria," *Ann. Math. Stat.* 40 (1969), 1635-1660.
- WHITT, W., "Stochastic Abelian and Tauberian Theorems," *Z. Wahrscheinlichkeitstheorie verw. Geb.* 22 (1972), 251-167.
- WHITT, W., "Planning Queueing Simulations," forthcoming technical report.

A193 584

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report No. 23	2. GOVT ACCESSION NO. N/A	3. RECIPIENT'S CATALOG NUMBER N/A
4. TITLE (and Subtitle) SIMULATING DISCOUNTED COSTS		5. TYPE OF REPORT & PERIOD COVERED Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Bennett L. Fox and Peter W. Glynn		8. CONTRACT OR GRANT NUMBER(s) DAAG29-84-K-0030
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Operations Research Stanford University Stanford, CA 94305-4022		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS N/A
11. CONTROLLING OFFICE NAME AND ADDRESS U.S. Army Research Office Post Office Box 12211 Research Triangle Park, NC 27709		12. REPORT DATE August 1987
		13. NUMBER OF PAGES 21 pp.
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) NA		
18. SUPPLEMENTARY NOTES The view, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision, unless so designated by other documentation.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Discounted costs, simulation, semi-Markov process, regenerative process		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) PLEASE SEE OTHER SIDE		

20. (Continued.)

We numerically estimate, via Monte Carlo simulation, the expected infinite-horizon discounted cost d of running a stochastic system; we permit the discount rate to be state dependent. By exploiting various types of stochastic structure, we beat the naive strategy of estimating a finite-horizon approximation to d . Efficient estimators are obtained for systems which are semi-Markov and/or regenerative. These estimators are then ranked with respect to asymptotic variance as a function of computer-time budget and discount rate.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

DATE
FILMED
8