

AD-A194 289


POISSON FUNCTIONALS OF MARKOV PROCESSES AND QUEUEING
NETWORKS(U) NORTH CAROLINA UNIV AT CHAPEL HILL DEPT OF
STATISTICS R F SERFOZO DEC 87 TR-218 AFOSR-TR-88-0352
F49620-85-C-0144


1/1

UNCLASSIFIED

F/G 12/3

NL





DTIC FILE COPY ②


AD-A194 289

REPORT DOCUMENTATION PAGE

Unclassified			1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE			Approved for public release; distribution unlimited.	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. 218			5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 88-0352	
6a. NAME OF PERFORMING ORGANIZATION University of North Carolina		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION AFOSR/NM	
6c. ADDRESS (City, State, and ZIP Code) Statistics Dept. 321-A Phillips Hall 039-A Chapel Hill, NC 27514			7b. ADDRESS (City, State, and ZIP Code) AFOSR/NM Bldg 410 Bolling AFB DC 20332-6448	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION AFOSR		8b. OFFICE SYMBOL (If applicable) NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR No. F49620 85C 0144.	
8c. ADDRESS (City, State, and ZIP Code) AFOSR/NM Bldg 410 Bolling AFB DC 20332-6448			10. SOURCE OF FUNDING NUMBERS	
			PROGRAM ELEMENT NO. 61102F	PROJECT NO. 2304
			TASK NO. AS	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Poisson functionals of Markov processes and queueing networks.				
12. PERSONAL AUTHOR(S) Serfozo, R.F.				
13a. TYPE OF REPORT preprint		13b. TIME COVERED FROM 9/87 TO 8/88		14. DATE OF REPORT (Year, Month, Day) December 1987
15. PAGE COUNT 24				
16. SUPPLEMENTARY NOTATION N/A				
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	Key Words & Phrases: Poisson process, multivariate compound Poisson process, functionals of Markov processes, queueing networks, time reversal.	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) We present conditions under which a point process of certain jump times of a Markov process is a Poisson process. One result is that if the Markov process is stationary and the compensator of the point process in reverse time has a constant intensity a , then the point process is Poisson with rate a . A classical example is that the output flow from a M/M/1 queueing system is Poisson. We also present similar Poisson characterizations of more general marked point process functionals of a Markov process. These results yield easy-to-use criteria for a collection of such processes to be multi-variate Poisson or marked Poisson with a specified dependence or independence. We give several applications to queueing systems, and indicate how our results extend to functionals of non-Markovian processes.				
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT ☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT ☐ DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified/unlimited	
22a. NAME OF RESPONSIBLE INDIVIDUAL Maj. Brian Woodruff			22b. TELEPHONE (Include Area Code) (202) 767-5026	22c. OFFICE SYMBOL NM

CENTER FOR STOCHASTIC PROCESSES

Department of Statistics
University of North Carolina
Chapel Hill, North Carolina



POISSON FUNCTIONALS OF MARKOV
PROCESSES AND QUEUEING NETWORKS

by

R.F. Serfozo

Technical Report No. 218

December 1987

141. T. Hsing, On the intensity of crossings by a shot noise process, July 86. Adv. Appl. Probab. 10, 1987, 743-745.
142. V. Mandrekar, On a limit theorem and invariance principle for symmetric statistics, July 86.
143. O. Kallenberg, Some new representations in bivariate exchangeability, July 86.
144. B.G. Nguyen, Correlation length and its critical exponents for percolation processes, July 86. J. Stat. Phys., to appear.
145. G. Kallianpur and V. Perez-Abreu, Stochastic evolution equations with values on the dual of a countably Hilbert nuclear space, July 86. Appl. Math. Optimization, to appear.
146. B.G. Nguyen, Fourier transform of the percolation free energy, July 86. Probab. Theor. Rel. Fields, to appear.
147. A.M. Hasofer, Distribution of the maximum of a Gaussian process by Monte Carlo, July 86. J. Sound Vibration, 112, 1987, 283-293.
148. T. Norberg, On the existence of probability measures on continuous semi-lattices, Aug. 86.
149. G. Samorodnitsky, Continuity of Gaussian processes, Aug. 86. Ann. Probability, to appear.
150. T. Hsing, J. Husler, and M.R. Leadbetter, Limits for exceedance point processes, Sept. 86. Prob. Theory and Related Fields, to appear.
151. S. Ombani, Random filters which preserve the stability of random inputs, Sept. 86. Adv. Appl. Probability, 1987, to appear.
152. O. Kallenberg, On the theory of conditioning in point processes, Sept. 86. Proceedings for the First World Congress of the Bernoulli Society, Tashkent, 1986.
153. G. Samorodnitsky, Local moduli of continuity for some classes of Gaussian processes, Sept. 86.
154. V. Mandrekar, On the validity of Beurling theorems in polydiscs, Sept. 86.
155. R.F. Serfozo, Extreme values of queue lengths in M/G/1 and GI/M/1 systems, Sept. 86.
156. B.G. Nguyen, Gap exponents for percolation processes with triangle condition, Sept. 86. J. Statist. Physics, 49, 1987.
157. G. Kallianpur and R. Wolpert, Weak convergence of stochastic neuronal models, Oct. 86. Stochastic Methods in Biology, M. Kimura et al., eds., Lecture Notes in Biomathematics, 70, Springer, 1987, 116-145.
158. G. Kallianpur, Stochastic differential equations in duals of nuclear spaces with some applications, Oct. 86. Inst. of Math. & Its Applications, 1986.
159. G. Kallianpur and R.L. Karandikar, The filtering problem for infinite dimensional stochastic processes, Jan. 87. Proc. Workshop on Stochastic Differential Systems, Stochastic Control Theory & Applications, Springer, to appear.
160. V. Perez-Abreu, Multiple stochastic integrals and their extensions to nuclear space valued Wiener processes, Oct. 86. Appl. Math. Optimization, to appear.
161. R.L. Karandikar, On the Feynman-Kac formula and its applications to filtering theory, Oct. 86. Appl. Math. Optimization, to appear.
162. R.L. Taylor and T.-C. Hu, Strong laws of large numbers for arrays of rowwise independent random elements, Nov. 86.
163. M. O'Sullivan and T.R. Fleming, Statistics for the two-sample survival analysis problem based on product limit estimators of the survival functions, Nov. 86.
164. F. Avram, On bilinear forms in Gaussian random variables, Toeplitz matrices and Parseval's relation, Nov. 86.
165. D.B.H. Cline, Joint stable attraction of two sums of products, Nov. 86. J. Multivariate Anal., to appear.
166. R.J. Wilson, Model fields in crossing theory—a weak convergence perspective, Nov. 86.
167. D.B.H. Cline, Consistency for least squares regression estimators with infinite variance data, Dec. 86.
168. L.L. Campbell, Phase distribution in a digital frequency modulation receiver, Nov. 86.
169. B.G. Nguyen, Typical cluster size for 2-dim percolation processes, Dec. 86. J. Statist. Physics, to appear.
170. H. Oodaira, Freidlein-Wentzell type estimates for a class of self-similar processes represented by multiple Wiener integrals, Dec. 86.
171. J. Nolan, Local properties of index- β stable fields, Dec. 86. Ann. Probability, to appear.
172. R. Menich and R.F. Serfozo, Optimality of shortest queue routing for dependent service stations, Dec. 86.
173. F. Avram and M.S. Taqqu, Probability bounds for M-Skorohod oscillations, Dec. 86.
174. F. Moricz and R.L. Taylor, Strong laws of large numbers for arrays of orthogonal random variables, Dec. 86.
175. G. Kallianpur and V. Perez-Abreu, Stochastic evolution driven by nuclear space valued martingales, Apr. 87.
176. E. Merzbach, Point processes in the plane, Feb. 87.
177. Y. Kasahara, M. Maejima and W. Vervaat, Log fractional stable processes, March 87.

Poisson Functionals of Markov Processes
and Queueing Networks

by

Richard F. Serfozo
University of North Carolina at Chapel Hill
and
Georgia Institute of Technology

ABSTRACT

We present conditions under which a point process of certain jump times of a Markov process is a Poisson process. One result is that if the Markov process is stationary and the compensator of the point process in reverse time has a constant intensity a , then the point process is Poisson with rate a . A classical example is that the output flow from a M/M/1 queueing system is Poisson. We also present similar Poisson characterizations of more general marked point process functionals of a Markov process. These results yield easy-to-use criteria for a collection of such processes to be multi-variate Poisson or marked Poisson with a specified dependence or independence. We give several applications to queueing systems, and indicate how our results extend to functionals of non-Markovian processes.

Keywords and phrases: Poisson process, multivariate compound Poisson process, functionals of Markov processes, queueing networks, time reversal.

This research was sponsored in part by Air Force Office of Scientific Research contracts S4-0367 and F49620 85 C 0144.

Accession For	
NTIS	GRA&I ☒
DTIC	TAB ☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	FOR INSPECTION

1. Introduction

There are a variety of point processes associated with the jump times of a Markov process. For instance, in a Markovian network of queues, one might be interested in the point process of times at which units move between two sectors of the network. More generally, if the network has synchronous movements of items, one might be interested in the marked point process of the times at which batches of units move between two sectors and the numbers of units in the batches (the batch size being the "mark" of the time of the movement). One can formulate such a point process as a functional of the Markov process representing the network. The typical aim is to describe the behavior of the point process in terms of the characteristic of the Markov process. Some immediate questions in this regard are: Is such a point process Poisson (or marked Poisson)? Is a collection of these point processes multi-variate Poisson (or marked Poisson); and what are the dependencies, if any, among them?

These are the issues this study addresses. We begin in Section 2 by presenting conditions under which a point process of certain jump times of a Markov process is a Poisson process. It is well known that a simple point process on the real line is Poisson with rate λ if its compensator has the constant intensity λ (Theorem 2.3(i)). We present a reverse-time version of this (Theorem 2.3(ii)). It says that if the Markov process is stationary and the compensator of the point process in reverse time has the constant intensity λ , then the point process is Poisson with rate λ .

This is an easy-to-use criterion for establishing whether a point process of jump times is Poisson. We also give necessary conditions for this Poissonness. In Section 3, we present similar conditions under which more general functionals of a pure jump Markov process are marked Poisson processes. Some of our results overlap those of Melamed (1979), Brémaud (1981), Variaya and Walrand (1981), and Disney and Kiessler (1987). Melamed and Disney and Kiessler derive their results using a Markov renewal argument and Brémaud, Variaya and Walrand use a filtering argument. We use a simpler approach based on the notion of the time reversibility of the compensator of a point process. This approach lays bare the characteristics underlying the Poisson property, and it readily extends to the more general settings in Sections 3-5.

The applications in this area have been primarily for queueing systems. Burke (1956) and Reich (1957) showed that, in a stationary M/M/1 queueing system, the output flow is a Poisson process with the same rate as the Poisson input flow. Similarly, the exit flows from the queues in a Jackson network are independent Poisson processes; this is discussed in the references in the preceding paragraphs and in Kelly (1979), Disney and König (1985) and Whittle (1986). In Sections 2-4, we discuss applications identifying Poisson, compound Poisson and multivariate-Poisson flows in single queues and in queueing networks with dependent nodes. We end in Section 5 by indicating how our results extend to Markov processes with general state spaces and to functionals of semi-Markovian processes.

2. Jump Times of a Markov Process that Form a Poisson Process

Let $X = \{X_t : t \in \mathbb{R}\}$ be a Markov process with a countable state space $\mathcal{I}$ and transition rates

$$q(x,y) = \lim_{t \downarrow 0} P\{X_t = y \mid X_0 = x\}/t, \quad x \neq y,$$

and $q(x,x) = 0$. We indicate later how our results extend to a general state space. We adopt the standard assumption that

$$q(x) := \sum_y q(x,y) < \infty, \quad x \in \mathcal{I},$$

and that each sample path of X is right continuous and has a finite number of jumps in any finite time period. Then the sojourn time of X in a state x is exponential with mean $q(x)^{-1}$ and, at the end of the sojourn, X jumps to some state y with probability $q(x,y)/q(x)$, $y \in \mathcal{I}$. For convenience, we assume that X is irreducible.

We shall study the point process N on $\mathbb{R}$ defined by

$$(2.1) \quad N(\Lambda) = \sum_{t \in \Lambda} f(X_{t-}, X_t)$$

where Λ is a Borel set in $\mathbb{R}$ and $f: \mathcal{I} \times \mathcal{I} \rightarrow \{0,1\}$ with $f(x,x) = 0$, $x \in \mathcal{I}$.

The $N(\Lambda)$ is the number of jumps of X from some x to some y for which $f(x,y) = 1$ in the time period Λ . Any such f is an indicator function $f(x,y) = 1((x,y) \in S)$ of a subset S of $\mathcal{I} \times \mathcal{I}$ that does not contain pairs of identical values; the $N(\Lambda)$ would then be the number of transitions of X in the period Λ that take place in the transition set S . Clearly $N(\Lambda)$ is finite when Λ is bounded and it may be infinite when Λ is unbounded.

We shall frequently use the function

$$(2.2) \quad \alpha(x) = \sum_y q(x,y)f(x,y), \quad x \in \mathcal{I}.$$

First note that the mean measure of N is

$$(2.3) \quad EN(\Lambda) = \int_{\Lambda} E\alpha(X_t) dt$$

which is Levy's formula (see for instance Benveniste and Jacod (1973)).

When X is stationary, an easy check shows that N is stationary (i.e. for any $\Lambda_1, \dots, \Lambda_n$, the distribution of $N(\Lambda_1+t), \dots, N(\Lambda_n+t)$ is independent of t). Consequently, $EN(s,t] = a(t-s)$, $s < t$ in $\mathbb{R}$, where

$$(2.4) \quad a = EN(0,1] = \sum_{x,y} \pi(x)q(x,y)f(x,y)$$

and $\pi(x) = P\{X_t=x\}$ is the equilibrium (or stationary) distribution of X .

Here $0 \leq a \leq \infty$. Also, with probability one, $t^{-1}N(s,s+t] \rightarrow a$ as $t \rightarrow \infty$.

One can use N to model a variety of event occurrences of X by appropriate selections of f . For instance, suppose X takes values in $\mathcal{X} = \mathbb{Z}_+^m$, the m -dimensional vectors with nonnegative integer-valued entries. Then N records the downward jumps of X when $f(x,y) = 1$ iff $x \neq y$ and $x_j \geq y_j$, $j=1, \dots, m$. Similarly, N records the jumps of X at which the maximum component does not change when $f(x,y) = 1$ iff $x \neq y$ and $\max_j x_j = \max_j y_j$.

We shall investigate conditions under which N is a Poisson process.

We say that the future of N is independent of the past of X , denoted

$N_+ \perp\!\!\!\perp X_-$, if $\{N(\Lambda): \Lambda \subset [t, \infty)\}$ is independent of $\{X_u: u \leq t\}$, $t \in \mathbb{R}$.

($\{X_u: u \leq t\}$ can be replaced simply by X_t since X is Markovian).

Similarly, $N_- \perp\!\!\!\perp X_+$ denotes that the past of N is independent of the

future of X . Our first result is that these conditions are sufficient

for N to be a Poisson process; the N might be a non-stationary Poisson

process. We write " N is $\mathcal{P}(a)$ " to mean that N is a stationary

(time-homogeneous) Poisson process with rate a . The degenerate case $a = 0$ corresponds to $N = 0$; the case $a = \infty$ is not possible.

Theorem 2.1. If $N_+ \perp\!\!\!\perp X_-$ or $N_- \perp\!\!\!\perp X_+$, then N is a Poisson process. In this case, N is $\mathcal{P}(a)$ if and only if $Ea(X_t) = a$, $t \in \mathbb{R}$.

Proof. The N is a Poisson process if it is simple (i.e., $N(\{t\}) = 0$ or 1 , $t \in \mathbb{R}$), it has no fixed atoms and has independent increments (i.e., $N(A_1), \dots, N(A_n)$ are independent for disjoint $A_1, \dots, A_n$); see p. 58 of Kallenberg (1983). Since the probability is zero that X has a jump at any specified time, it follows that N is simple with no fixed atoms. Now suppose $N_+ \perp\!\!\!\perp X_-$. Then, for any $s < t$ in $\mathbb{R}$,

$$\begin{aligned} P\{N(s, t] = n \mid N(A) : A \subset (-\infty, s]\} \\ = E[P\{N(s, t] = n \mid X_r : r \leq s\}] = P\{N(s, t] = n\}. \end{aligned}$$

Thus N has independent increments and hence N is Poisson. This conclusion also follows when $N_- \perp\!\!\!\perp X_+$ since

$$\begin{aligned} P\{N(s, t] = n \mid N(A) : A \subset [t, \infty)\} \\ = E[P\{N(s, t] = n \mid X_u : u \geq t\}] = P\{N(s, t] = n\}. \end{aligned}$$

The second assertion on the stationarity of N follows from (2.3). $\square$

Remark 2.2. From the proof, it is clear that the first assertion of Theorem 2.1 is true for any pure jump stochastic process X ; the Markovian property is used only for the second assertion. Also, both assertions are true when X is a Markov process that is not time homogeneous (its transition rates are time dependent). Melamed (1979) showed that $N_+ \perp\!\!\!\perp X_-$ implies that N is Poisson; his argument relies on the property that X and $\{(N(s, t], X_t) : t \geq s\}$ are Markovian.

We now discuss conditions on the parameters of the Markov process X under which N is $\mathcal{P}(a)$. When X is stationary with equilibrium distribution π , we shall frequently use the function

$$(2.5) \quad \alpha^*(x) = \pi(x)^{-1} \sum_y \pi(y) q(y, x) f(y, x), \quad x \in \mathcal{A}.$$

Theorem 2.3. (i) If $\alpha(x) = a$, $x \in \mathcal{A}$, then $N_+ \perp\!\!\!\perp X_-$ and N is $\mathcal{P}(a)$. (ii) If X is stationary with equilibrium distribution π , and $\alpha^*(x) = a$, $x \in \mathcal{A}$, then $N_- \perp\!\!\!\perp X_+$ and N is $\mathcal{P}(a)$.

Proof. Assertion (i) is a special case of Theorem 18.9 in Liptser and Shirayev (1978) (or Theorem T5 in Brémaud (1981)), which says that a simple point process on $\mathbb{R}$ is $\mathcal{P}(a)$ if its compensator has the non-random intensity a . In our setting, for each $s \in \mathbb{R}$, the process

$$(2.6) \quad M_t = N(s, s+t] - \int_s^{s+t} \alpha(X_u) du$$

is an $\mathcal{F}_t$ -martingale, where $\mathcal{F}_t = \sigma(X_u : s \leq u \leq s+t)$; the process $A_t = \int_s^{s+t} \alpha(X_u) du$ is the compensator of $N(s, s+t]$ and $\alpha(X_t)$ is the intensity of this compensator.

To prove assertion (ii), consider the process $X_t^* = \bar{X}_{-t}$, $t \in \mathbb{R}$, where $\bar{X}_t = X_{t-}$. The X^* is the right-continuous time-reversal of X . Each sample path of X^* is the same as a sample path of X traversed in the opposite direction, and vice versa. Since X is stationary, it follows that X^* is an irreducible, stationary Markov process and its transition rates are

$$q^*(x, y) = \pi(x)^{-1} \pi(y) q(y, x), \quad x, y \in \mathcal{A}.$$

Define the point process N^* on $\mathbb{R}$ by

$$N^*(\Lambda) = \sum_{t \in \Lambda} f^*(X_{t-}^*, X_t^*)$$

where $f^*(x, y) = f(y, x)$, $x, y \in \mathcal{A}$. Clearly $N^*(\Lambda) = N(-\Lambda)$, for each Λ , and

so N^* is the time-reversal of N . Consequently, N is $\mathcal{P}(a)$ if and only if N^* is $\mathcal{P}(a)$.

Now observe that $a^*(x) = \sum_y q^*(x,y) f^*(x,y)$. That is, a^* is the function a for the processes X^*, N^* . Then an application of (i) to X^*, N^* says that $N_+^* \perp\!\!\!\perp X_-^*$ and N^* is $\mathcal{P}(a)$. But these statements are equivalent to the respective statements that $N_- \perp\!\!\!\perp X_+$ and N is $\mathcal{P}(a)$, since X^*, N^* are the time reversals of X, N . $\square$

Remarks. (1) In Theorem 2.3, statement (i) is the well-known property of Markov processes that N is $\mathcal{P}(a)$ if the intensity $a(X_t)$ of N 's compensator equals the constant a . Statement (ii) is simply a reversed time version of (i): it is (i) in terms of X and N viewed in reverse time. The a^* is the reversed time version of a in the sense that $a^*(X_t^*)$ is the intensity of the compensator of N in reverse time. Also, $N_- \perp\!\!\!\perp X_+$ in (ii) is the time reversal of $N_+ \perp\!\!\!\perp X_-$ in (i). (2) Note that the process X in (i) may be transient, recurrent or non-stationary; but in (ii), X must be stationary. (3) In consulting the references for Theorem 2.3(i), one can see that this result is true for a non-Markovian process X for which $a: \mathcal{A} \rightarrow R_+$ is a function such that the process M_t in (2.6) is an $\mathcal{F}_t$ -martingale. Similarly, it follows that Theorem 2.3(ii) is true for a non-Markovian process X if $a^*: \mathcal{A} \rightarrow R_+$ is a function such that, for each $s \in R$,

$$M_t^* = N[s-t, s) - \int_{s-t}^s a^*(X_u) du$$

is an $\mathcal{F}_t^*$ -martingale where $\mathcal{F}_t^* = \sigma(X_u: s-t \leq u < s)$.

In applications, the Poisson property of N due to $\alpha(x) = a, x \in \mathcal{I}$, is usually foreseen, while the Poisson property of N due to $\alpha^*(x) = a, x \in \mathcal{I}$, might not be anticipated. Here is an example.

Example 2.4. M/M/1 and Batch Service Queues. Suppose the Markov process X has the state space Z_+ and transition rates

$$q(n, n+1) = \lambda \quad n \in Z_+$$

$$q(n, n-K) = \mu \quad n \geq K$$

$$q(n, 0) = \mu \quad n < K$$

where λ, μ and K are positive. This process represents the number of customers in a queueing system in which customers arrive singly at the rate λ and are served in batches such that when $n \geq K$ customers are in the system, then batches of K customers depart at the rate μ ; and when $n < K$ customers are present, then all of the customers depart at the rate μ . When $K = 1$, this is the M/M/1 queueing system.

Implicit in the description of this queueing system, the point process N of customer arrivals is $\mathcal{P}(\lambda)$, regardless of whether X is transient or recurrent. Indeed, this follows from Theorem 2.3 (i) since N is defined by (2.1) with $f(n, n') = 1$ iff $n' = n + 1$ and

$$\alpha(n) = \sum_{n'} q(n, n') f(n, n') = q(n, n+1) = \lambda.$$

Now, suppose that N denotes the point process of times at which batches of size K depart from the system. This is defined by (2.1) with $f(n, n') = 1$ iff $n' = n - K$ and $n \geq K$. In this case,

$$\alpha(n) = q(n, n-K) 1(n \geq K) = \mu 1(n \geq K).$$

This depends on n , and so Theorem 2.3 (i) does not ensure that N is Poisson. However, assume that X is stationary. Necessarily, $\lambda < \mu K$ and the equilibrium distribution of X is

$$\pi(n) = r^n(1 - r), \quad n \geq 0,$$

where $r \in (0,1)$ is the unique root of

$$\mu r^{K+1} - (\lambda + \mu)r + \lambda = 0$$

(see Section 3.2 of Gross and Harris (1985)). Clearly

$$\begin{aligned} \alpha^*(n) &= \pi(n)^{-1} \sum_{n'} \pi(n') q(n', n) f(n', n) \\ &= \pi(n)^{-1} \pi(n + K) q(n + K, n) \\ &= \mu r^K = \mu + \lambda(1 - 1/r). \end{aligned}$$

Thus, we conclude by Theorem 2.3 (ii) that the process N is $\mathcal{P}(\mu + \lambda(1 - 1/r))$. One would probably not anticipate this result from the description of the process, or even from earlier work in this area. For the special case in which X is the $M/M/1$ queue, we have $K = 1$, $r = \lambda/\mu$, and so N , which is the departure process, is $\mathcal{P}(\lambda)$. Burke (1956) and Reich (1957) were the first ones to prove this.

Example 2.5. Queues With Compound Poisson Arrivals and Poisson Departures. Suppose the Markov process has the state space Z_+ and transition rates

$$\begin{aligned} q(n, n + m) &= \lambda_n p^{m-1} (1 - p) & m \geq 1, \quad n \in Z_+ \\ q(n, n - 1) &= \mu_n & n \geq 1 \end{aligned}$$

where λ_n, μ_n are positive and $0 < p < 1$. This process represents the number of customers in a queueing system in which batches of customers arrive at the rate λ_n when n customers are present and the number of customers in a batch has a geometric distribution with parameter p . The customers depart at the rate μ_n when n are in the system. The equilibrium distribution of X is

$$\pi(n) = \pi(0) \lambda_0^{-1} \mu_1^{-1} \dots \mu_n^{-1} \prod_{k=1}^{n-1} (\lambda_k + p\mu_k), \quad n \geq 1,$$

provided the sum of these terms over n is finite, which we assume is true (see Kook (1988)).

Suppose that N is the point process of customer departures. This is defined by (2.1) with $f(n, n') = 1$ iff $n' = n - 1$. Clearly,

$$\begin{aligned}\alpha^*(0) &= \pi(0)^{-1} \pi(1) q(1, 0) = \lambda_0 \\ \alpha^*(n) &= \pi(n)^{-1} \pi(n+1) q(n+1, n) = \lambda_n + p\mu_n, \quad n \geq 1.\end{aligned}$$

Thus, if X is stationary and $\lambda_0 = a$ and $\lambda_n + p\mu_n = a$, $n \geq 1$, then the departure process N is $\mathcal{P}(a)$.

We end this section with elaborations on Theorem 2.3 that establish necessary conditions for N to be Poisson.

Theorem 2.6. When X is recurrent, the following statements are equivalent.

- (i) $N_+ \perp\!\!\!\perp X_-$ and N is $\mathcal{P}(a)$.
- (ii) $\alpha(x) = a$, $x \in \mathcal{X}$.
- (iii) $N_+ \perp\!\!\!\perp X_-$ and $EN(s, t] = a(t-s)$, $s < t$ in $\mathbb{R}$.
- (iv) $E[\alpha(X_t) \mid X_s = x] = a$, $x \in \mathcal{X}$, $s < t$ in $\mathbb{R}$.

Proof. Theorem 2.3 ensures that (ii) implies (i). Clearly (i) implies (iii). Now, if (iii) holds, then, for $s < t$ in $\mathbb{R}$,

$$\begin{aligned}a(t-s) &= EN(s, t] = E[N(s, t] \mid X_s = x] \\ &= \int_s^t E[\alpha(X_u) \mid X_s = x] du\end{aligned}$$

the last equality being Lévy's formula. Taking the derivative of this with respect to t yields (iv). Finally, if equation (iv) holds, then taking the derivative of it with respect to t and letting $t \downarrow s$, we obtain

$$\sum_y \alpha(y) q(x, y) / q(x) = \alpha(x) \quad \text{for each } x.$$

This says that α is a harmonic function of the Markov matrix $\{q(x,y)/q(x)\}$. Now, this matrix is recurrent under the assumption that X is recurrent. But we know (see for instance Section 7.2 of Çinlar (1975)) that harmonic functions of irreducible, recurrent Markov chains are constant. This observation and equation (iv) imply (ii). $\square$

Theorem 2.7. When X is stationary with equilibrium distribution π , the following statements are equivalent.

- (i) $N_- \perp\!\!\!\perp X_+$ and N is $\mathcal{P}(a)$.
- (ii) $\alpha^*(x) = a, \quad x \in \mathcal{X}$.
- (iii) $N_- \perp\!\!\!\perp X_+$ and $EN(0,1] = a$.
- (iv) $E[\alpha^*(X_s) \mid X_t = x] = a, \quad x \in \mathcal{X}, s < t \text{ in } \mathbb{R}$.
- (v) $\alpha^*(x) = \sum_y \alpha(y)\pi(y) = a, \quad x \in \mathcal{X}$.

Proof. The equivalence of (i) - (iv) follows from Theorem 2.6 applied to the reversed-time processes N^*, X^* defined in the proof of Theorem 2.3. Furthermore, (v) obviously implies (ii). And (ii) implies (v) since N is $\mathcal{P}(a)$ by (i) and by (2.4), we have

$$a = EN(0,1] = \int_0^1 E\alpha(X_t)dt = \sum_y \alpha(y)\pi(y).$$

Thus (i) - (v) are equivalent statements. $\square$

Remark 2.8. Statement (v), relating α^* to α , is the "independence condition" that is the focus of Melamed (1979), Brémaud (1981) and Varaiya and Walrand (1981). Their main result, which is implicit in Theorem 2.7, is as follows: If X is stationary with equilibrium distribution π , then $N_- \perp\!\!\!\perp X_+$ if and only if $\alpha^*(x) = \sum_y \alpha(y)\pi(y), \quad x \in \mathcal{X}$.

In this case, N is $\mathcal{P}(a)$ where a is the preceding sum.

Although we are considering processes N and X defined on the entire real line, our results herein also apply to such processes defined on any time interval I in R . Indeed, consider the process extended to all of R and then the results apply to these extended processes and hence to their restrictions to I .

3. General Poisson Functionals of Markov Processes

We have been studying the point process N that records the times at which the Markov process X jumps from some x to some y with $f(x,y) = 1$. We shall now study a more general point process that records additional information at these jumps. Suppose that h is a function from $\mathcal{X} \times \mathcal{X}$ to some space $\mathcal{X}'$. Consider the marked point process M on $R \times \mathcal{X}'$ defined by

$$(3.1) \quad M(A \times B) = \sum_{t \in A} f(X_{t-}, X_t) 1(h(X_{t-}, X_t) \in B)$$

where f is as above, $A \times B$ is in the σ -field of $R \times \mathcal{X}'$ and $1(\cdot \in B)$ is the indicator function of B . This M is a functional of X that records the "mark" $h(X_{t-}, X_t)$ at each jump time t of X at which $f(X_{t-}, X_t) = 1$. As a point process, M is simple and $M(\{t\} \times \mathcal{X}') = 0$ or 1 , $t \in R$. Also, $M(A \times \mathcal{X}') = N(A)$ records the times at which M records the marks.

In this section, we identify conditions on X under which M is Poisson with $EM((s,t] \times B) = a(t-s)F(B)$, where $a \geq 0$ and F is a probability measure on $\mathcal{X}'$. In this case, the process N of occurrence times of the marks of M , is $\mathcal{P}(a)$, and each mark has the distribution F . We call such an M a marked Poisson process with rate a and mark distribution F , and we simply say M is $\mathcal{MP}(a,F)$.

The approach we used in Section 2 for deriving Poisson characterizations of N readily extends to yield marked Poisson characterizations of M . The following results are analogues of Theorems 2.1, 2.3 and 2.7; the analogue of Theorem 2.6 can be seen from Theorem 3.3 and hence is not displayed. Their proofs are omitted since they follow the same line of reasoning as their counterparts in Section 2. Here we let

$$\alpha(x, B) = \sum_y q(x, y) f(x, y) I(h(x, y) \in B)$$

$$\alpha^*(x, B) = \pi(x)^{-1} \sum_y \pi(y) q(y, x) f(y, x) I(h(y, x) \in B).$$

Theorem 3.1. If $M_+ \perp\!\!\!\perp X_-$ or $M_- \perp\!\!\!\perp X_+$, then M is a Poisson process. In this case, M is $\mathcal{MP}(a, F)$ if and only if $E\alpha(X_t, B) = aF(B)$, for each t and B .

Theorem 3.2. (i) If $\alpha(x, B) = aF(B)$, for each x and B , then $M_+ \perp\!\!\!\perp X_-$ and M is $\mathcal{MP}(a, F)$. (ii) If $\alpha^*(x, B) = aF(B)$, for each x and B , then $M_- \perp\!\!\!\perp X_+$ and M is $\mathcal{MP}(a, F)$.

Theorem 3.3. When X is stationary with equilibrium distribution π , then the following statements are equivalent.

- (i) $M_- \perp\!\!\!\perp X_+$ and M is $\mathcal{MP}(a, F)$.
- (ii) $\alpha^*(x, B) = aF(B)$, for each x and B .
- (iii) $M_- \perp\!\!\!\perp X_+$ and $EM((0, 1] \times B) = aF(B)$ for each x and B .
- (iv) $E[\alpha^*(X_s, B) \mid X_t = x] = aF(B)$, for each x , B and $s < t$ in $\mathbb{R}$.
- (v) $\alpha^*(x, B) = \sum_y \alpha(y, B) \pi(y) = aF(B)$, for each x and B .

Point processes of the form (3.1) are useful for representing multivariate and compound point processes. Indeed, suppose one is interested in the n -dimensional compound point process $(M_1, \dots, M_n)$ defined by

$$(3.2) \quad M_j(\Lambda) = \sum_{t \in \Lambda} f(X_{t-}, X_t) h_j(X_{t-}, X_t)$$

where $h_j: \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$ (or any other group). These processes contain the

same information as M defined in (3.1) with $\mathcal{X}' = \mathbb{R}^n$ and

$h(x, y) = (h_1(x, y), \dots, h_n(x, y))$. More precisely, there is a one-to-one

correspondence between point processes M on $\mathbb{R} \times \mathbb{R}^n$ as in (3.1) and

n -dimensional compound point processes $(M_1, \dots, M_n)$ as in (3.2). Note

that M on $\mathbb{R} \times \mathbb{R}^n$ is $\mathcal{M}(a, F)$ if and only if its corresponding $(M_1, \dots, M_n)$

is an n -dimensional compound Poisson process with rate a and atom

distribution F on $\mathbb{R}^n$. In this case,

$$\begin{aligned} P\{M_1(s, t] \in B_1, \dots, M_n(s, t] \in B_n\} \\ = \sum_{k=0}^{\infty} F^{k*}(B_1 \times \dots \times B_n) a^k (t-s)^k e^{-a(t-s)} / k!. \end{aligned}$$

And each M_j is a compound Poisson process with rate $a_j = a(1 - F_j(0))$ and

atom distribution F_j , where $F_1, \dots, F_n$ are the marginal distributions of

F . Here are some special cases:

C1: $M_1, \dots, M_n$ are independent compound Poisson processes with rates $a_1, \dots, a_n$ and atom distributions $F_1, \dots, F_n$ if and only if $F = F_1 \times \dots \times F_n$.

C2: $(M_1, \dots, M_n)$ is an n -dimensional Poisson process with rate a and point allocation distribution F on $\{0, 1\}^n$ if and only if F has support on $\{0, 1\}^n$. In this case, each M_j is $\mathcal{P}(a_j)$.

C3: $M_1, \dots, M_n$ are independent Poisson processes with rates $a_1, \dots, a_n$ if and only if F has support on $\{e_1, \dots, e_n\}$, where e_j is the j th unit vector with 1 in entry j and 0's elsewhere.

Remark 3.4. Each of the preceding theorems hold for these cases C1, C2,

C3 when the F in the theorem is as specified in the case of interest.

For instance, Theorem 3.2(ii) for case C3 reads: If $a^*(x, \{e_j\}) = a_j$,

$j=1, \dots, m$, and $\alpha^*(x, \{z\}) = 0$ otherwise, $x \in \mathfrak{A}$, then $(M_1, \dots, M_n) \perp\!\!\!\perp X_+$ and $M_1, \dots, M_n$ are independent Poisson processes with respective rates $a_1, \dots, a_n$.

An obvious application of Theorem 3.2(i) shows that the process of customer arrivals in Example 2.5, when $\lambda_n = \lambda$, is a compound Poisson process with rate λ and geometric atom distribution $p^{m-1}(1-p)$. The following is an example of a not so obvious compound Poisson flow in a queueing process.

Example 3.5. A Batch-Service Queueing System With Poisson Arrivals and Compound Poisson Departures. Suppose the Markov process has the state space Z_+ and transition rates

$$\begin{aligned} q(0,1) &= \lambda(1-p), & q(n,n+1) &= \lambda, & n &\geq 1, \\ q(n,n-m) &= \mu p^{m-1}(1-p) & 1 \leq m < n, & & n &\geq 1, \\ q(n,0) &= \mu p^{n-1} & & & n &\geq 1, \end{aligned}$$

where λ, μ are positive and $0 < p < 1$. This process represents the number of customers in a queueing system in which customers arrive at the rate λ and are served in batches as follows. When there are customers in the system, "buses" arrive at a rate μ to take them immediately from the queue. Busing is a common practice in computer systems and material handling systems. The number of customers each bus can take is a random variable with the geometric distribution $p^{m-1}(1-p)$, $m \geq 1$. Also, when there are no customers in the queue and a customer arrives, then with probability p there is a bus available to take the customer without delay. The equilibrium distribution of X is

$$\pi(n) = \pi(0)(1-p)\lambda^n/(\mu + p\lambda)^n, \quad n \geq 1,$$

provided $\lambda < \mu + p\lambda$, which we assume is true (see Kook (1988)).

Consider the compound point process

$$D(\Lambda) = \sum_{t \in \Lambda} \max\{0, X_t - X_{t-}\}$$

that describes the total number of departures in the time period Λ . It records the times at which batches of customers depart and the batch sizes as well. This process corresponds to M in (3.1) with $f(n, n') = 1$ iff $n' \neq n$ and

$$h(n, n') = \begin{cases} n - n' & n' \leq n \\ 0 & n' > n \end{cases}.$$

Clearly, for each $n \geq 0$ and $m \geq 1$,

$$\begin{aligned} \alpha^*(n, \{m\}) &= \pi(n)^{-1} \sum_{n'} \pi(n') q(n', n) f(n', n) 1(h(n', n) = m) \\ &= \pi(n)^{-1} \pi(n + m) q(n + m, n) \\ &= \lambda(1 - p) F(\{m\}), \end{aligned}$$

where $F(\{m\}) = r^{m-1}(1-r)$, $m \geq 1$, and $r = p\lambda/(\mu + p\lambda)$. Thus, if X is stationary, then Theorem 3.2(ii) implies that M is $\mathcal{M}^0(\lambda(1-p), F)$. Hence D is a compound process with rate $\lambda(1-p)$ and geometric atom distribution F .

4. Poisson Flows in a Network of Queues

We shall consider a queueing network process defined as follows. Suppose that $X(t) = (X_1(t), \dots, X_m(t))$, $t \in \mathbb{R}$, is a queueing network process on m nodes, where $X_j(t)$ denotes the number of units (i.e. customers) at node j at time t . The process X takes on values $n = (n_1, \dots, n_m)$ in $\mathbb{Z}_+^m$. Let e_j denote the j th unit vector in $\mathbb{Z}_+^m$ with 1 in entry j and 0's elsewhere. As in Whittle (1986), we assume that units move among the m nodes such that X is a Markov process with transition rates

$$(4.1) \quad q(n, n + e_k) = \lambda_{0k}$$

$$q(n, n - e_j + e_k) = \lambda_{jk} \phi(n - e_j) / \phi(n), \quad n_j \geq 1,$$

$$q(n, n - e_j) = \lambda_{j0} \phi(n - e_j) / \phi(n), \quad n_j \geq 1,$$

and $q(n, n') = 0$ for all other states n' . Here $\phi: \mathbb{Z}_+^m \rightarrow (0, \infty)$, the λ 's are nonnegative, and the subscript 0 denotes the "outside" node. Under this assumption, units enter the nodes $1, \dots, m$ by independent Poisson processes with respective rates $\lambda_{01}, \dots, \lambda_{0m}$. When X is in state n , then $\sum_k \lambda_{jk} \phi(n - e_j) / \phi(n)$ is the departure rate of units from node j . The λ_{jk} is the "arc-dependent" routing intensity from node j to node k , and $\phi(n - e_j) / \phi(n)$ is the "system-dependent" departure intensity from node j (the ratio representing the potential difference between the system in state n and in state $n - e_j$ with one less customer at node j).

We shall assume that X is irreducible. This is equivalent to the irreducibility of the Markov routing matrix

$$(4.2) \quad p(j, k) = \lambda_{jk} / \sum_{\ell=0}^m \lambda_{j\ell}, \quad j, k = 0, \dots, m,$$

where $\lambda_{00} = 0$. The irreducibility of this matrix is equivalent to the existence of unique positive numbers $w_1, \dots, w_m$ that satisfy

$$(4.3) \quad \sum_{k=0}^m (w_j \lambda_{jk} - w_k \lambda_{kj}) = 0, \quad j = 0, \dots, m,$$

where $w_0 = 1$. We also assume that $\sum_n \phi(n) \prod_{j=1}^m w_j^{n_j}$ is finite. Then X is positive recurrent and has the equilibrium distribution (p.198 of Whittle (1986))

$$(4.4) \quad \pi(n) = c \phi(n) \prod_{j=1}^m w_j^{n_j}, \quad n \in \mathbb{Z}_+^m,$$

where c is the normalizing constant. Hereafter, we assume that X is stationary.

Now, consider the point process

$$N_{j0}(A) = \sum_{t \in A} 1(X(t) = X(t-) - e_j)$$

that represents the times at which units exit the network from node j .

Of course, $N_{j0} = 0$ when $\lambda_{j0} = 0$.

Theorem 4.1. The exit processes $N_{10}, \dots, N_{m0}$ are independent Poisson processes with rates $w_1 \lambda_{10}, \dots, w_m \lambda_{m0}$, and $(N_{10}, \dots, N_{m0}) \sim \Pi X_+$.

Whittle (1986) on p. 207 proved this by establishing that the reverse time process X^* of X is again a queueing network process and the exit processes of X are just the time reversals of the Poisson input processes of X^* . Theorem 4.1 also follows from Theorem 3.2(ii) and Remark 3.4 since, for each n and j ,

$$\alpha^*(n, \{e_j\}) = \pi(n)^{-1} \pi(n + e_j) q(n + e_j, n) = w_j \lambda_{j0}$$

and $\alpha^*(n, \{z\}) = 0$ elsewhere.

Along with these exit processes, consider the point processes

$$N_{jk}(A) = \sum_{t \in A} 1(X(t) = X(t-) - e_j + e_k)$$

of times at which units move from node j to node k . We shall now identify sets $J \subset \{1, \dots, m\}$ and $K \subset \{0, 1, \dots, m\}$ such that N_{jk} , $j \in J$, $k \in K$ are independent Poisson processes. Suppose that J and K satisfy the following assumptions:

A1: Each unit that exits J can never return to J . (To verify this one need only check the possible routing under λ_{jk} .)

A2: The system-dependent departure intensity for each node $j \in J$ is of the form

$$\phi(n - e_j) / \phi(n) = \phi_J(n_J - e_j) / \phi_J(n_J)$$

where $n_J = (n_j : j \in J)$, and ϕ_J is a positive function on such vectors.

A3: K is the largest subset of $\{0, 1, \dots, m\}$ such that each unit in K cannot enter J on a subsequent move. (Note that $0 \in K$ and

$$J \cap K = \{j \in J: \lambda_{j\ell} = 0, \ell \in J\}.$$

For some networks, $J = \{1, \dots, m\}$ may be the only set of nodes that satisfies A1. At the other extreme are networks in which each node is visited at most once by a unit, and so each subset of nodes satisfies A1.

Assumption A2 is equivalent to being able to factor Φ as

$\Phi(n) = \Phi_J(n_J)\Psi(n_K: k \in J)$. Upon selecting J conforming to A1, A2, it is advantageous to select K as large as possible as we did in A3.

Let $X_J(t) = \{X_j(t): j \in J\}$, $t \in \mathbb{R}$, denote the process X on the nodes J and let $\mathcal{A}_J$ denote its state space.

Theorem 4.2. The processes N_{jk} , $j \in J$, $k \in K$, are independent Poisson processes with respective rates $w_j \lambda_{jk}$, $j \in J$, $k \in K$. Furthermore,

$$\{N_{jk}: j \in J, k \in K\}_- \stackrel{d}{=} (X_J)_+.$$

Proof. Under the assumptions, the X_J is a queueing network process on the nodes J with transition rates $q_j(n, n')$ defined as in (4.1) with the last line replaced by

$$q_j(n, n - e_j) = \lambda_j \phi_j(n - e_j) / \phi_j(n), \quad n_j \geq 1, n \in \mathcal{A}_J,$$

where $\lambda_j = \sum_{k \in K} \lambda_{jk}$. Furthermore, X_J has the equilibrium distribution

$$\pi_J(n) = c_J \phi_J(n) \prod_{j \in J} \alpha_j^{n_j}, \quad n \in \mathcal{A}_J.$$

This is just the sum of $\pi(n)$ in (4.4) over all n_ℓ , $\ell \in J$. Thus, from

Theorem 4.1, we know that X_J 's exit processes $N_j = \sum_{k \in K} N_{jk}$, $j \in J$, are

independent Poisson processes with respective rates $w_j \lambda_j$, $j \in J$, and that

$$(N_j, j \in J)_- \stackrel{d}{=} (X_J)_+.$$

Next, observe that for each $j \in J$, the processes N_{jk} , $k \in K$, form a partition of N_j in which each point of N_j is assigned to the subprocess N_{jk} with probability λ_{jk}/Λ_j , independent of everything else. Consequently, N_{jk} , $k \in K$, are independent Poisson processes with respective rates $(\lambda_{jk}/\Lambda_j) (w_j \Lambda_j) = w_j \lambda_{jk}$, $k \in K$ (see for instance p.89 of Çinlar (1975)). This property and the preceding argument yield the assertions. $\square$

Example 4.3. Suppose the network has a node j such that each unit passing through the network visits j at most once. Also, assume that $\phi(n) = \phi_j(n_j) \psi(n_k: k \neq j)$, and let $J = \{j\}$. Then the departure processes $N_{j0}, N_{j1}, \dots, N_{jm}$ are independent Poisson processes as in Theorem 4.2. Now, suppose that each node of the network can be visited at most once by each unit and $\phi(n) = \phi_1(n_1) \dots \phi_m(n_m)$. Then each process N_{jk} is $\mathcal{P}(w_j \lambda_{jk})$; some of these processes may be dependent. Consider the arrival processes to a fixed node k . Let J denote the set of all nodes j that can never be reached from k . Then, under the preceding supposition, the arrival processes N_{jk} , $j \in J$, are independent Poisson processes as in Theorem 4.2.

Example 4.4. Multivariate Poisson Flows. Suppose $J_1, \dots, J_p$ are subsets of $\{1, \dots, m\}$ and $K_1, \dots, K_p$ are subsets of $\{0, 1, \dots, m\}$ (the subsets need not be disjoint). Consider the process $N_i = \sum_{\substack{j \in J_i \\ k \in K_i}} N_{jk}$ of times at which

units move from some node in J_i to some node in K_i . Suppose the sets $J = J_1 \cup \dots \cup J_p$ and $K = K_1 \cup \dots \cup K_p$ satisfy assumptions A1, A2, A3, or that they are contained in a pair of such sets. Then N_{jk} , $j \in J$, $k \in K$, are independent Poisson processes as in Theorem 4.2. Consequently,

$(N_1, \dots, N_p)$ is a p -dimensional Poisson process with rate $\alpha = \sum_{\substack{j \in J \\ k \in K}} w_j \lambda_{jk}$ and

point allocation distribution

$$F(\{\sum_{i \in I} e_i\}) = a^{-1} \sum_{j,k}^I w_{jk} \lambda_{jk}$$

where $I \subset \{1, \dots, v\}$, and the sum is over all j in $\bigcap_{i \in I} J_i$, and k in $\bigcap_{i \in I} K_i$.

The $N_1, \dots, N_v$ are independent if $\bigcap_i J_i = \emptyset$ and $\bigcap_i K_i = \emptyset$.

Example 4.5. Networks With Several Types of Units. Consider a network as above in which each unit carries a label from a finite set $\mathcal{A}$ of types and the label may change when the unit moves. We represent this network by the process $X(t) = \{X_{aj}(t) : a \in \mathcal{A}, j=1, \dots, m\}$, $t \in \mathbb{R}$, where $X_{aj}(t)$ is the number of type a units at node j at time t . A typical state of X is $n = (n_{aj} : a \in \mathcal{A}, j=1, \dots, m)$. We assume that X is a Markov process with transition rates (analogous to (4.1))

$$\begin{aligned} q(n, n + e_{bk}) &= \lambda_{0,bk} \\ q(n, n - e_{aj} + e_{bk}) &= \lambda_{aj,bk} \phi(n - e_{aj}) / \phi(n), \quad n_{aj} \geq 1, \\ q(n, n - e_{aj}) &= \lambda_{aj,0} \phi(n - e_{aj}) / \phi(n), \quad n_{aj} \geq 1. \end{aligned}$$

Under assumptions as above, the process X has the equilibrium distribution

$$\pi(n) = c \phi(n) \prod_{aj} w_{aj}^{n_{aj}}.$$

The results above hold for this network - one need only use double indices aj, bk in place of j, k and consider the point processes $N_{aj,bk}$, $aj \in J$, $bk \in K$, where J, K satisfy assumptions A1, A2, A3 with double indices. For instance, suppose $\Lambda \subset \mathcal{A}$ is such that a unit with a label from $\mathcal{A} \setminus \Lambda$ can never carry a label from Λ . Consider the point processes of times at which units change labels from Λ to $\mathcal{A} \setminus \Lambda$. These processes are $N_{aj,bk}$, $aj \in J$, $bk \in K$, where $J = \{aj : a \in \Lambda, j=1, \dots, m\}$ and $K = \{bk : b \in \mathcal{A} \setminus \Lambda, k=0, \dots, m\}$. Clearly J satisfies A1 and K satisfies A3.

Suppose J also satisfies A2. Then the preceding point processes are independent Poisson processes with respective rates $\{w_{aj}\lambda_{aj,bk}\}$, and the past of these processes is independent of $(X_j)_+$.

The preceding results are for networks in which only one unit can move at a time. In networks with simultaneous movement of units, the flows among the nodes may be compound Poisson processes. Kook (1988) has characterized such flows using Theorem 3.2.

5. Further Generalizations

The results in Sections 2 and 3 readily extend to more general processes. We discussed some of these situations in the remarks above. Here are some more generalizations.

Markov Processes With General State Spaces. Suppose that $X = \{X_t : t \in \mathbb{R}\}$ is a pure jump Markov process with a general state space E and associated σ -field $\mathcal{E}$, and its transition kernel is $K(x, B)$, $x \in E$, $B \in \mathcal{E}$. That is, the exponential sojourn time in state x has parameter $K(x, E)$ and the probability of X jumping from x into B is $K(x, B)/K(x, E)$. Then the results in Sections 2 and 3 hold with the sums replaced by integrals. For example, α^* is the Radon-Nikodym derivative

$$\alpha^*(x) = \int_E \pi(dy) K(y, dx) f(y, x) / \pi(dx).$$

Functionals Involving Sojourn Times. Suppose X is a Markov process as in Section 2 and

$$N(\Lambda) = \sum_{t \in \Lambda} f(X_t, X_{t-}, W_t)$$

where W_t is the waiting time in state X_{t-} and $f: \mathcal{E} \times \mathcal{E} \times \mathbb{R}_+ \rightarrow \{0, 1\}$. Then the results of Section 2 hold for this N with α, α^* defined by

$$\alpha(x) = \sum_y \int_0^\infty q(x,y) f(x,y,w) F_x(dw)$$

$$\alpha^*(x) = \pi(x)^{-1} \int_0^\infty \sum_y \pi(y) q(y,x) f(y,x,w) F_y(dw)$$

where $F_x(w) = 1 - \exp(-wq(x))$. Similarly, the results of Section 3 hold for M as a functional of W_t as well as of X_t, X_{t-} . One can generalize further by assuming that X is a semi-Markov process and replacing F_x by a general sojourn time distribution $F_{x,y}$.

References

- Benveniste, A. and J. Jacod (1973). Systemes de Lévy des processus de Markov. Invent. Math. 21, 183-98.
- Brémaud, P. (1981). Point Processes and Queues: Martingale Dynamics. Springer-Verlag, New York.
- Burke, P. J. (1956). The output of a queueing system. Operations Res. 4, 699-704.
- Çınlar, E. (1975). Introduction to Stochastic Processes. Prentice-Hall, Englewood Cliffs.
- Disney, R. L. and P. C. Kiessler (1987). Traffic Processes in Queueing Networks: A Markov Renewal Approach. Johns Hopkins University Press, Baltimore.
- Disney, R. L. and D. König (1985). Queueing networks: a survey of their random processes. SIAM Review, 27, 335-403.
- Gross, D. and C. M. Harris (1985). Fundamentals of Queueing Theory. Wiley, New York.
- Kallenberg, O. (1983). Random Measures. 3rd Ed. Akademie-Verlag, Berlin and Academic Press, New York.
- Karr, A. F. (1986). Point Processes and Their Statistical Inference. Dekker, New York.
- Kelly, F. P. (1979). Reversibility and Stochastic Networks. Wiley, New York.
- Kook, K. (1988). Ph.D. Dissertation. Industrial and Systems Engineering, Georgia Institute of Technology.

Liptser, R. Sh. and A. N. Shiriyayev (1978). Statistics of Random Processes. Vol II. Springer-Verlag, New York.

Melamed, B. (1979). On Poisson traffic processes in discrete-state Markovian systems with applications to queueing theory. Adv. Appl. Prob. 11, 218-239.

Reich, E. (1957). Waiting times when queues are in tandem. Ann. Math. Stat. 28, 768-773.

Varaiya, P. and J. Walrand (1981). Flows in queueing networks: a martingale approach. Math. Operations Res., 6, 387-404.

Whittle, P. (1986). Systems in Stochastic Equilibrium. Wiley, New York.

178. G. Kallianpur, A.G. Miamee and H. Niemi, On the prediction theory of two parameter stationary random fields, March 87.
179. R. Brigola, Remark on the multiple Wiener integral, Mar. 87.
180. R. Brigola, Stochastic filtering solutions for ill-posed linear problems and their extension to measurable transformations, Mar. 87.
181. G. Samorodnitsky, Maxima of symmetric stable processes, Mar. 87.
182. H.L. Hurd, Representation of harmonizable periodically correlated processes and their covariance, Apr. 87.
183. H.L. Hurd, Nonparametric time series analysis for periodically correlated processes, Apr. 87.
184. T. Mori and H. Oodaira, Freidlin-Wentzell estimates and the law of the iterated logarithm for a class of stochastic processes related to symmetric statistics, May 87.
185. R.F. Serfozo, Point processes, May 87. Operations Research Handbook on Stochastic Processes, to appear.
186. Z.D. Bai, W.Q. Liang and W. Vervaat, Strong representation of weak convergence, June 87.
187. O. Kallenberg, Decoupling identities and predictable transformations in exchangeability, June. 87.
188. O. Kallenberg, An elementary approach to the Danielli-Kolmogorov theorem and some related results, June 87. Math. Nachr., to appear.
189. G. Samorodnitsky, Extrema of skewed stable processes, June 87.
190. D. Nualart, M. Sanz and M. Zakai, On the relations between increasing functions associated with two-parameter continuous martingales, June 87.
191. F. Avram and M. Taqqu, Weak convergence of sums of moving averages in the α -stable domain of attraction, June 87.
192. M.R. Leadbetter, Harald Cramér (1893-1985), July 87. ISI Review, to appear.
193. R. LePage, Predicting transforms of stable noise, July 87.
194. R. LePage and B.M. Schreiber, Strategies based on maximizing expected log, July 87.
195. J. Rosinski, Series representations of infinitely divisible random vectors and a generalized shot noise in Banach spaces, July 87.
196. J. Szulga, On hypercontractivity of α -stable random variables, $0 < \alpha < 2$, July 87.
197. I. Kuznezova-Sholpo and S.T. Rachev, Explicit solutions of moment problems I, July 87.
198. T. Hsing, On the extreme order statistics for a stationary sequence, July 87.
199. T. Hsing, On the characterization of certain point processes, Aug. 87.
200. J.P. Nolan, Continuity of symmetric stable processes, Aug. 87.
201. M. Marques and S. Cambanis, Admissible and singular translates of stable processes, Aug. 87.
202. O. Kallenberg, One-dimensional uniqueness and convergence results for exchangeable processes, Aug. 87.
203. R.J. Adler, S. Cambanis and G. Samorodnitsky, On stable Markov processes, Sept. 87.
204. G. Kallianpur and V. Perez-Abreu, Stochastic evolution equations driven by nuclear space valued martingales, Sept. 87.
205. R.L. Smith, Approximations in extreme value theory, Sept. 87.
206. E. Willekens, Estimation of convolution tails, Sept. 87.
207. J. Rosinski, On path properties of certain infinitely divisible processes, Sept. 87.
208. A.H. Korezlioglu, Computation of filters by sampling and quantization, Sept. 87.
209. J. Bather, Stopping rules and observed significance levels, Sept. 87.
210. S.T. Rachev and J.E. Yukich, Convolution metrics and rates of convergence in the central limit theorem, Sept. 87.
211. T. Fujisaki, Normed Bellman equation with degenerate diffusion coefficients and its applications to differential equations, Oct. 87.
212. G. Simons, Y.C. Yao and X. Wu, Sequential tests for the drift of a Wiener process with a smooth prior, and the heat equation, Oct. 87.
213. R.L. Smith, Extreme value theory for dependent sequences via the Stein-Chen method of Poisson approximation, Oct. 87.
214. C. Houdré, A note on vector bimeasures, Nov. 87.
215. M.R. Leadbetter, On the exceedance random measures for stationary processes, Nov. 87.
216. M. Marques, A study on Lebesgue decomposition of measures induced by stable processes, Nov. 87.
217. M.T. Alpuim, High level exceedances in stationary sequences with extremal index, Dec. 87.
218. R.F. Serfozo, Poisson functionals of Markov processes and queueing networks, Dec. 87.
219. J. Bather, Stopping rules and ordered families of distributions, Dec. 87.

END

DATE

FILMED

8-88

DTIC