

NOSC

NAVAL OCEAN SYSTEMS CENTER San Diego, California 92152-5000

Technical Document 1220
January 1988

Laser Dye Synthesis — Group C Compounds

E. J. Grubbs

San Diego State University Foundation
San Diego State University



Approved for public release;
distribution is unlimited.

The views and conclusions contained in this report are those of the authors and should not be interpreted as representing the official policies, either expressed or implied, of the Naval Ocean Systems Center or the U.S. Government.

NAVAL OCEAN SYSTEMS CENTER

San Diego, California 92152-5000

E. G. SCHWEIZER, CAPT. USN
Commander

R. M. HILLYER
Technical Director

ADMINISTRATIVE INFORMATION

This task was performed for the U.S. Army, Army Missile Command, Redstone Arsenal, AL 35398. Work under contract N66001-85-D-0203 was done through the San Diego State University Foundation, San Diego, CA 92182. The Contracting Officer's Technical Representative was T.G. Pavlopoulos of the Chemistry/Biochemistry Branch, Naval Ocean Systems Center, San Diego, CA 92152-5000.

Released by
E. Lindner, Head
Chemistry/Biochemistry
Branch

Under authority of
S. Yamamoto, Head
Environmental Sciences
Division

UNCLASSIFIED


SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			Approved for public release; distribution is unlimited.		
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S) NOSC TD 1220		
6a. NAME OF PERFORMING ORGANIZATION San Diego State University Foundation San Diego State University		6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION Naval Ocean Systems Center		
6c. ADDRESS (City, State and ZIP Code) San Diego, CA 92182			7b. ADDRESS (City, State and ZIP Code) San Diego, CA 92152-5000		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION U.S. Army Army Missile Command		8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N66001-85-D-0203		
8c. ADDRESS (City, State and ZIP Code) Redstone Arsenal, AL 35398			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO. RDA	PROJECT NO. ARMY	TASK NO. ARMY
			AGENCY ACCESSION NO DN305 157		
11. TITLE (Include Security Classification) Laser Dye Synthesis — Group C Compounds					
12. PERSONAL AUTHOR(S) E.J. Grubbs					
13a. TYPE OF REPORT Final		13b. TIME COVERED FROM TO		14. DATE OF REPORT (Year, Month, Day) January 1988	
15. PAGE COUNT					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
			Laser dyes		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) This report describes progress on the syntheses of a number of new potential laser dyes. The goal is to generate new structures allowing the upscaling of flashlamp pumped dye lasers to high energies. These systems ideally should show a lower action threshold, operate with high efficiencies, and exhibit greater photochemical stabilities than do those presently available.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT ☐ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT ☐ DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL T.G. Pavlopoulos			22b. TELEPHONE (include Area Code) (619) 553-2792		22c. OFFICE SYMBOL Code 521

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)



DD FORM 1473, 84 JAN

UNCLASSIFIED


SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

This report describes progress on the syntheses of a number of new potential laser dyes. The goal is to generate new structures allowing the upscaling of flashlamp pumped dye lasers to high energies. These systems ideally should show a lower laser action threshold, operate with high efficiencies, and exhibit greater photochemical stabilities than do those presently available.

Pavlopoulos has proposed (a) that the parent aromatic chromophores be substituted by auxochromic groups such as hydroxy, alkoxy, and dialkylamino and (b) that improvement should result by restricting the chromophore and auxochromic group to coplanarity. This should reduce intersystem crossing and thus improve the quantum fluorescence yield. The potential dye lasers described below are designed to meet these objectives.


1. The Synthesis of 1,2-bis-(9,10-dihydro-2-phenanthryl) ethene (3).

This extended stilbene was prepared by converting dihydrophenanthrene to the aldehyde using dichloromethyl methyl ether and stannic chloride. The general method has been described by Rieche, Gross, and Höft.¹




The crude aldehyde was obtained in 85% yield but appears to be a mixture of the 2-carboxaldehyde (shown) and the 3-carboxaldehyde in a ratio of about 3:1. The proton nmr spectrum of this mixture shows the expected aromatic absorptions at $\delta 7.1$ - 7.4 (7-proton multiplet), a four proton multiplet center at $\delta 2.86$ and two aldehyde singlets at $\delta 9.86$ and $\delta 10.3$. The mass spectrum was consistent with the proposed isomeric mixture showing the molecular ion at m/e 208 and fragments corresponding to losses of the aldehydic hydrogen, the aldehyde group ($\text{M}^+ - \text{CHO}$, parent ion), and the CH_2CH_2 bridge.

The aldehyde mixture was subjected to the McMurry coupling procedure² as modified by Geise and coworkers.³




A variety of reagent ratios and reaction times were explored. Relatively low yields (approximately 10-20%) of a highly fluorescent product was obtained as a white solid. After chromatography over florisil and recrystallization from hexane the purified samples still melted over an eleven degree range (239.5-251°C). The proton nmr spectrum and mass spectrum are consistent with the extended stilbene (3) but do not provide quantitative information regarding the distribution of positional and geometric isomers in the mixture. Although future coupling studies employing purified samples of the 9,10-dihydro-2-phenanthrene carboxaldehyde are planned (pending results from laser spectroscopic studies on the delivered samples of 3 mixtures), the McMurry coupling is complicated by formation of substantial quantities of the corresponding diol 4.




II. Attempted Preparation of 1,2-bis-(2-fluorenyl) ethene (7).

A. Thermal Decomposition of Azine

Saint-Ruf and Buu-Hoi⁴ have reported the preparation of the bis fluorenyl ethene 7 by thermolysis of the corresponding azine in paraffin oil. Accordingly the azine was prepared in 95% yield by the usual method. However, in our hands several attempts to pyrolyze the azine either in paraffin oils or




in silicon oil led only to small yields of complex mixtures.





B. McMurry coupling of 2-fluorene carboxaldehyde.

Attempts to obtain **7** using the McMurry coupling procedure^{2,3} again led to complex mixtures from which 2-methylfluorene was separated as the only identifiable product.




C. Condensation-dehydrobromination of 2-bromomethylfluorene.

Pinck and Hilbert⁵ have described the preparation of 1,2-bis(1-fluorenyl) ethene by the treatment of 1-bromomethylfluorene with sodium methoxide in methanol. In hopes of applying this procedure to the 2-bromomethyl isomer, the following transformations were performed. The borohydride reduction of **5**




afforded purified 8 in 75% yield. Conversion of the alcohol to the bromide was accomplished using the general procedure of Braun and Engel.⁶ However, unlike the behavior of 1-bromomethylfluorene, the treatment of 9 with sodium methoxide afforded the corresponding methyl ether 10 (73%) as the only isolable product. Additional attempts using the condensation-dehydrobromination



route employing potassium *t*-butoxide or 1,8-diazabicyclo [5.4.0] undec-7-ene (DBU) as the base were unsuccessful.


III. Derivatives of 9,9-dimethylfluorene.

To remove complications arising from potential facile removal of protons (or H atoms) from C₉ during the course of various desired transformations, fluorene was converted to its 9,9-dimethyl derivative using the procedure described by Bavin.⁷




A. 9,9-Dimethyl-2-fluorene carboxyaldehyde (13).

The aldehyde was prepared in yields as high as 89% using the method of Rieche, Gross and Hoft.¹ The spectroscopic properties for 13 were in accord




with this proposed structure. Because of the uncertainty regarding the position of the aldehyde group in 13, the following sequence of transformations was undertaken. The cyclic acetal 14 obtained directly from 13 was shown to be



identical to that obtained starting with an authentic sample of fluorene 2-carboxaldehyde (5).


B. 1,2-bis-(9,9-dimethylfluoren-2-yl) ethene (16).

The aldehyde 13 was subjected to the McMurry coupling conditions (TiCl_3 , LiAlH_4). By this procedure the desired alkene was obtained in yields of approximately 10%. The dimethylated bis fluorenylethene



was obtained as a light-yellow highly fluorescent solid.

Samples of 3 and 16 as well as previously prepared samples of



have been delivered for testing.

References

1. A. Rieche, H. Gross and E. Höft, Ber., 1964, 97, 88.
2. J.E. McMurry, Acc. Chem. Res., 1974, 7, 281.
3. R. Dams, M. Malinowski, I. Westdorp and H.Y. Geise, J. Org. Chem., 1982, 47, 248.
4. G. S-Ruf and N.P. B-Hoi, Bull. Soc. Chim. Fr., 1970, 525.
5. L.A. Pinck and G.E. Hilbert, J. Am. Chem. Soc., 1946, 68, 751.
6. J. v.Braun and H. Engel, Ber., 1924, 57, 191.
7. P.M.G. Bavin, Can. J. Chem., 1960, 38, 882.