

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

DTIC FILE COPY

(2)

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

AD-A198 435

DTIC

CTE

0183 1988

D

1b. RESTRICTIVE MARKINGS
N/A3. DISTRIBUTION/AVAILABILITY OF REPORT
Approved for public release:
Distribution unlimited

4. PERFORMING ORGANIZATION REPORT NUMBER(S)

TR# 11

5. MONITORING ORGANIZATION REPORT NUMBER(S)

6a. NAME OF PERFORMING ORGANIZATION

The University of Chicago

6b. OFFICE SYMBOL
(If applicable)

7a. NAME OF MONITORING ORGANIZATION

Office of Naval Research

6c. ADDRESS (City, State, and ZIP Code)

5801 S. Ellis Avenue
Chicago, IL 60637

7b. ADDRESS (City, State, and ZIP Code)

800 N. Quincy Street
Arlington, VA 222178a. NAME OF FUNDING/SPONSORING
ORGANIZATION Office of Naval
Research Physics Program8b. OFFICE SYMBOL
(If applicable)

9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

Contract : N00014-86-K-0254

8c. ADDRESS (City, State, and ZIP Code)

800 N. Quincy Street
Arlington, VA 22217

10. SOURCE OF FUNDING NUMBERS

PROGRAM
ELEMENT NO.

Code 1112

PROJECT
NO.TASK
NO.

4121023

WORK UNIT
ACCESSION NO.

11. TITLE (Include Security Classification)

Molecular Beam Studies of Surface Phonons and Gas Surface Interactions

12. PERSONAL AUTHOR(S)

S.J. Sibener

13a. TYPE OF REPORT

Interim Technical

13b. TIME COVERED

FROM _____ TO _____

14. DATE OF REPORT (Year, Month, Day)

1988 June 15

15. PAGE COUNT

5

16. SUPPLEMENTARY NOTATION

Invited Progress Report: Proceedings of the Xth International
Symposium on Molecular Beams, Cannes, France

17. COSATI CODES

FIELD	GROUP	SUB-GROUP

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)
Surface phonon spectroscopy; thin film dynamics; phonon dispersion relations; physisorption interactions; gas-surface energy exchange; inelastic single phonon scattering; rare gas interaction potentials; molecular beams. (1112)

19. ABSTRACT (Continue on reverse if necessary and identify by block number) Angle and velocity resolved inelastic helium scattering is becoming an incisive tool for probing the vibrational characteristics of clean and adsorbate covered surfaces. These experiments can be visualized as being a two-dimensional (i.e., surface sensitive) analog to inelastic neutron scattering, which has played such a crucial role in elucidating the nature of elementary excitations in bulk crystals. In this paper a general review is presented of our recent work which demonstrates how the surface vibrational properties of epitaxially grown rare gas films evolve, on a layer-by-layer basis, towards those characteristic of thick crystals. Lattice dynamics calculations are used to quantitatively analyze the data. Keywords:

20. DISTRIBUTION/AVAILABILITY OF ABSTRACT

☒ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT ☐ DTIC USERS

21. ABSTRACT SECURITY CLASSIFICATION

Unclassified

22a. NAME OF RESPONSIBLE INDIVIDUAL

James S. Murday/Richard Brandt

22b. TELEPHONE (Include Area Code)

(202) 696-4206

22c. OFFICE SYMBOL

DD Form 1473, JUN 86

Previous editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

Unclassified

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0254

Task No. 4121023

Technical Report No. 11

Molecular Beam Studies of Surface Phonons
and Gas Surface Interactions

By

S.J. Sibener

Prepared for Publication

in the

Proceedings of the Xth International Symposium
on Molecular Beams

Department of Chemistry and the James Franck Institute
The University of Chicago
Chicago, Illinois 60637

June, 1988

Reproduction in whole or in part is permitted for
any purpose of the United States Government

This document has been approved for public release
and sale; its distribution is unlimited.



Accession For	
NTIS	CRA&I ☒
DTIC	TAB ☐
Unannounced	☐
Justification	
By	
Distribution	
Availability Codes	
For	Approved for Special
A-1	

Molecular Beam Studies of Surface Phonons
and Gas-Surface Interactions

Steven J. Sibener

The James Franck Institute and The Department of Chemistry
The University of Chicago
Chicago, Illinois 60637 U.S.A.

The scattering of atomic and molecular beams from well-characterized single crystal surfaces is providing important structural and dynamical information pertaining to clean and adsorbate covered systems. Prime objectives in these studies include the determination of gas-surface interaction potentials, the development of a microscopic understanding of collision induced gas-surface energy transfer, surface crystallography, the elucidation of surface reaction mechanisms, and high resolution ($\sim 1 \text{ cm}^{-1}$) surface vibrational spectroscopy. At the present time especial attention is being given to determining which surface modes actively participate in collision induced energy exchange, the mapping out of surface phonon dispersion relations, and determining under which energy regimes single or multiphonon transitions dominate the interaction.

It is the intent of this progress report to cover some of our recent activities in this field. In particular, the growing importance that angle and velocity resolved inelastic He scattering is playing as a surface spectroscopic tool will be highlighted. These experiments can be visualized as being a two-dimensional (i.e., surface sensitive) analog to inelastic neutron scattering, which has played such a crucial role in elucidating the nature of elementary excitations in bulk crystals.

Experimental data, lattice dynamics calculations, and quantum scattering calculations will be presented which reveal how the surface vibrational properties and geometric structures of ordered Ar, Kr, and Xe rare gas films

evolve, on a layer-by-layer basis, into those of thick crystals. These films form azimuthally aligned but translationally incommensurate structures on the support crystal, Ag(111). The monolayer SP_1 frequencies give valuable information necessary for constructing accurate rare gas-metal physisorption potentials. Deconvoluted inelastic single phonon scattering transition probabilities will also be discussed. Comparison of the experimental phonon dispersion curves with predictions derived from lattice dynamics calculations has allowed us to examine the validity of using gas phase rare gas pair potentials to model condensed phase systems. Examples of the types of data collected are shown in Figures 1 and 2, which respectfully contain time-of-flight spectra and bilayer phonon dispersion curves.

The inelastic scattering experiments are complemented by He diffraction and selective adsorption measurements, which yield information on the nature of He-(rare gas) surface potentials.

Finally, if time permits, a discussion of molecule-surface interactions, adsorbate island characterization, and preliminary results employing a new laser-based particle detector will be briefly presented.

It is with great pleasure that I acknowledge my collaborators and colleagues who made essential contributions to this progress report: K. D. Gibson and C. F. Yu for experimental measurements; K. B. Whaley, E. Pontius and J. C. Light for quantum scattering calculations; and B. M. Hall, D. Mills, and J. E. Black for the theoretical lattice dynamics results. This work was supported, in part, by the Office of Naval Research and the NSF-Materials Research Laboratory Program at The University of Chicago.




Figure 1. TOF spectra of Kr overlayers physisorbed on Ag(111), $T_s = 25$ K and $E_i = 18$ meV. Crosses are the experimental data, solid lines are least-squares fits to the data, and dashed lines are computer simulations assuming single phonon interactions with delta-function excitation lineshapes. Heights for the simulations were adjusted to be the same as that of the fits. Arrows indicate the position of elastic time-of-flight.




Figure 2. Comparison between the experimental and lattice dynamics results for the bilayer phonon dispersion curves. The crosses are the experimental points, and the slashes through selected crosses are error bars. The solid lines are the best fits to the data. The triangles are the results from lattice dynamics calculations which use as input the experimental lattice constants and realistic two-body Ar-Ar, Kr-Kr, and Xe-Xe pair potentials.