

4

DTIC FILE (A4)

A REVISION OF COUPLED MODE THEORY
FOR IRREGULAR ACOUSTIC WAVEGUIDES

Ronald F. Pannatoni

DR. RONALD F. PANNATONI
120 Mark Dowdle Road
Franklin, North Carolina 28734

8 August 1988

Final Report for Contract
N00014-83-C-0230

Approved for public release.
Distribution is unlimited.

Prepared for
OFFICE OF NAVAL RESEARCH
800 North Quincy Street
Arlington, Virginia 22217-5000

DCASMA ATLANTA
805 Walker Street
Marietta, Georgia 30060-2789

DTIC
UNCLASSIFIED
AUG 16 1988

D

H

Copyright © 1988 by Ronald F. Pannatoni.
This material may be reproduced by or for the U.S. Government.

88 8 17 1 2

AD-A198 466

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS NONE										
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; Distribution Unlimited										
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE												
4. PERFORMING ORGANIZATION REPORT NUMBER(S)		5. MONITORING ORGANIZATION REPORT NUMBER(S)										
6a. NAME OF PERFORMING ORGANIZATION DR. RONALD F. PANNATONI	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION DCASMA ATLANTA										
6c. ADDRESS (City, State and ZIP Code) 120 Mark Dowdle Road Franklin, North Carolina 28734		7b. ADDRESS (City, State and ZIP Code) 805 Walker Street Marietta, Georgia 30060-2789										
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-83-C-0230										
8c. ADDRESS (City, State and ZIP Code) 800 North Quincy Street Arlington, Virginia 22217-5000		10. SOURCE OF FUNDING NOS. <table border="1"> <tr> <td>PROGRAM ELEMENT NO.</td> <td>PROJECT NO.</td> <td>TASK NO.</td> <td>WORK UNIT NO.</td> </tr> </table>		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT NO.					
PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT NO.									
11. TITLE (Include Security Classification) (U) A revision of coupled mode theory for irregular acoustic waveguides												
12. PERSONAL AUTHOR(S) Ronald F. Pannatoni												
13a. TYPE OF REPORT Final	13b. TIME COVERED FROM 1983 TO 1987	14. DATE OF REPORT (Yr., Mo., Day) 1988 August 8	15. PAGE COUNT 6									
16. SUPPLEMENTARY NOTATION												
17. COSATI CODES <table border="1"> <tr> <td>FIELD</td> <td>GROUP</td> <td>SUB GR.</td> </tr> <tr> <td></td> <td></td> <td></td> </tr> <tr> <td></td> <td></td> <td></td> </tr> </table>		FIELD	GROUP	SUB GR.							18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) (U) Underwater Acoustics; (U) Coupled Mode Theory	
FIELD	GROUP	SUB GR.										
19. ABSTRACT (Continue on reverse if necessary and identify by block number) This note concerns new eigenfunction expansions for the fields in an irregular acoustic waveguide. The expansions are associated with a non-selfadjoint eigenvalue problem that has the eigenvalue in some of the boundary conditions. The waveguide consists of two layers of fluid that are separated by a frictionless free interface. The field expansions converge uniformly on both sides of the interface even if the fluid density is discontinuous at the interface. We use the eigenfunction expansions to convert the reduced wave equation and the conditions at the boundaries of the waveguide into a system of first-order, ordinary differential equations. These coupling equations are similar to Shevchenko's equations for coupling between local modes in the waveguide [Sov. Phys. Acoust. 7 (1962) pp. 392-397, eqs. (16-18)].												
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT. ☐ DTIC USERS ☐		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED										
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. Ronald F. Pannatoni		22b. TELEPHONE NUMBER (Include Area Code) (704) 524-7623	22c. OFFICE SYMBOL									

DD FORM 1473, 83 APR

EDITION OF 1 JAN 73 IS OBSOLETE.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

A revision of coupled mode theory for irregular acoustic waveguides

RONALD F. PANNATONI

Franklin, North Carolina, USA

August 8, 1988

1. INTRODUCTION

This note concerns new eigenfunction expansions for the fields in an irregular acoustic waveguide. The expansions are associated with a non-selfadjoint eigenvalue problem that has the eigenvalue in some of the boundary conditions. The waveguide consists of two layers of fluid that are separated by a frictionless free interface. The field expansions converge uniformly on both sides of the interface even if the fluid density is discontinuous at the interface.

We use the eigenfunction expansions to convert the reduced wave equation and the conditions at the boundaries of the waveguide into a system of first-order, ordinary differential equations. These coupling equations are similar to Shevchenko's equations for coupling between local modes in the waveguide [*Sov. Phys. Acoust.* 7(1962) pp. 392-397, eqs. (16-18)].

The eigenvalue problem and the field expansions are described in the next section. The coupling equations are stated in the last section.

2. THE EIGENFUNCTION EXPANSIONS

A. Notation and assumptions. The discussion in this note is limited to two-dimensional problems. The real variables x and z denote, respectively, the coordinates of horizontal and vertical position in the waveguide. The following notation is used for the indicated limits and jump discontinuities of an arbitrary function $F(x, z)$:

$$(1) \quad F(x, z+0) = \lim_{\substack{\epsilon \rightarrow 0 \\ \epsilon > 0}} F(x, z + \epsilon),$$

$$(2) \quad F(x, z-0) = \lim_{\substack{\epsilon \rightarrow 0 \\ \epsilon > 0}} F(x, z - \epsilon),$$

$$(3) \quad \text{jump}[F]_{(x,z)} = F(x, z+0) - F(x, z-0).$$





Fig. 1. Geometry of the waveguide and conditions at the boundaries.

A representative section of the waveguide is sketched in Figure 1. The surface boundary coincides with the horizontal line $z = 0$, and the bottom boundary coincides with the horizontal line $z = -L$. The interface between layers in the waveguide is described by the curve $z = -H(x)$. It is assumed that $0 < H(x) < L$ for all x and that the function $H(x)$ has a continuous second derivative.

The local sound speed $c(x, z)$ is continuous in each layer, and the local mass density $\rho(x, z)$ is continuously differentiable in each layer. These quantities may be discontinuous at the interface, but it is assumed that each of the four limits $c(x, -H(x) \pm 0)$ and $\rho(x, -H(x) \pm 0)$ exists and is positive.

Sound in the waveguide is assumed to depend harmonically on time with angular frequency ω . The acoustic pressure at time t can be expressed as the real part of the product $p(x, z) \exp(-i\omega t)$, where the complex pressure $p(x, z)$ solves a reduced wave equation in two dimensions: if $-L < z < -H(x)$ or $-H(x) < z < 0$ then

$$(4) \quad \rho(x, z) \frac{\partial}{\partial x} \left(\frac{1}{\rho(x, z)} \frac{\partial}{\partial x} p(x, z) \right) + \rho(x, z) \frac{\partial}{\partial z} \left(\frac{1}{\rho(x, z)} \frac{\partial}{\partial z} p(x, z) \right) + \frac{\omega^2}{c(x, z)^2} p(x, z) = S(x, z).$$



The forcing term $S(x, z)$ in eq.(4) represents distributed sources of sound inside the waveguide.

The acoustic pressure vanishes at the external boundaries of the waveguide, and it is continuous at the interface:

$$(5) \quad p(x, -L) = 0,$$

$$(6) \quad p(x, 0) = 0,$$

$$(7) \quad \text{jump}_{(x, -H(x))} [p] = 0.$$

The fluid velocity associated with sound in the waveguide is proportional to the acoustic pressure gradient. The components of velocity in the horizontal and vertical directions can be expressed, respectively, as the real parts of the products $u(x, z) \exp(-i\omega t)$ and $w(x, z) \exp(-i\omega t)$, where the complex velocity components $u(x, z)$ and $w(x, z)$ are

$$(8) \quad u(x, z) = \frac{1}{i\omega \rho(x, z)} \frac{\partial}{\partial x} p(x, z),$$

$$(9) \quad w(x, z) = \frac{1}{i\omega \rho(x, z)} \frac{\partial}{\partial z} p(x, z).$$

At the interface the component of fluid velocity normal to the interface is continuous. This requirement can be expressed in terms of the complex velocity components as

$$(10) \quad \text{jump}_{(x, -H(x))} [w + \dot{H}(x) u] = 0,$$

where $\dot{H}(x) = (d/dx)H(x)$.

B. The eigenvalue problem. At each position x the following equations determine countably infinite sets of eigenvalues $k_n(x)$ and eigenfunctions $\phi_n(x, z)$:

$$(11) \quad \rho(x, z) \frac{\partial}{\partial z} \left(\frac{1}{\rho(x, z)} \frac{\partial}{\partial z} \phi_n(x, z) \right) + \left(\frac{\omega^2}{c(x, z)^2} - k_n(x)^2 \right) \phi_n(x, z) = 0$$

if $-L < z < -H(x)$ or $-H(x) < z < 0$,

$$(12) \quad \phi_n(x, -L) = 0,$$

$$(13) \quad \phi_n(x, 0) = 0,$$

$$(14) \quad \text{jump}_{(x, -H(x))} \left[\dot{H}(x) \frac{\partial}{\partial z} \phi_n - i k_n(x) \phi_n \right] = 0,$$

$$(15) \quad \text{jump}_{(x, -H(x))} \left[\frac{1}{\rho} \left(\frac{\partial}{\partial z} \phi_n + \dot{H}(x) i k_n(x) \phi_n \right) \right] = 0.$$

The eigenfunctions are assumed to be normalized in such a way that the mixed partial derivatives $(\partial^2 / \partial x \partial z) \phi_n(x, 0)$ exist at the surface.

For every x all but finitely many eigenvalues $k_n(x)$ are purely imaginary. The indices n of the eigenvalues run through the integers $0, \pm 1, \pm 2, \dots$, and they can be assigned to the eigenvalues in such a way that for some integer $N(x)$ the following statements are true:

- (a) if $|n| > N(x)$ then $k_n(x)$ is imaginary and $|k_n(x)| = O(|n|)$;
- (b) if $n > N(x)$ then $-i k_n(x)$ is positive and increases in value as n increases;
- (c) if $n < -N(x)$ then $-i k_n(x)$ is negative and decreases in value as n decreases.

C. Acoustic field expansions. For each index n define

$$(16) \quad a_n(x) = \int_{-L}^0 \left(k_n(x) p(x, z) - i \frac{\partial}{\partial x} p(x, z) \right) \frac{1}{\rho(x, z)} \phi_n(x, z) dz + \frac{1}{k_n(x)} p(x, -H(x)) \text{jump}_{(x, -H(x))} \left[\frac{1}{\rho} \frac{\partial}{\partial z} \phi_n \right],$$

$$(17) \quad d_n(x) = 2 k_n(x) \int_{-L}^0 \frac{1}{\rho(x, z)} \phi_n(x, z)^2 dz + \frac{1}{k_n(x)} \text{jump}_{(x, -H(x))} \left[\phi_n \frac{1}{\rho} \frac{\partial}{\partial z} \phi_n \right].$$

The following expansions of the complex acoustic fields converge uniformly in the intervals $-L < z < -H(x)$ and $-H(x) < z < 0$:

$$(18) \quad p(x, z) = \sum_{n=-\infty}^{\infty} \frac{a_n(x)}{d_n(x)} \phi_n(x, z),$$

$$(19) \quad u(x, z) = \frac{1}{i\omega \rho(x, z)} \sum_{n=-\infty}^{\infty} \frac{a_n(x)}{d_n(x)} i k_n(x) \phi_n(x, z),$$

$$(20) \quad w(x, z) = \frac{1}{i\omega \rho(x, z)} \sum_{n=-\infty}^{\infty} \frac{a_n(x)}{d_n(x)} \frac{\partial}{\partial z} \phi_n(x, z).$$

In particular, expansions (19) and (20) do not exhibit Gibb's phenomenon at $z = -H(x)$ when the velocity components $u(x, z)$ and $w(x, z)$ are discontinuous at the interface.

3. THE COUPLING EQUATIONS

The functions $a_n(x)$ that appear as coefficients in the acoustic field expansions solve an infinite system of first-order, ordinary differential equations. Each equation in this system takes the form

$$(21) \quad \frac{d}{dx} a_n(x) = i k_n(x) a_n(x) - \sum_{m=-\infty}^{\infty} \frac{a_m(x)}{d_m(x)} c_{mn}(x) - i s_n(x),$$

(for $n = 0, \pm 1, \pm 2, \dots$)

where the forcing term $s_n(x)$ is a moment of the source distribution:

$$(22) \quad s_n(x) = \int_{-L}^0 S(x, z) \frac{1}{\rho(x, z)} \phi_n(x, z) dz.$$

It is possible to express the general coupling coefficient $c_{mn}(x)$ in terms of several factors to which physical meaning can be attributed. First, define the following quantities:

$$(23) \quad L_{mn}(x, z) = \frac{\partial}{\partial z} \phi_m(x, z) \frac{\partial}{\partial z} \phi_n(x, z) + \left(k_m(x) k_n(x) - \frac{\omega^2}{c(x, z)^2} \right) \phi_m(x, z) \phi_n(x, z),$$

$$(24) \quad M_{mn}(x, z) = \frac{\partial}{\partial z} \phi_m(x, z) \frac{\partial}{\partial z} \phi_n(x, z) - k_m(x) k_n(x) \phi_m(x, z) \phi_n(x, z),$$

$$(25) \quad N_{mn}(x, z) = \frac{\partial}{\partial z} \phi_m(x, z) \frac{\partial}{\partial z} \phi_n(x, z) + k_m(x) k_n(x) \phi_m(x, z) \phi_n(x, z) + \dot{H}(x) i \left(k_m(x) \phi_m(x, z) \frac{\partial}{\partial z} \phi_n(x, z) + \frac{\partial}{\partial z} \phi_m(x, z) k_n(x) \phi_n(x, z) \right),$$

$$(26) \quad D_{\parallel(z,z)}[1/\rho] = \frac{\partial}{\partial x} \left(\frac{1}{\rho(x,z)} \right) - \dot{H}(x) \frac{\partial}{\partial z} \left(\frac{1}{\rho(x,z)} \right).$$

Use these quantities to construct the following factors:

$$(27) \quad A_{mn}(x) = \int_{-L}^0 \frac{\partial}{\partial x} \left(\frac{\omega^2}{c(x,z)^2} \right) \frac{1}{\rho(x,z)} \phi_m(x,z) \phi_n(x,z) dz \\ - \int_{-L}^0 \frac{\partial}{\partial x} \left(\frac{1}{\rho(x,z)} \right) L_{mn}(x,z) dz,$$

$$(28) \quad I_{mn}(x) = \dot{H}(x) \underset{(x,-H(x))}{\text{jump}} \left[\frac{\omega^2}{c^2 \rho} \phi_m \phi_n \right],$$

$$(29) \quad J_{mn}(x) = \dot{H}(x) \underset{(x,-H(x))}{\text{jump}} [M_{mn}/\rho],$$

$$(30) \quad K_{mn}(x) = - \frac{\ddot{H}(x) i}{1 + \dot{H}(x)^2} \underset{(x,-H(x))}{\text{jump}} [M_{mn}/\rho] \\ + \frac{\dot{H}(x) i}{1 + \dot{H}(x)^2} \underset{(x,-H(x))}{\text{jump}} [N_{mn} D_{\parallel}[1/\rho]],$$

where $\ddot{H}(x) = (d^2/dx^2)H(x)$ and $m, n = 0, \pm 1, \pm 2, \dots$

The factor $A_{mn}(x)$ is associated with lateral variations in material properties of the fluids inside the waveguide. The factors $I_{mn}(x)$, $J_{mn}(x)$ and $K_{mn}(x)$ are associated with conditions at the interface. These factors can differ from zero only if material properties of the fluids are discontinuous at the interface, and if the slope or the curvature of the interface does not vanish. The following theorem states the relation of these factors to the general coupling coefficient $c_{mn}(x)$.

THEOREM. *If $m \neq n$ then*

$$(31) \quad c_{mn}(x) = \frac{1}{k_m(x) - k_n(x)} \left(A_{mn}(x) + I_{mn}(x) + \frac{k_m(x)}{k_n(x)} J_{mn}(x) \right. \\ \left. + \frac{1}{k_n(x)} K_{mn}(x) \right).$$

However, if $m = n$ then

$$(32) \quad c_{nn}(x) = -\frac{1}{2} \dot{d}_n(x) + \frac{1}{k_n(x)} J_{nn}(x) + \frac{1}{2k_n(x)^2} K_{nn}(x),$$

where $\dot{d}_n(x) = (d/dx)d_n(x)$.