

REPORT DOCUMENTATION PAGE

2

1a REPORT SECURITY CLASSIFICATION UNCLASSIFIED	1b RESTRICTIVE MARKINGS ELECTE
---	-----------------------------------

2 AUG 26 1988 LE	3 DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited.
---------------------	---

AD-A198 471

4 R(S) H	5 MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 88 - 0898
----------	---

6a NAME OF PERFORMING ORGANIZATION Colorado State University College of Vet Med	6b OFFICE SYMBOL (if applicable)	7a NAME OF MONITORING ORGANIZATION Air Force Office of Scientific Research/NL
---	-------------------------------------	--

6c ADDRESS (City, State, and ZIP Code) Department of Physiology Colorado State University Ft. Collins, Colorado 80523	7b ADDRESS (City, State, and ZIP Code) Building 410 Bolling AFB, DC 20332-6448
--	--

8a NAME OF FUNDING / SPONSORING ORGANIZATION AFOSR	8b OFFICE SYMBOL (if applicable) NL	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-86-0117
---	---	---

8c ADDRESS (City, State, and ZIP Code) Building 410 Bolling AFB, DC 20332-6448	10 SOURCE OF FUNDING NUMBERS		
	PROGRAM ELEMENT NO 61102F	PROJECT NO. 2312	TASK NO A2

11 TITLE (Include Security Classification)
Biochemical Mechanisms Controlling Bioreactivity of Adrenal Chromaffin Cells

12 PERSONAL AUTHOR(S)
P. R. Vulliet

13a TYPE OF REPORT Final	13b TIME COVERED FROM 4/86 TO 5/88	14 DATE OF REPORT (Year, Month, Day) - June 17, 1988	15 PAGE COUNT 6
-----------------------------	---------------------------------------	---	--------------------

16 SUPPLEMENTARY NOTATION

17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Adrenal Gland, Chromaffin Cells, Phosphorylation, Stress, Molecula Mechanisms
FIELD	GROUP	SUB-GROUP	
0601			

19 ABSTRACT (Continue on reverse if necessary and identify by block number)

This project has examined the biochemical mechanisms regulating the response of the rat adrenal medulla and chromaffin tissue. The responsivity of the tissue will change depending upon the treatment that the animal has received and the time after the treatment. The alteration in reactivity appears to be correlated with changes in tissue levels of the catecholamine neurotransmitters. Tissues that demonstrate altered reactivity have profound morphological changes that return to control with the return of normal reactivity. Additional biochemical factors that may regulate reactivity are currently being examined in a variety of moael systems of the chromaffin cell.

20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS	21 ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED
---	---

22a NAME OF RESPONSIBLE INDIVIDUAL Dr. William O. Berry	22b TELEPHONE (Include Area Code) (202) 767-5021	22c OFFICE SYMBOL NL
--	---	-------------------------

88 3

FINAL TECHNICAL REPORT

U.S.A.F. OFFICE OF SCIENTIFIC RESEARCH
PROJECT #86-NL-0122

**BIOCHEMICAL MECHANISMS CONTROLLING BIOREACTIVITY
OF ADRENAL CHROMAFFIN CELLS**

1. SUMMARY:

This project has examined the biochemical mechanisms regulating the response of the rat adrenal medulla and chromaffin tissue. We have found that the responsivity of the tissue will change depending upon the treatment that the animal has received and the time after the treatment. The alteration in reactivity appears to be correlated with changes in tissue levels of the catecholamine neurotransmitters. Tissues that demonstrate altered reactivity have profound morphological changes that return to control with the return of normal reactivity. Additional biochemical factors that may regulate reactivity are currently being examined in a variety of model systems of the chromaffin cell.

2. RESEARCH OBJECTIVES:

1. Optimize Conditions to provide maximal alterations in the bioreactivity of the adrenal medulla.

2. Evaluate disassociated chromaffin cells isolated from animals exhibiting altered adrenal reactivity

3. Compare the localization (compartmentalization) and activity of protein kinases and their substrates in control tissues and adrenal glands demonstrating altered reactivity or from disassociated cells derived from these tissues.

4. Quantitate the biosynthesis and number of chromaffin granules, catecholamine content, and evaluate the morphological characteristics of adrenal chromaffin vesicles exposed to treatments that alter the bioreactivity.

5. Purify and evaluate chromaffin vesicles from the adrenal medullary cells with respect to catecholamine content, protein content, protein kinase substrate content and ability to sequester calcium.

6. Quantitate the degree of phosphorylation and specific sites of phosphorylation of functionally significant proteins such as tyrosine hydroxylase, protein III, protein F1, and vesicle associated phosphoproteins from control, hyper- and hypo-reactive adrenal glands.

3. STATUS OF RESEARCH:

The research efforts from 3/1/86 to 3/1/88 have focussed on identifying factors that regulate the responsivity of the rat adrenal medulla and biochemical mechanisms that regulate the PC12 cells. The specific areas of research include: a) Development of the isolated perfused rat adrenal gland preparation b) Examination of the morphological changes that accompany changes in bioreactivity of the rat adrenal medulla c) Development of the perfused rat adrenal slice technique d) Identification of a novel protein kinase activity in PC12 cells e) Investigation of the role of protein kinase C in regulating neurite outgrowth f) Application of monoclonal antibodies for the immunocytochemical localization of tyrosine hydroxylase and g) Development of techniques for quantitating TH mRNA. Each of these areas will be briefly discussed individually below.

A. Development of the isolated perfused rat adrenal gland preparation:

The ability of the rat adrenal sympathoadrenal axis to alter its reactivity in vivo had been previously reported. To determine if this change in reactivity occurred at the level of the adrenal gland, we employed the isolated perfused adrenal gland. This preparation enabled the investigator to perform experimental manipulations on the intact animal and then assess whether molecular or biochemical changes had occurred in the isolated perfused adrenal gland. We found with experimental manipulation that the amount of catecholamines released by a maximal dose of acetylcholine could be decreased by up to 50 percent or increased by 40 percent when compared with controls. Following chronic periods of insulin induced hypoglycemia, the adrenal glands isolated from the animals were found to have altered responsivity to standard doses of acetylcholine. The changes in responsivity was found to parallel the tissue content of catecholamines.

This change in reactivity was not only dependent upon the type of experimental manipulation, but also the time after the experiment treatment. The amount of epinephrine and norepinephrine released from the isolated perfused rat adrenal gland by a fixed dose of acetylcholine was found to be decreased at one day. The responsivity of the gland gradually returned to normal values at day 5. Initial experiments were performed using immobilization stress which increased the amount of catecholamine release by up to 60 percent above controls. Hypoglycemic stress caused a decrease in both the catecholamine content of the adrenal gland and the amount that was released by acetylcholine mediated depolarization. These series of experiments clearly demonstrated that the reactivity of the adrenal gland could be altered by physiological conditioning in the laboratory rat.

B. Examination of the morphological changes that accompany changes in bioreactivity of the rat adrenal medulla:

When adrenal glands isolated from these animals were examined by electronmicroscopy, profound differences were observed. The most striking difference was in the cellular content of chromaffin granules. At day one of treatment, the number of granules was significantly decreased. At day one of



Dist		
A-1		

recovery, the granule depletion was so severe that the cells contained only a few granules. The granule content of the glands isolated from the hypoglycemic treated animals slowly returned to normal by day 7. The Golgi apparatus was markedly affected by the hypoglycemic treatment. During the early stages of cellular response, the Golgi apparatus is enlarged and appeared to be in an active secretory stage. Numerous multivesiculate bodies were observed in the cytosol, suggesting that a massive secretory event had occurred. Currently, a complete morphological analysis is being performed on these tissue to more fully quantitate the observed differences.

C. Development of the perfused rat adrenal slice technique.

To more fully characterize functional changes that occur to the adrenal medulla at the biochemical level, we have developed a novel preparation using adrenal glands isolated from the treated animals. The glands are removed and sliced in 300 um slices, perfused with an oxygenated physiological buffer solution for 30 minutes. The slices are then depolarized with acetylcholine, the endogenous secretagogue, and the amount of epinephrine or norepinephrine released from this preparation is quantitated by HPLC/EC methods. We find that the patterns of altered reactivity with this preparation are similar to those observed with the perfused rat adrenal gland.

D. Identification of a novel protein kinase activity in PC12 cells.

During the investigation of the regulation of tyrosine hydroxylase, the rate limiting enzyme in the biosynthesis of the catecholamine neurotransmitters, we discovered that this enzyme was phosphorylated in a novel site that would not be phosphorylated by any other known protein kinase. We have developed methods for the assay of this kinase activity using a synthetic peptide that is homologous with the N-terminal region of TH. We have succeeded in partially purifying this kinase activity and find that it has a molecular weight of approximately 50,000 daltons. Additional research using intact cells labelled with radioactive phosphate suggests that this kinase may play an important role in mediating some of the actions of the growth factors.

E. Investigation of the role of protein kinase C in regulating neurite outgrowth

While investigating the regulation of cell phosphorylation in PC12 cells following treatment with nerve growth factor (NGF), we observed that the NGF-mediated neurite outgrowth was potentiated by simultaneous treatment with phorbol esters. Inclusion of a specific inhibitor of protein kinase C blocked the outgrowth of these neurite processes, suggesting that protein kinase C activity is essential for neurite outgrowth. The role of other protein phosphorylation in regulating the responsiveness of the PC12 cells is currently being investigated.

F. Application of monoclonal antibodies for the immunocytochemical localization of tyrosine hydroxylase.

During the earlier stages of this project, we developed a monoclonal antibody specifically directed against rat tyrosine hydroxylase. TH was purified from rat pheochromocytoma and injected into mice and lymphocytes from the spleen isolated and fused with myeloma cells. We screened and cloned a cell line that produces a monoclonal antibody that is specific for tyrosine hydroxylase. We have used this antibody to localize TH in the growth cone of developing neurites. In other laboratories, this antibody has been used to identify and localize TH in caudate nucleus, the regenerating rat spinal cord, the retina of the fish eye, and a variety of other tissues.

G. Development of techniques for quantitating TH mRNA.

We have recently obtained a copy of cDNA coding for tyrosine hydroxylase. This DNA is in the Bluescript plasmid and also contains the endogenous promoter for the rat gene. This segment of DNA has been used to generate specific fragments that are radioactively labelled that have used to quantiate the amount of TH mRNA present in the adrenal gland. We find that during periods of chronic stress, the level of TH message increases nearly two fold and that this increase precedes the appearance of increased TH activity.

4. WRITTEN PUBLICATIONS:

Masserano, J. M., Vulliet, P. R., Tank, A. W. and Weiner, N. The Role of Tyrosine Hydroxylase in the Regulation of Catecholamine Synthesis. In: Handbook of Experimental Pharmacology, Catecholamines II. (Heidelberg, Springer-Verlag) In press, 1988.

Carbonaro, D. L., Mitchell, J. P., Hall, F. L. and Vulliet, P. Altered Bioreactivity of the Adrenal Medulla. Brain Research Bulletin, In press, 1988.

Vulliet, P. R., Hall, F. L., Mitchell, J. P. and Hardie. A Novel Growth Factor Sensitive Protein Kinase in Pheochromocytoma. Proceeding Western Pharmacology Society. 31: 255-258, 1988.

Vulliet, P. R., Fontenot, G., Mitchell, J., Kwan, S., Abel, C. and Hall, F. Induction of Tyrosine Hydroxylase and its Localization in Neurites and Growth cones of PC12 Cells Following NGF Treatment. (In preparation, to be submitted to Mol. Pharmacol. June 1988).

Hall, F., Fernyhough, P., Ishii, D. and Vulliet, P.R. Suppression of Neurite Outgrowth in PC12 Cells by Sphingosine, a Specific Inhibitor of Protein Kinase. C. J. Biol. Chem. 263: 4460-4466, 1988.

Kwan, S., Vulliet, P., Denney, R., Shen, R., Westland, K. and Abell, C. Characterization and Application of Monoclonal Antibodies to Rat Tyrosine Hydroxylase (To be submitted to J. Neurosci Methods, August 1988).

Carbonaro-Law, D., Hall, F. and Vulliet, P. Morphological Alterations of the Rat Adrenal Medulla Following Periods of Chronic Hypoglycemic Stress. In: Stress: Neurochemical and Humoral Mechanisms. ed. by G. van Loon, R. Kvetnansky and J. Axelrod. Gordon and Breach Publishers. In press, July 1988.

Vulliet, P. and D. G. Hardie. Phosphorylation Sites on Tyrosine Hydroxylase. In: Signal Transduction and Protein Phosphorylation. ed. by L. G. Heilmeyer (New York, Plenum Press, 1987) pp 265-269.

Fontenot, G. K., Cass, W. and Vulliet, P. R. Changes in Rat Adrenal Tyrosine Hydroxylase mRNA Levels following Chronic Administration of Amphetamine. Proc. Western Pharmacol. Soc. 30: 263-265, 1987.

Hall, F. L. and Vulliet, P. R. Immunocytochemical Localization of Tyrosine Hydroxylase in the Growth Cones of PC12 Pheochromocytoma Cells. Proc. Western Pharmacol. Soc. 30: 45-49, 1987.

Mitchell, J. and Vulliet, P. R. Altered Reactivity of the Rat Adrenal Medulla Following Periods of Chronic Stress. In: Molecular Mechanisms of Neuronal Responsiveness ed. by Y. Ehrlich, W. Berry and Robert Lennox (New York, Plenum Press, 1987) pp 367-374.

Vulliet, P. R., Loskutoff, N. and Kraemer, D. A Technique of Embryo Transfer in the Laboratory Rat. Proc. Western Pharmacol. Soc. 29: 315-318, 1986.

Campbell, D. C., Hardie, D. G. and Vulliet, P. R. Identification of four Phosphorylation Sites in the N-terminal Region of Tyrosine Hydroxylase. J. Biological Chem. 261: 10489-10492, 1986.

Roskoski, R., Vulliet, P. and Glass, D. Phosphorylation of Tyrosine Hydroxylase by Cyclic GMP-dependent Protein Kinase. J. Neurochem. 48: 840-845, 1987.

Shen, R., Hamilton-Byrd, E., Vulliet, P., Kwan, S and Abell, C. An Improved ^{14}C -Trapping Microassay for Tyrosine Hydroxylase Activity in Rat Striatal Synaptosomes. J. Neuroscience Methods 16: 163-173, 1986.

Vulliet, P. R., Woodgett, J. R. Ferrari, S. and Hardie, D. G. Characterization of the Sites Phosphorylated on Tyrosine Hydroxylase by the Ca^{2+} and Phospholipid-dependent Protein Kinase, Calmodulin-dependent Multiprotein Kinase and Cyclic AMP dependent Protein Kinase. FEBS Letter 182: 335-339, 1985.

5. PROFESSIONAL PERSONNEL:

Philip R. Vulliet, Ph.D., DVM -50 % time

Fred Hall, Ph.D. 100 % time

Denise Carbonaro, M. S. 100 % time

Greg Fontenot, M. S. 100 % time

Jeff Mitchell, B. S., 100 % time

6. PROFESSIONAL INTERACTIONS:

Dr. Creed Abell
Department of Medicinal Chemistry
School of Pharmacy
University of Texas
Austin, TX

Dr. Philip Cohen, FRS, FRSE
MRC Protein Phosphorylation Group
Department of Biochemistry
University of Dundee
Dundee DD1 4HN, Scotland

Dr. Graham Hardie
MRC Protein Phosphorylation Group
Department of Biochemistry
University of Dundee
Dundee DD1 4HN, Scotland

Dr. Karen O'Malley
Department of Anatomy and Neurobiology
Washington University
St. Louis, MO

Dr. Howard Norness
Department of Anatomy and Neurobiology
Colorado State University
Ft. Collins, CO 80523

END

DATE

FILMED

DTIC

10-88