

AD-A198 581

①

OFFICE OF NAVAL RESEARCH

CONTRACT NO. N00014-86-K-0772

TECHNICAL REPORT NO. 11

Reversed-Phase Separation of a Standard PAH Mixture
on an Adamantyl Surface

by

S. S. Yang and R. K. Gilpin

Liquid Crystal Institute*
Kent State University
Kent, OH 44242

*Subcontractor to
Liquid Crystalline Polymer Research Center
University of Connecticut
Storrs, CT 06268

Prepared for Publication

in

Journal of Chromatography

July 20, 1988

REPRODUCTION IN WHOLE OR IN PART IS PERMITTED FOR ANY
PURPOSE OF THE UNITED STATES GOVERNMENT.

THIS DOCUMENT HAS BEEN APPROVED FOR PUBLIC RELEASE
AND SALE; ITS DISTRIBUTION IS UNLIMITED.

DTIC
ELECTE
AUG 02 1988
S H D

REPORT DOCUMENTATION PAGE

1a REPORT SECURITY CLASSIFICATION Unclassified		1b RESTRICTIVE MARKINGS None	
2a SECURITY CLASSIFICATION AUTHORITY		3 DISTRIBUTION/AVAILABILITY OF REPORT Approved for Public Release, Distribution Unlimited	
2b DECLASSIFICATION/DOWNGRADING SCHEDULE			
4 PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. 11		5 MONITORING ORGANIZATION REPORT NUMBER(S)	
6a NAME OF PERFORMING ORGANIZATION University of Connecticut	6b OFFICE SYMBOL (if applicable)	7a NAME OF MONITORING ORGANIZATION Office of Naval Research	
6c ADDRESS (City, State, and ZIP Code) Storrs, CT 06268		7b ADDRESS (City, State, and ZIP Code) 800 North Quincy Avenue Arlington, VA 22217	
8a NAME OF FUNDING/SPONSORING ORGANIZATION	8b OFFICE SYMBOL (if applicable) ONR	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-86-K-0772	
8c ADDRESS (City, State, and ZIP Code) 800 North Quincy Avenue Arlington, VA 22217		10 SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO	PROJECT NO
		TASK NO	WORK UNIT ACCESSION NO
11 TITLE (Include Security Classification) Reversed-Phase Separation of a Standard PAH Mixture on an Adamantyl Surface (Unclassified)			
12 PERSONAL AUTHOR(S) S. S. Yang and R. K. Gilpin			
13a TYPE OF REPORT Interim, Technical	13b TIME COVERED FROM TO 07/20/88	14 DATE OF REPORT (Year Month, Day) 88-July-20	15 PAGE COUNT 13
16 SUPPLEMENTARY NOTES Prepared for publication in <u>J. of Chromatography</u> Research carried out at Kent State University (Subcontractor) (LCPRC Publication No.)			
17 COSATI CODES		18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB GROUP	
		High Performance Liquid Chromatography (HPLC)	
		Polycyclic Aromatic Hydrocarbons (PAH)	
19 ABSTRACT (Continue on reverse if necessary and identify by block number) High performance liquid chromatography (HPLC) is an important technique for the analysis of polycyclic aromatic hydrocarbons (PAH). In carrying out the chromatographic separations, octadecyl (C ₁₈) phases often are used because of their excellent selectivity towards PAH. Recently, Sander and Wise have studied and attempted to classify the various parameters which influence solute retention for this class of compounds. Surface coverage and solute shape have been found to be major factors which influence retention behavior. The elution order for polycyclic aromatic hydrocarbons which differ in planarity may reverse depending on structural differences in the C ₁₈ phase being used. As a result of their studies, Sander and Wise have proposed that selected PHAS solutes (e.g., benzo[a]pyrene - BaP, phenanthro[3,4-c]phenanthrene-PhPh, and 1,2:3,4:5,6:7,8-tetrabenzonaphthalene - TBN.) can be used to classify phases in terms of their selectivity. (continued on reverse ->)			
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21 ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a NAME OF RESPONSIBLE INDIVIDUAL Dr. Kenneth J. Wynne		22b TELEPHONE (Include Area Code) (202) 696-4410	22c OFFICE SYMBOL ONR

Block 19. (continued from front)

Recently we have reported an unique stationary phase synthesized by exhaustively modifying porous silica with bulky rigid molecules of adamantane. This surface has been shown to produce good peak shapes for basic solutes even under less than favorable chromatographic conditions. In the current work the selectivity of the adamantyl surface has been studied. In carrying out the evaluation, the polycyclic aromatic hydrocarbon mixture recommended by Sander and Wise has been used.

Reversed-Phase Separation of a Standard PAH mixture
on an Adamantyl Surface

S. S. Yang and R. K. Gilpin*

Department of Chemistry

Kent State University

Kent, OH 44242

* author to whom correspondence should be addressed

In Press J. of Chromatogr.

aromatic hydrocarbon mixture recommended by Sander and Wise
has been used.

EXPERIMENTAL SECTION

Reagent: Adamantylethyltrichlorosilane was purchased from Petrarch System Inc. (Levittown, PA) and was used as received. LiChrosorb Si100 (dp ~ 10 μ m) was obtained from E. Merck (Darmstadt, West Germany). HPLC grade acetonitrile was purchased from Aldrich Chemical Co. (Milwaukee, WI). Water was purified using a Milli-Q water system (Millipore Co., El Paso, TX). A test mixture containing Bap, PhPh, and TBN in acetonitrile was supplied by the National Bureau of Standards (NBS)(Gaithersburg, MD)

Preparation of adamantyl phase: Adamantylethyltrichlorosilane was used to chemically modify silica as previously described (4). Subsequently, this material was packed into a 4.6 mm i.d. X 150 mm stainless steel column using a dynamic packing procedure (5).

Chromatography: Chromatographic experiments were carried out using a LC/9560 ternary gradient liquid chromatograph (IBM Instruments Inc.) equipped with a UV detector. The column was evaluated using a mobile phase of acetonitrile/water (v/v, 85:15) at a flow rate 2 ml/min.

RESULT AND DISCUSSION

A representative separation of the NBS test mixture on the adamantyl phase is shown in figure 1. Capacity factors (k') of each PAH compound and selectivity values (α) are summarized in table 1. For comparison purposes, data obtained from chromatograms, supplied with the test mixture (6) also are listed for several different C_{18} phases.

The elution order of the PAH compounds (BaP < PhPh < TBN) is similar to a commercial monomeric C_{18} phase (e.g., Beckman Ultrasphere ODS phase). Likewise, selectivities, $\alpha(\text{TBN/BaP}) = 2.10$, $\alpha(\text{PhPh/BaP}) = 1.47$, are closest to this same commercial column. However, the k' data are about 1/3 that obtained from the Beckman column.

Presumably based on the above data a number of other solutes may be expected to behave similarly and the adamantyl surface will be applicable to separations now being carried out on C_{18} columns with similar selectivity. In a more general sense, this probably should not hold true based on geometrical considerations and the proposed "slot" model (2) for PAH solutes. Because of its ball-like structure the adamantyl molecule effectively shields unreacted silanols even for small basic solutes such as aniline (3). Since the surface structure is rigid compared to conventional linear n -alkyl phases, it should not only be useful for carrying out

numerous applications but also should provide an important tool for testing interaction/interfacial models.

REFERENCES

- (1) L. C. Sander and S. A. Wise, *Anal. Chem.*, 56(1984)504.
- (2) L. C. Sander and S. A. Wise, *Advances in Chromatography*,
(J. C. Gidding, ed.) Marcel Dekker, Inc., New York,
1986, vol.25, pp 139.
- (3) S. S. Yang and R. K. Gilpin, *Anal. Chem.*, 59(1987)2750.
- (4) R. K. Gilpin and J. A. Squires, *J. Chromatogr. Sci.*,
19(1981)195.
- (5) R. K. Gilpin and W. R. Sisco, *J. Chromatogr.*,
194(1980)285.
- (6) L. C. Sander and S. A. Wise, *J. High Resolut. Chromatogr.*
Commun., in press.

Table 1. Comparison of PAH retention & column selectivity

Column*	capacity factor (k')#			selectivity (α)	
	Bap	PhPh	TBN	PhPh/BaP	TBN/BAP
A	1.55	2.28	3.26	1.47	2.10
B	4.94	5.52	9.77	1.12	1.98
C	3.96	3.96	7.20	1.00	1.82
D	5.56	4.42	7.76	0.79	1.40
E	2.48	1.65	2.68	0.67	1.08
F	3.85	1.97	2.86	0.51	0.74
G	5.57	1.49	1.85	0.27	0.33

*. Columns: A = adamnatyl, B = Beckman Ultrasphere ODS,
 C = Whatman Partisphere C₁₈, D = Whatman Partisil ODS-2,
 E = Vydac TP low load custom synthesis, F = Vydac 201TP,
 G = Vydac TP high load custom synthesis.

#. Solutes: BaP = benzo[a]pyrene,
 PhPh = phenanthro[3,4-c]phenanthrene,
 TBN = 1,2:3,4:5,6:7,8-tetrabenzonaphthalene.
 k' data except column A are from ref.(6).


Figure Caption

Figure 1: Chromatogram of the NBS test mixture.

Column: adamantyl (4.6 mm i.d. X 150 mm).

Mobile phase: acetonitrile-water, 85/15, v/v.

Flow rate: 2 ml/min.



ACKNOWLEDGEMENT

Support from DARPA-ONR Contract N0014-86-K-0772 is
acknowledged.

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Duda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code LS2 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Bojic Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 92132	1

