

30 5265-MA-01-5
②

AD-A198 586

RESEARCH PROJECT: Numerical Optimization
Principal Investigator: Francesco ZIRILLI
Contractor: Università di Roma "La Sapienza"
Contract Number: DAJA45-86-C-0028
Fifth Periodic Report
Jan. '88 - July '88

DTIC FILE COPY

The research reported in this document has been made possible through the support and sponsorship of the U.S. Government through its European Research Office of the U.S. Army.

This report is intended only for the internal management use of the Contractor and the U.S. Government.

DTIC
ELECTE
S AUG 24 1988 D
E

ADA 198586

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Numerical Optimization		5. TYPE OF REPORT & PERIOD COVERED Periodic Jan 88-Jul 88
7. AUTHOR(s) Francesco Zirilli		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Dipartimento di Matematica G.Castelnuovo Università di Roma "La Sapienza" 00185 ROMA ITALY		8. CONTRACT OR GRANT NUMBER(s) DAJA 45-86-C-0028
11. CONTROLLING OFFICE NAME AND ADDRESS European research office of the U.S. Army 223 Old Marylebone Road London NW1 5TH (ENGLAND)		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE Jul. 88
		13. NUMBER OF PAGES
		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) UNLIMITED		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) UNLIMITED		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) NUMERICAL OPTIMIZATION COMPLEMENTARITY PROBLEMS LINEAR PROGRAMMING		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This is the Fifth Periodic Report on Contract DAJA 45-86-C-0028; contains: (1) Statement of scientific work done during the reporting period		

DD FORM 1473 1 JAN 73 EDITION OF 1 NOV 65 IS OBSOLETE

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

- A -

I N D E X

1. Statement of scientific work done during the reporting period.....pag. 1

2. Research plans for the immediate future..... " 5

3. Administrative actions..... " 6

4. Appendix 1: L. Misici, F. Zirilli: "The inverse gravimetry problem:-An application to the northern San Francisco craton granite" submitted to J. of the Geological Society London..... " 7

5. Appendix 2..... " 20

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



§1. STATEMENT OF SCIENTIFIC WORK DURING THE REPORTING PERIOD.

Let \mathbb{R}^N be the N dimensional real euclidean space, $\underline{x} = (x_1, x_2, \dots, x_N)^T \in \mathbb{R}^N$ a generic vector, the superscript T means transpose, $\langle \cdot, \cdot \rangle$ is the euclidean scalar product and $\|\cdot\|$ the euclidean norm.

We have considered the linear programming problem; that is:

Problem 1 (Linear programming) Given $\underline{b}, \underline{c} \in \mathbb{R}^N$ and A an $m \times N$ matrix ($m < N$), solve the following minimization problem:

$$\begin{aligned} & \text{minimize } \langle \underline{c}, \underline{x} \rangle \\ & \text{for } \underline{x} \text{ such that} \\ & \underline{x} \geq \underline{0} \\ & A\underline{x} - \underline{b} \geq \underline{0} \end{aligned}$$

where $\underline{x} \geq \underline{0}$ means that each component of \underline{x} is greater or equal to zero and similarly $A\underline{x} - \underline{b} \geq \underline{0}$.

The Linear programming problem is transformed into a Linear Complementarity Problem using the Lagrange multipliers as follows:

Let

$$\underline{z} = \begin{pmatrix} \underline{x} \\ \underline{y} \end{pmatrix} \quad \underline{\tilde{b}} = \begin{pmatrix} -\underline{c} \\ \underline{b} \end{pmatrix} \quad \tilde{A} = \begin{pmatrix} 0 & -A^T \\ A & 0 \end{pmatrix}$$

Problem 1 becomes

Problem 2 (Linear Complementarity Problem). Find \underline{z} such that:

$$\begin{aligned} \underline{z} &\geq 0 \\ \tilde{A}\underline{z} - \tilde{b} &\geq 0 \\ \langle \underline{z}, \tilde{A}\underline{z} - \tilde{b} \rangle &= 0 \end{aligned}$$

Finally the Linear Complementarity Problem is transformed into a global optimization problem via the transformation given at page 1 of the Fourth Periodic Report. In this way the linear programming problem is reduced to the problem of finding the minimizers of a function $F(\underline{z}) \geq 0$ such that $F = 0$ at the minimizers.

If the original linear programming problem has a unique solution the corresponding minimization problem has a unique global minimizer, but unfortunately the function F may have a rather complicated set of local minimizers including "cylindrical valleys". A direct application of a local minimization techniques such as GRACON (see Fourth Periodic Report) is unsuccessful. So that several modified algorithms have been considered the most promising ones are:

- (i) A perturbation method. Let $\epsilon \in \mathbb{R}, \epsilon > 0$, the matrix \tilde{A} is substituted with $\tilde{A}(\epsilon) = \tilde{A} + \epsilon I$ where I is the identity matrix. The idea is to find the solution $\underline{z}(\epsilon)$ of the problem correspondent to the matrix $\tilde{A}(\epsilon)$ and to let ϵ go to zero. If ϵ goes to zero too quickly the conjugate gradient method may remain trapped in a local minimizer, if ϵ goes to zero too slowly the computational cost becomes

excessive. In Table 1 the results obtained with this method are indicated with D0D4 D0D;

- (ii) Avoiding the cylindrical valleys. When the conjugate gradient method remains trapped in a cylindrical valley the minimization procedure is stopped, the direction of the valley is computed and a large step is taken in the direction of the valley in the "descending" direction, finally the conjugate gradient minimization algorithm is started again. In Table 1 the results obtained with this method are indicated with D0D4 D0D9.

The test problems considered in Table 1 have been constructed following the suggestions of De Leone and Mangasarian (see Appendix 2). The numerical experience has been obtained on a VAX 8530 with VMS 4,5 Operating System.

TABLE 1

DOD4 DCD			DOD4 DOD9		
M	NFEV	T	M	NFEV	T
50	694	2".97	50	315	2".24
100	1502	7".87	100	716	4".21
500	4176	1'.35".08	500	2154	43".44
1000	9620	7'. 4".74	1000	1333	55".39
			5000	12548	46'.26".80
			10000	14241	1 ^h .54'.18".51

Legenda

M dimension of the complementarity problem ($z \in \mathbb{R}^M$)

NFEV number of function and gradient evaluation of the function that must be minimized

T running time on the VAX 8530 (Example 1^h.54'.18".51 = 1 hour 54 minutes, 18 seconds, 51 seconds/100.

Tolerance required: Final function value $\leq 10^{-20}$

Finally some geophysical application of linear programming has been considered (see Appendix 1) since in our opinion the random generated test problems are not completely satisfactory.

§2. RESEARCH PLANS FOR THE IMMEDIATE FUTURE.

In the immediate future we plan to pursue the following objectives:

- (i) study Karmarkar and Renegar methods for linear programming in the context of continuation methods;
- (ii) study the nonlinear complementarity problem.

§3. ADMINISTRATIVE ACTIONS.

None.

§4. Appendix 1: L. Misici, F. Zirilli: "The inverse gravimetry problem: An application to the northern San Francisco craton granite" submitted to J. of the Geological Society London.

**THE INVERSE GRAVIMETRY PROBLEM: AN APPLICATION
TO THE NORTHERN SAN FRANCISCO CRATON GRANITE***

Luciano Misici
*Dipartimento di Matematica e Fisica
Università di Camerino
62032 Camerino (MC) Italy*

Francesco Zirilli
*Dipartimento di Matematica "G. Castelnuovo"
Università di Roma "La Sapienza"
00185 Roma Italy*

Abstract

From the knowledge of the anomalies of the gravitational field we reconstruct the mass density distribution in a large region of the state of Bahia (Brazil). This inverse gravimetry problem has been translated in a linear programming problem and solved using the simplex method. Both two and three dimensional models have been considered.

1. Introduction

In recent years a great deal of attention has been attracted by the study of several inverse problems in science and technology. In particular in science we mention the inverse problem of quantum mechanics, that is the reconstruction of a potential from its scattering data [1] and the famous problem "can you hear the shape of a drum", that is the reconstruction of the shape of a region in Euclidean space from the knowledge of spectral properties of suitable differential operators [2]. In technology we mention the radar technology that is the use of electromagnetic waves to detect not visible objects, the sonar and ecography technology that is the use of acoustics waves to detect properties of regions that are not directly accessible because they are underwater or inside the human body.

* The research reported in this document has been made possible through the support and sponsorship of the U.S. Government through the European Research Office of the U.S. Army under contract n. DAJA 45-86-C-0028 at the University of Roma "La Sapienza" and of the Ministero Pubblica Istruzione under contract 60% 1987 at the University of Camerino.

There are many inverse problems in geophysics both of fundamental interest such as the reconstruction of the Earth structure from knowledge of elastic, gravimetric and geomagnetic data or of more applied nature such as the use of similar data in geophysical prospecting to localize gas or oil.

In this paper we consider an inverse problem in gravimetry. Let x, y, z be cartesian coordinate in the three dimensional space, given a mass density $\rho(x, y, z)$ in a certain region Ω , the direct problem of gravimetry consists in finding the gravitational potential $V(x, y, z)$ generated by ρ , that is in solving the Poisson's equation:

$$\Delta V = -4\pi f \rho \quad \text{in } \Omega \quad (1.1)$$

where $\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$ is the laplacian, $f = 6.67 \cdot 10^{-8} \text{ cm}^3 \cdot \text{g}/\text{sec}^2$ is the gravitational constant and suitable boundary conditions on the boundary of Ω , $\partial\Omega$, are specified. Vice versa given the gravitational potential $V(x, y, z)$ in a certain region Ω , the inverse gravimetry problem consists in finding the mass density $\rho(x, y, z)$ that generates V , that is in solving the inverse Poisson's equation

$$\Delta^{-1} \rho = -\frac{1}{4\pi f} V \quad \text{in } \Omega \quad (1.2)$$

where Δ^{-1} is the inverse laplacian with the appropriate boundary conditions on $\partial\Omega$.

Almost all inverse problems and in particular the inverse gravimetry problem are ill posed, that is, to an arbitrarily small perturbation of the data V can correspond an arbitrarily large perturbation of the solution ρ .

The great difficult of solving the inverse gravimetry problem in practical situations is due to its ill posedness since the values of the gravitational field, that is the data, are obtained from experiments and so are affected by errors. In order to avoid this difficulty and restore a well behaved dependence of the solution ρ from the data V it is very useful to introduce some a priori constraints that ρ should satisfy.

In [3], [4] [5] using these ideas the inverse gravimetry problem has been reduced to a linear programming problem in a way that we will see later. In this paper using this linear programming formulation of the inverse gravimetry problem we study the northern San Francisco craton granite in Brazil on the basis of measurements of the residual Bouguer anomaly of the gravitational field taken from [6] and [7], see Fig.1. From these data we reconstruct a two dimensional section of the mass density along the BB' segment of Fig.1. This section is about 270Km long and 20Km deep, see Fig.3. Moreover we reconstruct a three dimensional section of the mass density in the region bounded by the dotted rectangle of Fig.1. This section on the surface of the Earth is a rectangle of about 240Km \times 189Km and is about 20Km deep, see Fig.4.

We have found that in the center of this three dimensional region there is a body of granite of mass density $\rho = 2.57\text{g}/\text{cm}^3$ while the average mass density of this region is $\rho = 2.67\text{g}/\text{cm}^3$. This body in the BB' segment is surfacing for about 30Km and is about 67Km long on its bottom which is about 16Km deep, see Fig.3. These results are confirmed by the three dimensional reconstruction of ρ (Fig.4) and they are in good agreement with the results obtained by Ussami and Bott in [6]. The granite body found by Ussami and Bott in [6] is a little bit smaller than ours.

In section 2 we describe our mathematical model and we reduce the inverse gravimetry problem to a linear programming problem. In section 3 we present the data that we have used and the results obtained in the northern San Francisco craton granite (Brazil) using the mathematical model described in section 2.

2. The Mathematical model

Here we will follow the work of Safon, Vasseur and Cuer [4], it is well known that, using the Green's function $G(x, y, z)$ [8], the equation (1,2) can be reformulated as an integral equation as follows

$$\int_{\Omega} G(x, y, z, x', y', z') \rho(x', y', z') dx' dy' dz' = V(x, y, z) \quad (2.1)$$

Let us cut the domain Ω in a great number N of cubic domains ω_n , the function $\rho(x, y, z)$ will be approximated on each cube ω_n by a constant ρ_n and the Green's function G will be approximated by a matrix $\mathcal{G} = ((g_{ij}))$; $i = 1, 2, \dots, M$; $j = 1, 2, \dots, N$. Finally let V_i , $i = 1, 2, \dots, M$ ($M \leq N$) be the measurements of the gravitational field $V(x, y, z)$ in x_i, y_i, z_i that belongs to the cubic domain $\omega_i, i = 1, 2, \dots, M$. So that equation (2.1) can be discretized as follows:

$$\sum_{j=1}^N g_{ij} \rho_j = V_i, \quad i = 1, 2, \dots, M \quad (2.2)$$

where $g_{ij} = \int_{\omega_j} G(x_i, y_i, z_i, x', y', z') dx' dy' dz'$. In general is natural to assume that the data V_i are available only on cubic domains ω_i that have one side on the accessible surface of the region Ω , so that $M < N$, see Fig.4 and the linear system (2.2) is underdetermined. Moreover since the data V_i , $i = 1, 2, \dots, M$ are known only with a certain error $\epsilon_i > 0$, $i = 1, 2, \dots, M$ we substitute the system of linear equations (2.2) with the system of linear inequalities

$$V_i - \epsilon_i \leq \sum_{j=1}^N g_{ij} \rho_j \leq V_i + \epsilon_i, \quad i = 1, 2, \dots, M \quad (2.3)$$

The mass density ρ is greater or equal to zero, so that we may impose

$$\rho_j \geq 0, \quad j = 1, 2, \dots, N \quad (2.4)$$

or more realistically in a geophysical problem

$$\rho_j^{\min} \leq \rho_j \leq \rho_j^{\max}, \quad j = 1, 2, \dots, N \quad (2.5)$$

where ρ_j^{\min} and ρ_j^{\max} , $j = 1, 2, \dots, N$ are the minimum and the maximum values for ρ , assigned on the basis of physical intuition. Both the linear system (2.2) or the more realistic system of inequalities (2.3), (2.5) if compatible have in general an infinite number of solutions.

In order to restore uniqueness we introduce a functional to be optimized on the set of points satisfying (2.3), (2.5), that is

$$J = \sum_{k=1}^L \rho_k \text{vol}(\omega_k) \quad (2.6)$$

where $vol(\omega_{k_i})$ is the volume of the cube ω_{k_i} , so that J is the total mass of the body Ω or of some part of it.

The linear programming problem that we consider is the following one: minimize the functional J given by (2.6) subject to the constraints (2.3), (2.5).

In section 3 we will also use a two-dimensional model, in this case we assume that the region Ω has as infinite extension in the horizontal x coordinate and that ρ is a function of z , the vertical coordinate, and y only. Reasoning as in the three-dimensional case with an appropriate Green's function G and data V_i , $i = 1, 2, \dots, M$ it is easy to obtain a linear programming problem analogous to the one obtained for the three-dimensional model given by (2.3), (2.5), (2.6).

3. Results and conclusions

Here we will consider two special applications of the method described in section 2. That is a two dimensional vertical section and a three dimensional slice of a large region in the state of Bahia in Brazil. In our computations the gravitational anomaly is measured in *mgals* and the distances in *Km*. The gravitational anomalies, that is the residual Bouguer anomalies, are reported in Table 1 for the three-dimensional problem and in Fig.2 for the two-dimensional problem. Moreover we assume that the average mass density in the regions considered is $\rho_0 = 2.67g/cm^3$. The data of table 1 and Fig.2 have been regularized in order to avoid numerical instability due to the ill posedness of the problem. In particular the data of Fig.2, that are taken along the BB' segment of Fig.1, are highly irregular outside the central region of Fig.2 where the maximum of the gravitational anomaly is attained so that, as shown in Fig.2, the data have been regularized outside this region. For the three-dimensional case the numbers in bold face of Table 1 are real gravitational anomalies, these are the data taken in the rectangle AA'C'C of Fig.1, while the numbers not in bold face in Table 1, that is the data of the region between the dotted rectangle and AA'C'C of Fig.1, have been regularized solving some suitable direct problem. In order to choose the functional J given by (2.6) we have used an empiric formula [9] that estimates the maximum deepness Δz of a body that generates a certain gravitational anomaly V_i , $i = 1, 2, \dots, M$ on the surface, that is

$$\Delta z = 0.65 \frac{\max_i |V_i|}{\max_i |\nabla V_i|} \quad (3.1)$$

where ∇V_i is a numerical approximation to the gradient of V_i expressed in *mgals/Km*. With our data we obtain $\Delta z \simeq 16Km$. So that on the basis of this estimate for Δz and of the conclusion of [6] that the body is surfacing we have chosen

$$J = \sum_{\{j | 0Km \leq z \leq 16Km\}} \rho_j vol \omega_j \quad (3.2)$$

In particular in the two-dimensional case along the BB' segment of Fig.1 we have considered 37 data points taken on the surface at a distance of 7.5Km one from the other. Moreover we assume that the body we are looking for is localized in the region $z \leq 20Km$. So that the rectangle we are working on, as shown in Fig.3, is $(7.5Km \times 37 = 277.5Km) \times 20Km$ and has been divided in small rectangles ω_j , $j = 1, 2, \dots, 370$ where ω_j is $7.5Km \times 2Km$ (see Fig.3) and j runs from left

to right and from top to bottom from 1 to 370. The bound (2.5) assumed on ρ_j , $j = 1, 2, \dots, 370$ have been chosen so that the feasible region is as small as possible.

The gravimetric problem that has been solved for the two dimensional case is the following one: find the values of ρ_j , $j = 1, 2, \dots, 370$ that minimize

$$J = \sum_{j=1}^{370} \rho_j \text{vol} \omega_j \quad (3.3)$$

subject to

$$\begin{aligned} V_i - \epsilon_i &\leq \sum_{j=1}^{370} g_{ij} \rho_j \leq V_i + \epsilon_i \quad ; \quad i = 1, 2, \dots, 37 \\ 2.57g/cm^3 &\leq \rho_j \leq 2.69g/cm^3 \quad ; \quad 1 \leq j \leq 37 \\ 2.57g/cm^3 &\leq \rho_j \leq 2.675g/cm^3 \quad ; \quad 38 \leq j \leq 74 \\ 2.57g/cm^3 &\leq \rho_j \leq 2.67g/cm^3 \quad ; \quad 75 \leq j \leq 370 \end{aligned} \quad (3.4)$$

where $V_i = \delta_i + \gamma_i$ with $\delta_i = 2.67g/cm^3 \sum_{j=1}^{370} g_{ij}$ and γ_i the anomalies shown in Fig.2 while $\epsilon_i = 0.05\gamma_i$. This linear programming problem (3.3),(3.4) has been solved using the FORTRAN package [10] and the solution found is shown in Fig.3.

In the three dimensional problem the dotted rectangle of Fig.1 is of dimension 240Km in the BB' direction (y direction) times 189Km in the AC direction (x direction). This dotted rectangle has been divided in $16 \times 7 = 112$ small rectangles of dimension 15Km in the BB' direction times 27Km in the AC direction, the gravitational anomaly data γ_i are taken in the central points of these small rectangles $(x_i, y_j, 0)$, $i = 1, 2, \dots, 7$, $j = 1, 2, \dots, 16$.

TABLE 1.

	y ₁	y ₂	y ₃	y ₄	y ₅	y ₆	y ₇	y ₈	y ₉	y ₁₀	y ₁₁	y ₁₂	y ₁₃	y ₁₄	y ₁₅	y ₁₆
x ₁	-0.18	-0.25	-0.35	-0.52	-0.82	-1.41	-2.69	-4.57	-5.34	-4.57	-2.69	-1.41	-0.82	-0.52	-0.35	-0.25
x ₂	-0.23	-0.32	-0.48	-0.77	-1.37	-9.60	-17.65	-22.65	-48.76	-37.63	-23.04	-2.94	-1.37	-0.77	-0.48	-0.32
x ₃	-0.26	-0.38	-0.58	-0.95	-1.73	-18.24	-28.98	-40.13	-53.18	-35.14	-14.40	-3.74	-1.73	-0.95	-0.58	-0.38
x ₄	-0.27	-0.40	-0.61	-1.01	-1.83	-26.88	-40.32	-57.60	-57.60	-32.64	-5.76	-3.92	-1.83	-1.01	-0.61	-0.40
x ₅	-0.26	-0.38	-0.58	-0.95	-1.73	-12.10	-23.04	-42.05	-52.80	-39.36	-22.08	-3.74	-1.73	-0.95	-0.58	-0.38
x ₆	-0.23	-0.32	-0.48	-0.77	-1.37	2.69	-5.76	-26.50	-48.00	-46.08	-38.04	-2.94	-1.37	-0.77	-0.48	-0.32
x ₇	-0.18	-0.25	-0.35	-0.52	-0.82	-1.41	-2.69	-4.57	-5.34	-4.57	-2.69	-1.41	-0.82	-0.52	-0.35	-0.25

The parallelepiped considered is the one of base the dotted rectangle of Fig.1 and deepness 20Km. This parallelepiped has been divided in $7 \times 16 \times 5 = 560$ small parallelepipeds ω_j , $j = 1, 2, \dots, 560$ of dimensions 27Km \times 15Km \times 4Km in the x, y, z directions respectively (see Fig. 4) and j runs as shown in Fig. 4.

The gravimetric problem that has been solved with the FORTRAN package [10] is the following one: find the values of ρ_j , $j = 1, 2, \dots, 560$ that minimize J given by (3.2) subject to:

$$V_i - \epsilon_i \leq \sum_{j=1}^{560} g_{i,j} \rho_j \leq V_i + \epsilon_i \quad i = 1, 2, \dots, 112$$
$$2.57g/cm^3 \leq \rho_j \leq 2.697g/cm^3 \quad ; \quad 1 \leq j \leq 112$$
$$2.57g/cm^3 \leq \rho_j \leq 2.672g/cm^3 \quad ; \quad 113 \leq j \leq 224$$
$$2.57g/cm^3 \leq \rho_j \leq 2.67g/cm^3 \quad ; \quad 225 \leq j \leq 560$$

where ϵ_i and $V_i = \delta_i + \gamma_i$ are defined as in the two-dimensional case. The results obtained are shown in Fig.4.

As shown in Fig.3 for the two-dimensional problem we have found a granite body about 16Km deep with density $2.57g/cm^3$, that is $0.1g/cm^3$ smaller than the surrounding medium, this body is about 30Km long on the surface and 67Km long on the bottom. The results of the three-dimensional study, shown in Fig.4, confirm the results of Fig.3. This results are in qualitative agreement with the ones obtained by Ussami and Bot in [6].

REFERENCES

- [1] I.M.Gelfand, B.M.Levitan, *On the determination of a differential equation from its spectral function.* Amer. Math. Soc. Transl., (1955), 253-304.
- [2] M.H.Protter, *Can you hear the shape of a drum? Revisited.* SIAM Review 29, (1987), 185-199.
- [3] V.Richard, R.Baier, M.Cuer, *An attempt to formulate well posed questions in gravity. Application of linear inverse techniques to mining exploration.* Geophysics 49, 10, 1781-1793.
- [4] C.Safon, G.Vasseur, M.Cuer, *Some applications of linear programming to the inverse gravity problem.* Geophysics 42, (1977), 1215-1229.
- [5] M.Cuer, *Des questions bien posees dans des problemes inverses de gravimetrie et geomagnetisme. Une nouvelle application de la programmation lineaire.* (1984) These, Université des Sciences et Techniques du Languedoc Montpellier.
- [6] N.Ussami, M.H.P.Bott, *A gravity study of the northern San Francisco Craton Granite (Brazil)* to appear in J. of the geological society of London.
- [7] A.C. Motta, R.A. Gomes, I.M. Delgado, L.P. Siqueira, A.J. Pedreira, *Feições gravimétricas e magnéticas do craton do São Francisco* in J.F. Mascarenhas (Ed.), Anais do simpósio sobre o craton do São Francisco e suas faixas marginais. SBG/CPM, Salvador, 18-20.
- [8] R. Courant, D. Hilbert, *Methods of Mathematical physics*, 2, chapter IV. Interscience Publishers New York 1962.
- [9] M.H.P. Bott, R.A. Smith, *The Estimation of the Limiting Depth of Gravitating Bodies.* Geophysical Prospecting, 6, (1958), 1-10.
- [10] M. Cuer, R. Bayer *Fortran Routines for Linear Inverse Problems.* Geophysics, 45, (1980), 1706-1719.

FIGURE CAPTIONS

Fig.1 A large region in the state of Bahia (Brazil) is considered between 40-44 degrees of longitude and 10-15 degrees of latitude. The iso-Bouguer anomaly lines are shown. The BB' direction is the one considered in the two-dimensional problem. The dotted rectangle is the region considered in the three-dimensional problem. In the three-dimensional problem measured data have been used in the AA'C'C rectangle, regularized data have been used in the remaining region.

Fig.2 The measured residual Bouguer anomaly in the BB' segment and the regularized one is shown.

Fig.3 The mass density distribution obtained from the data of Fig.2 solving the linear programming problem (3.3),(3.4) in a vertical section, 20Km deep, along the BB' direction is shown.

Fig.4 The mass density distribution obtained from the data of Table 1 solving the linear programming problem (3.2),(3.5) is shown.




fig. 1




fig. 2




fig. 3







fig. 4

§5. Apendix 2:

**COMPUTER SCIENCES
DEPARTMENT**

**University of Wisconsin -
Madison**



**SERIAL AND PARALLEL SOLUTION OF LARGE
SCALE LINEAR PROGRAMS BY AUGMENTED
LAGRANGIAN SUCCESSIVE OVERRELAXATION,**

by

R. De Leone and O. L. Mangasarian

Computer Sciences Technical Report #701

June 1987

Handwritten note:
This report is a presentation of the results of the
two-dimensional models, and
three-dimensional models.

**Serial and Parallel Solution of Large
Scale Linear Programs by Augmented
Lagrangian Successive Overrelaxation¹⁾**

R. De Leone & O. L. Mangasarian

Computer Sciences Department
University of Wisconsin
Madison, Wisconsin 53706

Technical Report #701

June 1987

Abstract. Serial and parallel successive overrelaxation (SOR) methods are proposed for the solution of the augmented Lagrangian formulation of the dual of a linear program. With the proposed serial version of the method we have solved linear programs with as many as 125,000 constraints and 500,000 variables in less than 72 hours on a MicroVax II. A parallel implementation of the method was carried out on a Sequent Balance 21000 multiprocessor with speedup efficiency of over 65% for problem sizes of up to 10,000 constraints, 40,000 variables and 1,400,000 nonzero matrix elements.

Key Words: Linear programming, SOR, augmented Lagrangian, parallel algorithms

Abbreviated Title: SOR Solution of Linear Programs

¹⁾This material is based on research supported by National Science Foundation Grants DCR-8420963 and DCR-8521228 and Air Force Office of Scientific Research Grants AFOSR-86-0172 and AFOSR-86-0255.

1. Introduction

In [8, 9, 10, 12,] successive overrelaxation methods are proposed for solving the dual of the problem of finding the least 2-norm solution of a linear program. This leads to an exterior penalty formulation of the dual of the original linear program with the interesting property that the penalty parameter need not approach zero in order to obtain an exact solution of the primal linear program [1, 14]. Thus the penalty parameter need only be less than a certain threshold value in order to obtain an exact solution to the primal linear program. However, the penalty parameter must approach zero in order to obtain a solution to the dual problem. Although this approach has been used effectively in conjunction with successive overrelaxation methods both on serial [10] and parallel machines [11, 12], we propose here the use of an augmented Lagrangian on the dual problem instead of an exterior penalty function in order to alleviate the twin difficulties of determining the threshold value of the penalty parameter required for an exact primal solution, and the need for the penalty parameter to approach zero in order to obtain a dual solution. The first proposal for using an augmented Lagrangian formulation for solving linear programs was made in [22]. In [18] Polyak and Tretiyakov made the remarkable discovery that after a finite number of steps of the augmented Lagrangian algorithm, an exact solution to the primal and dual linear programs is obtained. In [3] Golshtein proposed a projected Gauss-Seidel method in conjunction with an augmented Lagrangian formulation and gave computational results for linear programs with sizes up to 352 variables and 166 constraints. No convergence proofs of the projected Gauss-Seidel method was given in [3], nor of the closely related iterative method of Syrov and Churkreidze in [18]. We propose here the use of a projected successive overrelaxation method in conjunction with an augmented Lagrangian formulation. The convergence of the projected SOR method established in [7] is general enough to cover both a serial and a parallel implementation of the method. Since SOR methods are inherently serial in nature, their parallelization is not a routine matter. In [11, 13] two related methods were proposed for the parallelization of SOR methods. The more recent method [13] utilizes an unreduced relaxation factor interval of $(0, 2)$ which we shall employ here with an augmented Lagrangian algorithm for the dual linear program.

The paper is organized as follows. In Section 2 we give the necessary theoretical

background and convergence results for the proposed augmented Lagrangian method applied to the dual linear program. In Section 3 we describe a serial SOR implementation of the method and establish its convergence. In Section 4 we describe our parallel SOR implementation and in Section 5 we present computational results for both the serial and parallel methods.

We briefly describe our notation now. For a vector x in the n -dimensional real space R^n , x_+ will denote the vector in R^n with components $(x_+)_i = \max \{x_i, 0\}$, $i = 1, \dots, n$. The scalar product of two vectors x and y in R^n will be simply denoted by xy . For $1 \leq p \leq \infty$, the p -norm $\left(\sum_{i=1}^n |x_i|^p\right)^{1/p}$ of a vector in R^n will be denoted by $\|x\|_p$. For the 2-norm the subscript 2 will be dropped. R_+^n will denote the nonnegative orthant or the set of points in R^n with nonnegative components, while $R^{m \times n}$ will denote the set of all $m \times n$ real matrices. For $A \in R^{m \times n}$, A^T will denote the transpose, A_i will denote the i th row, $A_{i,j}$ the element in row i and column j , and for $I \subset \{1, \dots, m\}$, $J \subset \{1, \dots, n\}$, A_I will denote the submatrix of A with rows A_i , $i \in I$, while A_{IJ} will denote the submatrix of A with elements $A_{i,j}$, $i \in I$, $j \in J$. Similarly for $x \in R^n$ and $I_\ell \subset \{1, \dots, n\}$, x_{I_ℓ} will denote x_i , $i \in I_\ell$. The set $\{I_1, I_2, \dots, I_k\}$ is said to be a consecutive partition of $\{1, \dots, n\}$ if it is a partition of $\{1, \dots, n\}$ such that $i < j$ for $i \in I_\ell$, $j \in I_{\ell+1}$ and $\ell = 1, \dots, k-1$. Here and throughout the symbols $:=$ and \equiv denote definition of the term on the left and right sides of each symbol respectively.

2. Theoretical Background

We consider the linear program

$$(2.1) \quad \min cx \quad \text{subject to } Ax \geq b, x \geq 0$$

where $c \in R^n$, $b \in R^m$ and $A \in R^{m \times n}$ and its dual

$$(2.2) \quad \max_{(u,v) \geq 0} bu \quad \text{subject to } v = -A^T u + c \geq 0$$

For simplicity we exclude trivial constraints with $A_i = 0$. In [8, 9] the exterior penalty problem associated with the dual problem (2.2)

$$(2.3) \quad \max_{(u,v) \geq 0} \varepsilon bu - \frac{1}{2} \|A^T u + v - c\|^2$$

was solved by an SOR procedure for a sufficiently small value of the penalty parameter ε to obtain $(u(\varepsilon), v(\varepsilon))$. The unique least 2-norm solution \bar{x} of the linear program (2.1) was obtained by using the equation

$$(2.4) \quad x(\varepsilon) = \frac{1}{\varepsilon} (A^T u(\varepsilon) + v(\varepsilon) - c)$$

which relates an optimal solution $(u(\varepsilon), v(\varepsilon))$ of the dual penalty problem (2.3) and the unique solution $x(\varepsilon)$ of the corresponding quadratic primal problem [5]

$$(2.5) \quad \min cx + \frac{\varepsilon}{2} xx \quad \text{subject to } Ax \geq b, x \geq 0$$

In particular it follows [14] that the least 2-norm solution \bar{x} of the linear program (2.1) is related to $x(\varepsilon)$ of (2.4) by

$$(2.6) \quad \bar{x} = x(\varepsilon) \quad \text{for } \varepsilon \in (0, \bar{\varepsilon}] \quad \text{for some } \bar{\varepsilon} > 0$$

Thus the penalty parameter ε of the dual penalty problem (2.3) is the perturbation parameter of the perturbed primal problem (2.5). In order to avoid possible difficulties associated with determining the threshold value $\bar{\varepsilon}$ we consider instead of the exterior penalty problem (2.3) the augmented Lagrangian associated with the dual linear program (2.2)

$$(2.7) \quad L(u, v, x, \gamma) := bu - \frac{1}{2\gamma} \|A^T u + v - c\|^2 - x(A^T u + v - c)$$

It is a standard result [6, 2, 19] that for any $\gamma > 0$, a primal-dual solution $(\hat{x}, \hat{u}, \hat{v})$ of (2.1)-(2.2) is equivalent to a stationary point of the following saddlepoint problem of (2.7): Find an $(\hat{x}, \hat{u}, \hat{v}) \in R^n \times R_+^m \times R_+^n$ such that for all $x \in R^n$ and all $(u, v) \in R_+^m \times R_+^n$,

$$(2.8) \quad L(u, v, \hat{x}, \gamma) \leq L(\hat{u}, \hat{v}, \hat{x}, \gamma) \leq L(\hat{u}, \hat{v}, x, \gamma)$$

The standard augmented Lagrangian algorithm [2, 19] consists of a maximization step in the (u, v) space $R_+^m \times R_+^n$ followed by an unconstrained gradient descent step in the x space R^n . In particular we have the following.

2.1 Augmented Lagrangian Algorithm

Start with any $x^0 \in R^n$. Having x^i determine x^{i+1} as follows

$$(2.9) \quad \begin{cases} \text{(a) } L(u^i, v^i, x^i, \gamma^i) = \max_{(u,v) \geq 0} L(u, v, x^i, \gamma^i) \\ \text{(b) } x^{i+1} = x^i - \frac{1}{\gamma^i} \nabla_x L(u^i, v^i, x^i, \gamma^i) = x^i + \frac{1}{\gamma^i} (A^T u^i + v^i - c) \end{cases}$$

where $\{\gamma^i\}$ is any bounded sequence of positive numbers.

For this iterative linear programming algorithm Polyak and Tretiyakov [18] have given the following important finite convergence theorem.

2.2 Augmented Lagrangian Algorithm Finite Termination Theorem [18]

For any bounded positive sequence $\{\gamma^i\}$ and $x^0 \in R^n$, Algorithm 2.1 is finite, that is there exists an integer k such that (x^k, u^k, v^k) solve the dual linear programs (2.1)-(2.2). Furthermore for each x^0 there exists a $\hat{\gamma} > 0$ such that for $0 < \gamma^0 < \hat{\gamma}$, the method will terminate in one step, that is (x^1, u^1, v^1) will solve the dual linear programs (2.1)-(2.2).

Note that in the above theorem, two exact maximizations over the (u, v) space $R_+^m \times R_+^n$ are required in order to obtain $(x^{i+1}, u^{i+1}, v^{i+1})$ from x^i .

We note that since by duality theory [5]

$$(2.10) \quad \max_{(u,v) \geq 0} L(u, v, x, \gamma) = \min_x \left\{ cz + \frac{\gamma}{2} \|z - x\|^2 \mid Az \geq b, z \geq 0 \right\} =: \phi(x)$$

the Augmented Lagrangian Algorithm 2.1 is equivalent to the following gradient method applied to the proximal point function $\phi(x)$

$$(2.11) \quad \begin{cases} x^{i+1} = x^i - \frac{1}{\gamma^i} \nabla \phi(x^i) = \text{Prox}(x^i) \\ \text{where } \text{Prox}(x^i) \text{ is the solution of} \\ \min_x \left\{ cz + \frac{\gamma^i}{2} \|z - x^i\|^2 \mid Az \geq b, z \geq 0 \right\} \end{cases}$$

Bertsekas [1] and Rockafellar [21] also obtain finite termination for (2.11) from proximal point theory considerations for the linear programming case.

With this background we are prepared now to state and prove the convergence of our serial and parallel SOR algorithms.

3. Serial Successive Overrelaxation Algorithm

The proposed serial algorithm consists of applying the projected SOR method of [7] to the maximization step (2.9a) of the Augmented Lagrangian Algorithm 2.1 for a decreasing sequence of positive numbers $\{\gamma^i\}$. It follows from the Finite Termination Theorem 2.2 and a theorem of Pang [16, Theorem 3.1], that for any x^i the projected SOR method will generate a sequence of points converging to an x^{i+1} that solves the primal linear program (2.1), provided that γ^i is sufficiently small. There are no easily implementable and theoretically justifiable ways of determining how to choose γ^i sufficiently small [20, †], however we shall prescribe some computationally effective ways for doing that.

The serial projected SOR method has been proposed [7, 9] for solving the quadratic minimization problem

$$(3.1) \quad \min_{z \geq 0} \theta(z) := \min_{z \geq 0} \frac{1}{2} z M z + q z$$

where $M \in R^{k \times k}$ is symmetric and positive semidefinite. This is precisely problem (2.9a) of Algorithm 2.1 if we make the identifications

$$(3.2) \quad M := \frac{1}{\gamma^i} \begin{bmatrix} A \\ I \end{bmatrix} \begin{bmatrix} A^T & I \end{bmatrix}, \quad q := \begin{bmatrix} A(x^i - \frac{c}{\gamma^i}) - b \\ x^i - \frac{c}{\gamma^i} \end{bmatrix}, \quad z := \begin{bmatrix} u \\ v \end{bmatrix}$$

The projected SOR algorithm consists of the following.

$$(3.3) \quad z_j^{t+1} = \left(z_j^t - \omega (\nabla^2 \theta(z^t))_{jj}^{-1} \nabla_{z_j} \theta(z_1^{t+1}, \dots, z_{j-1}^{t+1}, z_j^t, \dots, z_k^t) \right)_+ \\ \omega \in (0, 2), \quad j = 1, \dots, k$$

More specifically for $\theta(z) = \frac{1}{2} z M z + q z$ we have the following.

3.1 Serial SOR Algorithm for $\min_{z \geq 0} \frac{1}{2} z M z + q z$

Choose $z^0 \in R_+^k$, $\omega \in (0, 2)$. Having z^t compute z^{t+1} as follows

$$(3.4) \quad z_j^{t+1} = \left(z_j^t - \omega M_{jj}^{-1} \left(\sum_{\substack{\ell=1 \\ \text{for } j > 1}}^{j-1} M_{j\ell} z_\ell^{t+1} + \sum_{\ell=j}^k M_{j\ell} z_\ell^t + q_j \right) \right)_+ \\ j = 1, \dots, k$$

We are ready to state and establish the convergence of our augmented Lagrangian serial SOR algorithm.

3.2 Augmented Lagrangian Serial SOR Algorithm

Let $\{\gamma^i\} \downarrow \bar{\gamma}$ for some $\bar{\gamma} > 0$, let $\{\delta^i\} \downarrow 0$ and let $x^0 \in R^n$. Having x^i determine x^{i+1} as follows:

- (a) Apply the Serial SOR Algorithm 3.1 to solve (2.9a) with the identifications (3.2), and let $(u^i(t), v^i(t))$ be the t iterate of this SOR algorithm. Stop if for some $t = t^i$ the following inequality is satisfied.

$$(3.5) \quad \begin{aligned} & \left| u^i(t) \nabla_u L(u^i(t), v^i(t), x^i, \gamma^i) \right| + \left| v^i(t) \nabla_v L(u^i(t), v^i(t), x^i, \gamma^i) \right| \\ & + \left\| \left(\nabla_u L(u^i(t), v^i(t), x^i, \gamma^i) \right)_+ \right\| + \left\| \left(\nabla_v L(u^i(t), v^i(t), x^i, \gamma^i) \right)_+ \right\| \leq \delta^i \end{aligned}$$

- (b) Set $x^{i+1} = x^i(t^i)$ where

$$(3.6) \quad x^i(t) := x^i + \frac{1}{\gamma^i} (A^T u^i(t) + v^i(t) - c)$$

3.3 Augmented Lagrangian Serial SOR Convergence Theorem

Let $\{\gamma^i\} \downarrow \bar{\gamma} > 0$ be a sufficiently rapidly decreasing sequence of positive numbers and $\bar{\gamma}$ sufficiently small. Then either

- (a) For some integer k , the sequence $\{x^k(t)\}$ converges to an \bar{x}^k that solves the linear program (2.1), or
- (b) For each subsequence of $\{(x^i, u^i(t^i), v^i(t^i))\}$ converging to some $(\bar{x}, \bar{u}, \bar{v})$, the corresponding subsequence $\{x^{i+1} = x^i(t^i)\}$ converges to an \hat{x} such that \hat{x} solves the linear program (2.1). If $\hat{x} = \bar{x}$, then (\bar{u}, \bar{v}) solves the dual linear program (2.2).

Proof Either the inequality (3.5) of the algorithm is satisfied at each iteration i for some t^i or not. Accordingly we have the alternatives (b) and (a) below respectively.

- (a) For some iteration $i = k$ the inequality (3.5) is never satisfied. Hence by Pang's Theorem 3.1 [16], since $L(u, v, x^k, \gamma^k)$ is by duality theory bounded above for $(u, v) \geq 0$ by $cx + \gamma^k \|x - x^k\|^2$ for any $x \geq 0$ such that $Ax \geq b$, it follows that the sequence

$$(3.7) \quad \{x^k(t)\} := \left\{ x^k + \frac{A^T u^k(t) + v^k(t) - c}{\gamma^k} \right\}, \quad t = 0, 1, 2, \dots,$$

where t is the SOR iteration index, converges to a vector \bar{x}^k defined by

$$(3.8) \quad \bar{x}^k := x^k + \frac{A^T \bar{u}^k + \bar{v}^k - c}{\gamma^k}$$

for some (\bar{u}^k, \bar{v}^k) which solves $\max_{(u,v) \geq 0} L(u, v, x^k, \gamma^k)$. Note however that $\{u^k(t), v^k(t)\}$ need not converge to (\bar{u}^k, \bar{v}^k) . If γ^k is sufficiently small (and this is what is meant by requiring that $\{\gamma^i\}$ decreases sufficiently rapidly) it follows by Theorem 2.2 that $x^{k+1} = \bar{x}^k$, is a solution of the primal linear program (2.1). (Note that (\bar{u}^k, \bar{v}^k) need not be a solution of the dual linear program (2.2). To obtain such a dual optimal (\bar{u}^k, \bar{v}^k) we need to solve $\min_{(u,v) \geq 0} L(u, v, \bar{x}^k, \gamma^k)$ exactly. See Corollary 3.5 below.)

(b) If inequality (3.5) holds for each i for some t^i , then since $\{\delta^i\} \downarrow 0$ and $\{\gamma^i\} \downarrow \bar{\gamma} > 0$, we have that for any subsequence of $\{(x^i, u^i(t^i), v^i(t^i))\}$ converging to some $(\bar{x}, \bar{u}, \bar{v})$, the point (\bar{u}, \bar{v}) solves the problem $\min_{(u,v) \geq 0} L(u, v, \bar{x}, \bar{\gamma})$ (because $\{\delta^i\} \downarrow 0$), and moreover the corresponding subsequence $\{x^{i+1}\}$ defined by (3.6) and part (b) of Algorithm 3.2 converges to an \hat{x} defined by

$$(3.9) \quad \hat{x} := \bar{x} + \frac{A^T \bar{u} + \bar{v} - c}{\bar{\gamma}}$$

If $\bar{\gamma}$ is sufficiently small, then it follows by Theorem 2.2 that \hat{x} solves the linear program (2.1). If in addition $\hat{x} = \bar{x}$, then (\bar{u}, \bar{v}) is feasible for the dual linear program (2.2) and is also optimal because $b\bar{u} = c\hat{x}$. ■

It is useful to point out that when the sequence $\{(u^i(t), v^i(t))\}$ of Algorithm 3.2 is bounded then it has an accumulation point and by Theorem 2.1 [7], each such accumulation point satisfies the optimality conditions for (2.9a). Consequently for each i inequality (3.5) of Algorithm 3.2 is satisfied after a finite number of steps of part (a) of the algorithm. However by Theorem 2(ii) of [9] we have that if the linear program (2.1) satisfies the Slater constraint qualification, then the sequence $\{(u^i(t), v^i(t))\}$ is indeed bounded. Therefore we have the following.

3.4 Augmented Lagrangian Serial SOR Convergence Corollary

Let the linear program (2.1) satisfy a Slater constraint qualification, that is $Ax > b$ for some $x \geq 0$. Then Theorem 3.3 holds with outcome (b).

Another useful observation follows from the fact [18, Lemma 1], [19] that minimizing the augmented Lagrangian $L(u, v, x, \gamma)$ with an optimal value of x and any $\gamma > 0$ gives a solution to the dual linear program (2.2). Hence we have the following.

3.5 Dual LP SOR Solution Corollary

Under the assumptions of Theorem 3.3 a solution to the dual linear program (2.2) can be obtained for either outcome (a) or (b) of Theorem 3.3 by solving respectively

$$(a) \quad \min_{(u,v) \geq 0} L(u, v, \bar{x}^k, \gamma^k)$$

or

$$(b) \quad \min_{(u,v) \geq 0} L(u, v, \bar{x}, \bar{\gamma})$$

We note immediately that we have left open the procedure by which the sequence $\{\gamma^i\}$ is decreased. This is an inherent theoretical difficulty that arises when using inexact minimization in the subproblems of proximal point or augmented Lagrangian algorithms. Thus the approximate minimization criteria of [21, Criteria A, B, A', B'] are not implementable for our problem, while the assumptions of [2, Section 2.5] are not verifiable for our problem. Computationally we have overcome this difficulty by using the following scheme, often used for updating the penalty parameter in augmented Lagrangian algorithm

$$(3.10) \quad \gamma^{i+1} = \begin{cases} \gamma^i & \text{if } \|x^{i+1} - x^i\| \leq \mu \|x^i - x^{i-1}\|, 0 < \mu < 1 \\ \nu \gamma^i & \text{otherwise, } 0 < \nu < 1 \end{cases}$$

This scheme works effectively for the solution of very large sparse linear programs as our computational results indicate.

4. Parallel Successive Overrelaxation Algorithm

The key to our parallel SOR algorithm is the use of the parallel gradient projection SOR (GP-SOR) that we proposed in [13] for the solution of (3.1) and which we outline below. Partition the matrix M of (3.1) into r contiguous horizontal blocks as follows:

$$(4.1) \quad M =: \begin{bmatrix} M_{I_1} \\ M_{I_2} \\ \vdots \\ M_{I_r} \end{bmatrix}$$

where the blocks M_{I_j} correspond to the variables z_{I_j} , and $\{I_1, I_2, \dots, I_r\}$ is a consecutive partition of $\{1, 2, \dots, k\}$. Now partition M_{I_j} as follows

$$(4.2) \quad M_{I_j} =: [M_{I_j, I_j} \quad M_{I_j, \bar{I}_j}]$$

where \bar{I}_j is the complement of I_j in $\{1, 2, \dots, k\}$. Thus M_{I_j, I_j} is a principal square submatrix of M with elements M_{st} , $s \in I_j$ and $t \in I_j$. We further partition M_{I_j, I_j} into the sum of its strictly lower triangular part L_{I_j, I_j} , its diagonal part D_{I_j, I_j} , and its strictly upper triangle part U_{I_j, I_j} as follows

$$(4.3) \quad M_{I_j, I_j} =: L_{I_j, I_j} + D_{I_j, I_j} + U_{I_j, I_j}$$

Now define a block diagonal matrix K as follows

$$(4.4) \quad K =: \begin{bmatrix} L_{I_1, I_1} & & & \\ & L_{I_2, I_2} & & \\ & & \ddots & \\ & & & L_{I_r, I_r} \end{bmatrix}$$

where each L_{I_j, I_j} is the strictly lower triangular part of M_{I_j, I_j} . An SOR algorithm can now be performed for each row block I_j , $j = 1, \dots, r$, simultaneously, that is in parallel. Note that the block diagonal matrix K replaces the traditional strictly lower triangular matrix of the serial SOR. Specifically we have the following.

4.1 Parallel GP-SOR Algorithm for (3.1)

Let $\{I_1, I_2, \dots, I_r\}$ be a consecutive partition of $\{1, 2, \dots, k\}$, let E be a positive diagonal matrix in $R^{k \times k}$ and let $z^0 \geq 0$. For $i = 0, 1, 2, \dots$, do the following

Direction Generation Define the direction

$$(4.5) \quad d^i = p(z^i) - z^i = \begin{pmatrix} p_{I_1}(z^i) - z_{I_1}^i \\ \vdots \\ p_{I_r}(z^i) - z_{I_r}^i \end{pmatrix}$$

such that $p(z^i)$ satisfies

$$(4.6) \quad p_{I_j}(z^i) = \left(z_{I_j}^i - \omega E_{I_j, I_j} (M_{I_j} z^i + q_{I_j} + L_{I_j, I_j} (p_{I_j}(z^i) - z_{I_j}^i)) \right)_+, \quad j = 1, \dots, r$$

where $\omega > 0$ is chosen such that for some $\nu > 0$

$$(4.7) \quad z_{I_j} \left((\omega E_{I_j, I_j})^{-1} + L_{I_j, I_j} \right) z_{I_j} \geq \nu \|z_{I_j}\|^2, \quad \forall z_{I_j}, \quad j = 1, \dots, r$$

Stop if $d^i = 0$, else continue.

Stepsize Generation $z^{i+1} = z^i + \lambda^i d^i$

where

$$(4.8) \quad f(z^i + \lambda^i d^i) = \min_{\lambda} \{ f(z^i + \lambda d^i) \mid z^i + \lambda d^i \geq 0 \}$$

4.2 Remark The principal part of this algorithm consists of the direction generation part (4.6), which can be performed in parallel on r processors. Once this is done the stepsize generation (4.8) is performed and the new value z^{i+1} is shared between the r processors.

The following convergence results were derived in [13] for Algorithm 4.1.

4.3 Theorem (Convergence of the Parallel GP-SOR Algorithm) Let M be symmetric and positive semidefinite. Either the sequence $\{z^i\}$ generated by the Parallel GP-SOR Algorithm 4.1 terminates at a solution of (3.1) or each of its accumulation points solves (3.1).

4.4 Corollary (Parallel GP-SOR special cases) Condition 4.7 of Algorithm 4.1 holds under either of the following two assumptions:

$$(4.9) \quad 0 < \omega < \min_{j=1, \dots, r} \min_{\substack{t \in I_j \\ t \neq i}} \frac{2}{E_{ii} \sum_{\substack{t \in I_j \\ t \neq i}} |M_{it}|}$$

$$(4.10) \quad 0 < \omega < 2, E = D^{-1} \text{ and } M \text{ is positive semidefinite}$$

Our parallel augmented Lagrangian method consists of replacing the Serial SOR Algorithm 3.1 by the Parallel GP-SOR Algorithm 4.1 in Algorithm 3.2 with option (4.10) for the choice of ω and E . We formally state the algorithm below.

4.5 Augmented Lagrangian Parallel SOR Algorithm

Identical to Algorithm 3.2 except that the Serial SOR Algorithm 3.1 in Algorithm 3.2 is replaced by the Parallel GP-SOR Algorithm 4.1 with condition (4.10) above in force.

5. Computational Results

Our algorithms were tested on random linear programs which were generated as follows. First the matrix A of the linear program (2.1) was generated. Each of its nonzero elements was a random number generated by a uniform distribution on the interval $[-100, 100]$. The number of nonzero elements in each row was in accordance to a prescribed density and the random position of each nonzero element was determined according to a uniform distribution on the column indices of the matrix. Next a primal-dual solution vector (\bar{x}, \bar{u}) was generated from a uniform distribution on the interval $[0, 10]$ with 80% of the components being nonzero. Finally the vectors b and c were chosen so that (\bar{x}, \bar{u}) is optimal.

In Figure 1 we give a summary of our computational results for 6 problems with the number of constraints varying between 25,000 and 125,000 and the number of variables varying between 100,000 and 500,000. All tests were performed on a MicroVax II with 16 megabytes of memory and an expanded disk swap space. We are not aware of any other linear programming software that can handle problems of the size that we have attempted on a comparable machine. MINOS [15], a state-of-the-art pivotal linear programming package, cannot handle any of the problems listed in Figure 1 because they are too big for the machine memory using the MINOS configuration. The largest problem attempted on the MicroVax II with MINOS was a problem with 5000 variables of 20,000 constraints and a matrix density of 0.2% with about 200,000 nonzero elements. MINOS was used with the standard partial pricing and scaling options. After 3150 iterations and 49 hours 54 minutes of machine time, the point was infeasible and the objective function was in error by 59% of the exact minimum. By comparison our Algorithm 3.2 solved the same problem in 1 hour and 4 seconds with a primal-dual objective function accuracy of 7 figures, and relative accuracy of not less than 10^{-9} as defined in Figure 1.

The Parallel SOR Algorithm 4.5 was implemented on the Sequent Balance 21000, a multiprocessor that incorporates eight NS32032 processors running at 10MHz, each with a floating point unit, memory management and an 8-kbyte cache sharing a global memory via a 32-bit wide pipelined bus. The machine has 8-Mbytes of physical memory. The operating system DYNIX, is a version of Berkeley 4.2 bsd unix. The computational results are depicted in Figures 2 and 3.

Figure 2 shows the total computing time versus number of processors for four different densities: $d=1, 2, 7$ and 10 percent for a linear program with 1000 constraints and 4000 variables. All problems were solved to a 7-figure accuracy of the primal-dual objective function and relative accuracy better than 10^{-7} as defined in Figure 1. We observe that the optimal number of processors, that is the one that solves the problem in minimum total time, increases with density as expected. This number is 3 for 1% and 2% densities, 6 for 7% density, and 7 or more for 10% density. This means that for denser problems, a larger number of processors is needed in order to arrive at the shortest solution time. This also means that for denser problems, the communication cost does not become a dominant and hence prohibitive factor until a larger number of processors are used.

In Figure 3 we show results for the case with 10,000 constraints and 40,000 variables, with density of 0.35% and about 1,400,000 nonzero elements. To our knowledge this is one of the largest linear programs solved on this relatively modest sized multiprocessor. One of the reasons that we were able to solve larger problems on the MicroVax II, is that the latter had twice the total memory size of the Balance 21000 and furthermore, the MicroVax was essentially a single-user machine dedicated to the serial SOR algorithm. The optimal number of processors for the low-density case shown in Figure 3 is 4. Just as in the cases of Figure 2, the optimal number of processors should increase with problem density.

We conclude with some observations on the speedup efficiency of our Parallel SOR Algorithm 4.5. We define the speedup efficiency $E(r)$ as the ratio of the actual to the theoretical speedup of the algorithm using r processors instead of 1 processor, thus

$$(5.1) \quad E(r) := \frac{T(1)}{rT(r)}$$

where $T(r)$ is the total time for solving a given problem using r parallel processors. Figure 4 shows the speedup efficiencies for a typical case of a linear program with 1000 constraints, 4000 variables and 2% density. The reason why some efficiencies are over 100% was pointed out in [13]. The explanation is that our Parallel SOR Algorithm 4.5 changes with the number of processors used, because the matrix K defined by (4.4) changes with the number of blocks into which M is divided. Thus we are not comparing identical algorithms when we evaluate the ration $T(1)/rT(r)$ of (5.1). Nevertheless the expression is a valid measure of efficiency in the sense of comparing the theoretical reduced time $T(1)/r$ to the observed time $T(r)$ for an algorithm with a variable K that depends on the partition

of M . If the matrix K is held fixed for $r = 1$ and $r > 1$, then we obtain efficiencies for identical algorithms, and they would all be less than 100%. This was demonstrated in [13]. Nevertheless the present efficiencies of over 100% are indeed very encouraging and also give the additional and somewhat surprising result that a serial implementation of our r -block Parallel SOR Algorithm 4.5 on a single machine, will give for some r a better computing time than the single block Serial SOR Algorithm 3.2. For the specific case of Figure 4, a serial implementation of the r -block parallel SOR with $r=2, 3, 4$ and 5 will be faster than the single block serial SOR.

Prob. No.	No. constr. $m \times 10^{-3}$	No. var. $n \times 10^{-3}$	Nonzero elements per row	Total Nonzero elements		Time hr:min	Fig. Accur. Obj. Func.		Log ₁₀ Rel. Accur.			
				$\times 10^{-5}$	Iter		P	D	1	2	3	4
							(a)	(b)	(c)	(d)	(e)	(f)
1	25	100	20	5	309	2:31	7	7	-7	-10	-9	-10
2	30	120	25	7.5	392	4:34	7	7	-10	-14	-12	-14
3	40	160	20	8	363	4:56	7	7	-9	-11	-7	-8
4	50	200	16	8	525	7:47	7	7	-9	-8	-9	-8
5	100	400	12	12	967	22:35	5	7	-3	-5	-5	-7
6	125	500	9	11.25	3100	71:40	6	7	-3	-5	-5	-7

(a) Number of correct figures in primal objective

(b) Number of correct figures in dual objective

(c) $\|(-Ax + b)_+\|_\infty / \|(b)_+\|_\infty$ (Relative Accuracy)

(d) $\|(A^T u - c)_+\|_\infty / \|(-c)_+\|_\infty$ (Relative Accuracy)

(e) $|cx - c\bar{x}| / |cx + c\bar{x}|$, \bar{x} : exact, x : computed (Relative Accuracy)

(f) $|bu - b\bar{u}| / |bu + b\bar{u}|$, \bar{u} : exact, u : computed (Relative Accuracy)

Fig. 1. MICROVAX II: Serial SOR Algorithm 3.2 test results.




Fig. 2. BALANCE 21000: Total time for Parallel SOR Algorithm 4.5 to solve linear program versus number of processors for various densities d . (Average of 4 randomly generated cases with 1000 constraints and 4000 variables.)




Fig. 3. BALANCE 21000: Total time for Parallel SOR Algorithm 4.5 to solve linear program versus number of processors. (d =density, m =number of constraints, n =number of variables.)

Density $d = 2\%$, $m = 1000$, $n = 4000$		
No. Processes r	Time Sec. $T(r)$	Speedup Efficiency $E(r) = T(1)/rT(r)$
1	3946	—
2	709	278%
3	638	206%
4	716	138%
5	711	111%
6	840	78%
7	850	66%

Fig. 4. BALANCE 21000: Speedup efficiency $E(r)$ for the Parallel SOR Algorithm 4.5 for an LP with 1000 constraints and 4000 variables and 2% density.

References

1. D. P. Bertsekas: "Necessary and sufficient conditions for a penalty method to be exact", *Mathematical Programming* 9, 1975, 87-99.
2. D. P. Bertsekas: "Constrained optimization and Lagrange multiplier methods", Academic Press, New York 1982.
3. E. G. Golshtein: "An iterative linear programming algorithm based on the use of modified Lagrange function", US-USSR Seminar on Distribution and Production Problems of Industry and Enterprise, October 15-17, 1980, University of Maryland, College Park, Proceedings Published in College Park, Maryland 1982.
4. Y. Y. Lin & J. S. Pang: "Iterative methods for large convex quadratic programs: A survey", *SIAM Journal on Control and Optimization* 25, 1987, 383-411.
5. O. L. Mangasarian: "Nonlinear programming", McGraw-Hill, New York 1969.
6. O. L. Mangasarian: "Unconstrained Lagrangians in nonlinear programming", *SIAM Journal on Control* 13, 1975, 772-791.
7. O. L. Mangasarian: "Solution of symmetric linear complementarity problems by iterative methods", *Journal of Optimization Theory and Applications* 22, 1977, 465-485.
8. O. L. Mangasarian: "Iterative solution of linear programs", *SIAM Journal on Numerical Analysis* 18, 1981, 606-614.
9. O. L. Mangasarian: "Sparsity-preserving SOR algorithms for separable quadratic and linear programs", *Computers and Operations Research* 11, 1984, 105-112.
10. O. L. Mangasarian: "Normal solution of linear programs", *Mathematical Programming Study* 22, 1984, 206-216.
11. O. L. Mangasarian & R. De Leone: "Parallel successive overrelaxation methods for symmetric linear complementarity problems and linear programs", *Journal of Optimization Theory and Applications* 54(3), September 1988.
12. O. L. Mangasarian & R. De Leone: "Error bounds for strongly convex programs and (super)linearly convergent iterative schemes for the least 2-norm solution of linear programs", *Applied Mathematics and Optimization*, 1988.
13. O. L. Mangasarian & R. De Leone: "Parallel gradient projection successive overrelaxation for symmetric linear complementarity problems and linear programs", University of Wisconsin Computer Sciences Department Tech. Rept. 659, August 1987, submitted to *Annals of Operations Research*.
14. O. L. Mangasarian & R. R. Meyer: "Nonlinear perturbation of linear programs", *SIAM Journal on Control and Optimization* 17, 1979, 745-752.

15. B. A. Murtagh & M. A. Saunders: "MINOS 5.0 user's guide", Stanford University Technical Report SOL 83.20, December 1983.
16. J. S. Pang: "More results on the convergence of iterative methods for the symmetric linear complementarity problem", *Journal of Optimization Theory and Applications* 49, 1986, 107-134.
17. B. T. Polyak: "Introduction to optimization", Optimization Software, Inc., New York 1987 (translated from Russian).
18. B. T. Polyak & N. V. Tretyakov: "Concerning an iterative method for linear programming and its economic interpretation", *Economics and Mathematical Methods* 8(5) 1972, 740-751 (Russian).
19. R. T. Rockafellar: "A dual approach to solving nonlinear programming problems by unconstrained optimization", *Mathematical Programming* 5, 1973, 354-373.
20. R. T. Rockafellar: "Monotone operators and the proximal point algorithm", *SIAM Journal on Control and Optimization* 14, 1976, 877-898.
21. R. T. Rockafellar: "Augmented Lagrangians and applications of the proximal point algorithm in convex programming", *Mathematics of Operations Research* 1, 1976, 97-116.
22. Yu. P. Syrov & Sh. S. Churkreidze: "Questions on the optimization of inter-branch and inter-regional connections in the planning for the growth of a unified national-economic system", Institute of National Economy of Irkutsk, Irkutsk 1970 (Russian).

END

DATE

FILMED

10 - 88