

UNCLASSIFIED

20030128388

AD-A204 443

## REPORT DOCUMENTATION PAGE

(2)

|                                                                              |  |  |                                                                                   |  |  |
|------------------------------------------------------------------------------|--|--|-----------------------------------------------------------------------------------|--|--|
| 1a. SECURITY CLASSIFICATION AUTHORITY                                        |  |  | 1b. RESTRICTIVE MARKINGS                                                          |  |  |
| 1c. DECLASSIFICATION/DOWNGRADING SCHEDULE                                    |  |  | 3. DISTRIBUTION AVAILABILITY OF REPORT                                            |  |  |
| 4. PERFORMING ORGANIZATION REPORT NUMBER(S)                                  |  |  | 5. MONITORING ORGANIZATION REPORT NUMBER(S)                                       |  |  |
| 6a. NAME OF PERFORMING ORGANIZATION                                          |  |  | 7a. NAME OF MONITORING ORGANIZATION                                               |  |  |
| 6b. OFFICE SYMBOL (if applicable)                                            |  |  | 7b. ADDRESS (City, State, and ZIP Code)                                           |  |  |
| 8a. NAME OF FUNDING/SPONSORING ORGANIZATION                                  |  |  | 9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER                                   |  |  |
| 8b. OFFICE SYMBOL (if applicable)                                            |  |  | 10. SOURCE OF FUNDING NUMBERS                                                     |  |  |
| 8c. ADDRESS (City, State, and ZIP Code)                                      |  |  | PROGRAM ELEMENT NO.                                                               |  |  |
|                                                                              |  |  | PROJECT NO.                                                                       |  |  |
|                                                                              |  |  | TASK NO.                                                                          |  |  |
|                                                                              |  |  | WORK UNIT ACCESSION NO.                                                           |  |  |
| 11. TITLE (Include Security Classification)                                  |  |  |                                                                                   |  |  |
| 12. PERSONAL AUTHOR(S)                                                       |  |  |                                                                                   |  |  |
| 13a. TYPE OF REPORT                                                          |  |  |                                                                                   |  |  |
| 13b. TIME COVERED                                                            |  |  |                                                                                   |  |  |
| 14. DATE OF REPORT (Year, Month, Day)                                        |  |  |                                                                                   |  |  |
| 15. PAGE COUNT                                                               |  |  |                                                                                   |  |  |
| 16. SUPPLEMENTARY NOTATION                                                   |  |  |                                                                                   |  |  |
| 17. COSATI CODES                                                             |  |  | 18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) |  |  |
| FIELD                                                                        |  |  | GROUP                                                                             |  |  |
| SUB-GROUP                                                                    |  |  |                                                                                   |  |  |
| 19. ABSTRACT (Continue on reverse if necessary and identify by block number) |  |  |                                                                                   |  |  |
| 20. DISTRIBUTION/AVAILABILITY OF ABSTRACT                                    |  |  |                                                                                   |  |  |
| 21. ABSTRACT SECURITY CLASSIFICATION                                         |  |  |                                                                                   |  |  |
| 22a. NAME OF RESPONSIBLE INDIVIDUAL                                          |  |  |                                                                                   |  |  |
| 22b. TELEPHONE NUMBER                                                        |  |  |                                                                                   |  |  |
| 22c. OFFICE SYMBOL                                                           |  |  |                                                                                   |  |  |

DD FORM 1473, 84 M/R

82 APR edition may be used until exhausted.  
All other editions are obsolete.SECURITY CLASSIFICATION OF THIS PAGE  
UNCLASSIFIED

THE KEY INVOLVEMENT OF POLY (ADP-RIBSYLATION) IN DEFENSE AGAINST  
TOXIC AGENTS: MOLECULAR BIOLOGY STUDIES

Post Available Copy

## SUMMARY AND RESEARCH OBJECTIVES:

Poly (ADP-Rib) polymerase requires DNA for activity and it is significant that the catalytic activity of this enzyme is directly coordinated to the number of DNA strand breaks in DNA, both *in vitro* as well as *in vivo*. The poly (ADP-Ribosylation) modification of chromatin-associated proteins thus functions during various biological reactions involving DNA repair, replication).

The project initially had three aims. Some of these aims have been modified during the two years of the course of the project. Most recently, we have been mainly interested in how poly (ADP-ribosylation) helps us to protect cells from toxic agents which interact with DNA, by utilizing new genetic tools which have been developed in our laboratory. Specifically, we have been the first group to have reported the cloning of the cDNA and gene for this enzyme. Our aim during the last years has been mainly concerned with how one might manipulate the hyperexpression of the cloned gene for this enzyme in order to first make cells more resistant to toxic agents and secondly, to begin to understand the underlining mechanism by which ADP-Ribosylation alters chromatin around DNA strand breaks to help the cell to recover from such damage.

## STATUS OF THE RESEARCH

Overview of Results and Progress

Progress has been made in a number of related areas concerning the mechanism of the involvement of poly(ADP-ribosylation) in recovery of cells from toxic agents and secondly on the cloning, sequencing chromosomal localization and other aspects of the molecular biology of poly(ADP-Rib) polymerase.

|      |                   |
|------|-------------------|
| Dist | and/or<br>Special |
| A-1  |                   |

DETAILED REPORT1. Immunofractionation Characterization of Single Strand DNA Breaks  
Adjacent to Sites of Poly(ADP-Rib).

Smulson, M., Poly (ADP-Ribosylation) of Nucleosomal Chromatin: Electrophoretic and Immunological Methods, "Methods in Enzymology" 106, 933-943, 1984, Edited by F. Wold Academic Press

89 2 15 149

Using the method above, we analyzed the nucleosomal populations adjacent and distal to poly (ADP-Rib) for internal single strand breaks by using two-dimensional electrophoretic gels. Relevant to the current project was the observation that immunofractionated poly (ADP-Rib) oligonucleosomal DNA contains significant amounts of internal single-strand breaks compared with bulk chromatin.

Overview: This is the first direct data that demonstrates that poly (ADP-ribosylated) nucleosomes are adjacent to strand breaks.

## 2. Use of Immunofractionation on Anti-Poly (ADP-Rib) Antibody to Study DNA Repair

Thraves, P. J., Kasid, U., and Smulson, M. E., Selective Isolation of Domains of Chromatin Proximal to both the Carcinogen-induced DNA Damage and Poly (ADP-Ribosylation), Cancer Research, 45, 386-391 (1985).

We observed an enhancement (2-fold) in the specific retention of hyper-ADP-ribosylated, in vivo [ $^3\text{H}$ ]-TdR labelled chromatin, following treatment with MNU. In addition we noted that the incorporation of the chain terminator Ara-C was also enriched in the poly (ADP-ribosylated) strand breaks. Furthermore, the retention of these chromatin regions to the antibody column was due to the increased synthesis of longer polymer chains on the protein acceptors of these chromatin domains. This methodology thus offers a useful means of isolating the substance under analysis the dynamic domains of chromatin undergoing DNA synthesis and/or repair. Selected data from this paper are shown below.




Table 1

Association in vivo of [ $^3\text{H}$ ]Ara-C with ADP-ribosylated chromatin

HeLa S<sub>2</sub> cells were resuspended in fresh medium at a concentration of  $8 \times 10^6$ /ml and labeled with [ $^3\text{H}$ ]Ara-C (250  $\mu\text{Ci}/250 \mu\text{l}$ ) at 8  $\mu\text{Ci}/\text{ml}$  for 10 min. The cells were then harvested and nuclei purified, and micrococcal nuclease-soluble chromatin were prepared and immunofractionated as described in Chart 1.

| Fraction                 | [ $^{32}\text{P}$ ]Poly(ADP-Rib) (cpm) | [ $^3\text{H}$ ]Ara-C (cpm) | $A_{\text{fraction}}$ <sup>a</sup> | Specific activity of [ $^3\text{H}$ ]Ara-C |
|--------------------------|----------------------------------------|-----------------------------|------------------------------------|--------------------------------------------|
| Unfractionated chromatin | 1843                                   | 3061                        | 4.85                               | 631                                        |
| Unbound                  | 625 (34) <sup>b</sup>                  | 1903 (62)                   | 4.47 (92)                          | 425                                        |
| Bound                    | 1218 (66)                              | 1158 (38)                   | 0.38 (8)                           | 3047                                       |

<sup>a</sup> Calculated from a separate unlabeled column run, performed in parallel.



<sup>b</sup> Numbers in parentheses, percentage.

### 3. Pauses in Poly (ADP-Rib) Subsequent to DNA Strand Breaks Increase 3T3 Cell Transformation, and Oncogene Expression.

Kasid, U. N., Stefanik, D., Lubet, R. A., Dritschilo, A., and Smulson, M. E.; Relationship between DNA Strand Breaks and Inhibition of poly (ADP-ribosylation): Enhancement of Carcinogen-Induced Transformation, Carcinogenesis, 7, 327-330 (1986).

Inhibition of poly (ADP-Rib) of nuclear proteins increases the persistence of DNA strand breaks elicited by DNA damaging agents, and markedly increases SCE exchange reactions. Accordingly, it seemed logical to test, at the molecular level, whether these cellular events cause rearrangements or alterations of specific sequences such as oncogenes in DNA. We utilized transformation of BALB/3T3 cells as a selective system to obtain homogeneous samples of DNA after damaging cellular DNA by X-rays and/or inhibition of poly (ADP-Rib).

Inhibition of poly (ADP-Rib) by benzamide or 3AB for a brief period following DNA damage due to either X-ray or MNNG in BALB/3T3 cells significantly (3-30X) enhanced transformation frequency.



We established 14 transformed cell lines, after having been characterized for growth in soft agar and tumor induction in nude mice. No gross rearrangements of 7 representative oncogenes including the more frequently activated c-kis-ras, was observed (see above, right). DNA dot blot hybridization suggested a 2-4 fold amplification of c-Ha-ras gene in 3 transformed cell lines.

Kasid, U. N., Hough, C., Thraves, P., Dritschilo, A., and Smulson, M.E.; The Association of Human c-Ha-ras Sequences with Chromatin and Nuclear Proteins. *Biochem. Biophys. Res. Commun.*, **128**, 226-232, 1985.

Southern-blot hybridization of DNA isolated from various nucleosomal chromatin fractions to probes of either c-Ha-ras or the inactive B-globin gene yielded the following results: (1) c-Ha-ras sequences segregated predominantly in the nuclease-sensitive chromatin fractions. (2) Unlike the B-globin (inactive gene), C-Ha-ras associated chromatin lacked typical nucleosomal structure. In these preliminary studies we also immuno-fractionated nucleosomes on anti-poly(ADP-Rib) Sepharose to examine the association of this oncogene with sites of poly(ADP-Rib).

**Overview:** These past studies provide an approach to determine, at the molecular level (i.e. by use of anti-sense), whether inhibitors of poly. (ADP-Rib) polymerase or any other mechanism to reduce DNA repair alone, or in conjunction with damaging agents can cause alterations in oncogene sequences.

#### 4. Poly ADP-Ribosylation of an Oncogene Protein.

Baksi, K., Alkhatib, H., and Smulson, M. E.; *In Vivo* characterization of The Poly ADP-Ribosylation of SV 40 chromatin and Large T Antigen by Immunofractionation, *Expt'l. Cell Res.*, **172**, 110-123, 1987.

We have confirmed the poly(ADP-ribosylation) of large T antigen of SV40 by using antibodies to both large T antigen and poly(ADP-ribose) and consequently have begun to characterize how this post-translational nuclear modification of the viral protein modulates its biological functions. SV40 minichromosomal subpopulation containing replicative intermediate DNA was shown to have a significantly higher affinity for anti-poly(ADP-Rib)-Sepharose than viral chromatin fractions containing mature minichromosomal DNA. An anti-large T-Sepharose column was used to isolate T antigen from crude extracts by two different approaches: (1) large T antigen was labeled with [<sup>35</sup>S]methionine *in vivo* and the infected cell extract was immunofractionated to isolate large T antigen and (2) large T antigen from infected cell extracts was immunofractionated followed by immunostaining. Using these techniques, 1-10% of the total T antigen from infected cells was found to be poly(ADP-ribosylated). Minichromosome preparations *per se* were also subjected to immunofractionation on anti-large T-Sepharose. The high level of retention of poly(ADP-ribosylated) species of minichromosomes on this matrix suggested that this post-translational modification of viral chromatin may be related to those steps in viral replication and transcription under regulation by large T antigen. © 1987 Academic Press, Inc.




Fig. 3. The natural occurrence of poly(ADP-ribosylation) large T antigen of SV40 virus as determined by immunofractionation. [<sup>35</sup>S]Methionine-labeled large T antigen isolated on an anti-large T antibody column (Fig. 1) was applied to 6 ml of antipoly(ADP-Rib)-immunoglobulin G-Sepharose 4B as described in Table 1. Acid insoluble [<sup>35</sup>S]methionine incorporation was monitored in 50  $\mu$ l of each sample (●). The unbound and bound samples were concentrated and analyzed by SDS-PAGE. The insert shows the autoradiogram of the fractions. Unbound (lane 1); bound (lane 2).

Proc. Natl. Acad. Sci. USA  
Vol. 84, pp. 1224-1228, March 1987  
Biochemistry

## Cloning and expression of cDNA for human poly(ADP-ribose) polymerase

(DNA repair/COS cell transfection/cell cycle/DNA strand breaks)

HUSSEIN M. ALKHATIB\*, DEFENG CHEN\*, BARRY CHERNEY\*, KISHOR BHATIA\*, VICENTE NOTARIO\*,  
CHANDRAKANT GIRI†, GARY STEIN‡, ELIZABETH SLATTERY§§, ROBERT G. ROEDER§, AND MARK E. SMULSON\*||

\*Department of Biochemistry, Georgetown University Schools of Medicine and Dentistry, Washington, DC 20007; †Laboratory of Biochemistry and Molecular Biology, The Rockefeller University, New York, NY 10021; and ‡Division of Virology, Food and Drug Administration, Bethesda, MD 21205; and §Department of Biochemistry and Molecular Biology, University of Florida College of Medicine, Gainesville, FL 32610

Communicated by Arthur B. Pardee, November 10, 1986

**ABSTRACT** cDNAs encoding poly(ADP-ribose) polymerase from a human hepatoma  $\lambda$ gt11 cDNA library were isolated by immunological screening. One insert of 1.3 kilobases (kb) consistently hybridized on RNA gel blots to an mRNA species of 3.6–3.7 kb, which is consistent with the size of RNA necessary to code for the polymerase protein (116 kDa). This insert was subsequently used in both *in vitro* hybrid selection and hybrid-arrested translation studies. An mRNA species from HeLa cells of 3.6–3.7 kb was selected that was translated into a 116-kDa protein, which was selectively immunoprecipitated with anti-poly(ADP-ribose) polymerase. To confirm that the 1.3-kb insert from  $\lambda$ gt11 encodes for poly(ADP-ribose) polymerase, the insert was used to screen a 3- to 4-kb subset of a transformed human fibroblast cDNA library in the

Okayama–Berg vector. One of these vectors [pcD-p(ADPR)P; 3.6 kb] was tested in transient transfection experiments in COS cells. This cDNA insert contained the complete coding sequence for polymerase as indicated by the following criteria: (i) A 3-fold increase in *in vitro* activity was noted in extracts from transfected cells compared to mock or pSV2-CAT transfected cells. (ii) A 6-fold increase in polymerase activity in pcD-p(ADPR)P transfected cell extracts compared to controls was observed by “activity gel” analysis on gels of electrophoretically separated proteins at 116 kDa. (iii) A 10- to 15-fold increase in newly synthesized polymerase was detected by immunoprecipitation of labeled transfected cell extracts. Using pcD-p(ADPR)P as probe, it was observed that the level of poly(ADP-ribose) polymerase mRNA was elevated at 5 and 7 hr of S phase of the HeLa cell cycle, but was unaltered when artificial DNA strand breaks are introduced in HeLa cells by alkylating agents.

In the above work we showed that the full-length cDNA for poly(ADP-Rib) polymerase, in an appropriate expression vector, is capable of causing hyperexpression of this enzymatic activity in cells already possessing endogenous activity for the enzyme. For example, in the figure from this paper shown below, cells were transfected with the polymerase cDNA under transient transfection conditions and 48 hrs after the initial transfection the cells were isolated and analyzed for both activity and also concentration of poly(ADP-Rib) polymerase caused by the cloned cDNA.

In Panel B in the figure below the enzymatic activity of polymerase was assayed in extracts from Cos cells as well as other cells which had been transfected with control plasmids or mock transfected. In Lane 4 it is obvious that greatly enhanced enzymatic potential for poly ADP-ribosylation existed due to the cloned gene. This was followed up by immunoprecipitating the newly synthesized polymerase due to the cloned gene in the various extracts. Again, it is obvious in Lane 9 that considerably increased amounts of poly(ADP-Rib) polymerase are synthesized in these cells due to the expression vector.




FIG. 5. Transfection of COS cells with pcD-12 produces enhanced expression of polymerase activity and immunoprecipitable polymerase protein. COS cells ( $10^6$  cells) in duplicate flasks were treated in the presence or absence of plasmid DNA (25  $\mu$ g per 175-mm flask) for 4 hr at 37°C in the presence of DEAE-dextran as described (25). After 48 hr, the cells were rinsed, scraped from plates, sonicated, and assayed for

Best Available Copy

# cDNA sequence, protein structure, and chromosomal location of the human gene for poly(ADP-ribose) polymerase

(DNA binding protein/DNA-strand-break repair/chromosomes 1, 13, and 14/restriction-fragment-length polymorphism)

BARRY W. CHERNLY\*, O. WESLEY MCBRIDE†, DEFFENG CHEN\*, HUSSEIN ALKHATIB\*, KISHOR BHATIA\*, PRESTON HENSLEY\*, AND MARK E. SMULSON\*

\*Department of Biochemistry, Georgetown University Schools of Medicine and Dentistry, Washington, DC 20007, and †Laboratory of Biochemistry, National Cancer Institute, National Institutes of Health, Bethesda, MD 20892

Communicated by Roscoe Brady, August 18, 1987 (received for review July 1, 1987)

**ABSTRACT** Recently we described a full-length cDNA for the human nuclear enzyme poly(ADP-ribose) polymerase. Here, we report the chromosomal localization and partial map of the human gene for this enzyme as well as the complete coding sequence for this protein. The nucleotide sequence reveals a single 3042-base open reading frame encoding a protein with a predicted  $M_r$  of 113,135. A comparison of this deduced amino acid sequence with the amino acid sequence of three peptides derived from human poly(ADP-ribose) polymerase revealed a match of 27 amino acid residues. A computer-derived structural analysis of the enzyme and a search for similarities with other proteins confirmed that the polymerase belongs to a subfamily of DNA/NAD-binding proteins and DNA-repair proteins. Possible  $Zn^{2+}$ -binding "fingers," a nucleotide-binding fold, and a nuclear transport signal were noted. Additionally, chromosomal mapping has identified polymerase-hybridizing sequences on human chromosomes 1 (the active gene), 13, and 14 (processed pseudogenes). Using the polymerase cDNA as a probe, we also have detected several DNA restriction fragment length polymorphisms in normal humans.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

FIG. 2. Nucleotide sequence of the Ovarian-Berg pCD-ADP-ribose polymerase cDNA insert and the deduced amino acid sequence of the 113,135-Da protein. The protein contains sequences coding for three poly(ADP-ribose) polymerase peptides (underlined and numbered), in the 3' untranslated region, a putative mRNA processing signal (AATAAA) is underlined. In the 5' region two nucleotides correspond to the Kozak criteria for initiation are underlined (12).




FIG. 3. Genomic organization of poly(ADP-ribose) polymerase. The alignment of similar *EcoRI* sites present in the genomic and cDNA maps is illustrated together with cDNA probes used to order the *EcoRI* fragments. The approximate location in the genomic map for the *PstI* cDNA site is also shown. (Separate scales have been used to construct the genomic and cDNA maps.) B, Base.

Best Available Copy

Overview: These two studies (as supported by this as well as other sources) have culminated a three year program on the total sequencing of the cDNA for poly(ADP-Rib) polymerase as well as a formal study on the chromosome localization of the polymerase gene. We are thus now able to analyze the various functional domains (i.e., active site, DNA binding domain, etc.) of the protein. We plan to use this information to engineer important regions of the enzyme into various expression vectors such to manipulate, in cells, the various biological functions of poly(ADP-Rib) polymerase. Additionally, site-directed mutants will be useful for new experiments. The chromosomal localization of polymerase gene has allowed us to assign restriction sites to isolate the total active gene for the enzyme. The isolation of the gene may be very useful in future years in order to develop cells containing not only the cDNA sequences but also active gene sequences including introns.


#### 6. Preliminary Progress on Regulation of Expression of Polymerase Gene After DNA Strand Breakage.

Bhatia, K., Stein, G. S., Bustin, M., Fornace, A. J., Imaizumi, M., Breitman, T. R., Cherney, B. W., and Smulson, M. E.; Expression of The Poly(ADP-Ribose) Polymerase Gene: The Cell Cycle, Differentiation, and DNA Repair (Submitted), J. Biol. Chem. 1987

Having isolated the full-length cDNA for the polymerase, we have now evaluated the role of endogenously and exogenously induced DNA strand breaks on the transcriptional control of this enzyme. During cell replication and differentiation significant changes were found in mRNA levels for the polymerase. In a synchronized population of HeLa cells or in serum-stimulated WI-38 cells, steady state levels of the polymerase mRNA were highest at mid S and S-G<sub>2</sub> phases and negligible in early S phase. Transcription by *in vitro* nuclear run-off, showed a 4-fold increased level of newly synthesized RNA occurring in mid S phase. Unlike histones, the polymerase run-off transcription was independent of the continued replication of DNA during S phase. Similar to a number of growth related gene transcripts that are selectively degraded, the polymerase mRNA has an AUUUA motif in the 3' untranslated region. We conclude from this that mRNA pools for the polymerase are regulated both at the transcriptional and post-transcriptional levels.

During retinoic acid or DMSO-induced differentiation of HL 60 cells mRNA levels for the polymerase increased very early and remained high for up to 48 hours following which it decreased to pre-induced levels. None of these changes, however, were found to accompany the induction of exogenously caused DNA strand breaks as occasioned by either dimethylsulfate, X-irradiation, or UV-irradiation. It appears that in sharp contrast to the catalytic requirement of the polymerase, the induction of transcription of this gene may not be a strand break-dependent process.

Best Available Copy



## 7. Hyper-Expression of Polymerase Causes Increased DNA Repair.

Bhatia, K., Giri, C., Pommier, Y., Cherney, B., Dritschilo, A., Alkhatib, H., and Smulson, M.; Enhanced rate of DNA strand break repair by hyper expression of recombinant human poly(ADP-Rib) polymerase. (Submitted), 1988.

### ABSTRACT

The nuclear enzyme poly(ADP-ribose) polymerase has been implicated in modulating the repair process of mammalian cells exposed to various DNA insults. Studies of poly ADP-ribosylation *in vivo* have also supported a mechanism by which the generation of DNA strand breaks is a requirement to activate the enzyme. Inhibitors of the poly ADP-ribosylation reaction cause a reduction in the rejoining of DNA strand breaks in cells after DNA damage, the potentiation of the cytotoxicity of alkylating agents, an increase in the extent of sister chromatid exchange reactions as well as ionizing radiation induced cell transformation. Recently we described the first cloning of poly(ADP-ribose) polymerase cDNA from a human *gt-10* library. A 3.7-kb full length cDNA contained in the OLeyman-Berg expression vector was subsequently isolated and tested for transient expression in Cos cells. Results indicate that a 3 fold increase in specific activity of poly(ADP-ribose) polymerase is present over endogenous levels in transfected cells. The high level of expression thus provided for the first time a direct test for the hypothesis of an involvement of poly ADP ribosylation in DNA repair by a positive enhancement of enzyme levels rather than indirectly by the use only of inhibitors. Cos cells allowed to overexpress the polymerase were X-irradiated and subsequently allowed to repair. Residual single strand breaks in transfected cells were significantly lower at 15 mins following repair as compared to single strand breaks in mock transfected or non transfected cells. When Cos cells were transfected with plasmid containing a nonfunctional polymerase cDNA the enhanced repair capacity was abolished. Increased repair capacities of overexpressing Cos cells were also compromised in the presence of 3-aminobenzamide.

Best Available Copy

PUBLICATIONS DURING THE LAST 2 1/2 YEARS (11/85-2/88)

From this and other support in the laboratory

1. FULL PAPERS:

Thraves, P. J., Kasid, U., and Smulson, M. E.; Selective Isolation of Domains of Chromatin Proximal to both Carcinogen-Induced DNA Damage and Poly(Adenosine-Diphosphate-Ribosylation), Cancer Research, 45, 386-391 (1985).

Kasid, U. N., Hough, C., Thraves, P., Dritschilo, A., and Smulson, M. E.; The Association of Human c-Ha-ras Sequences with Chromatin and Nuclear Proteins, Biochem. Biophys. Res. Commun., 128, 226-232 (1985).

Kasid, U. N., Stefanik, D., Lubet, R. A., Dritschilo, A., and Smulson, M. E.; Relationship between DNA Strand Breaks and Inhibition of poly (ADP-ribosylation): Enhancement of Carcinogen-Induced Transformation, Carcinogenesis, 7, 327-330 (1986).

Baksi, K., Alkhatib, H., and Smulson, M. E.; *In Vivo* characterization of The Poly ADP-Ribosylation of SV 40 chromatin and Large T Antigen By Immunofractionation, Expt'l. Cell Res., 172, 110-123 (1987).

Alkhatib, H., Chen, D., Cherney, B., Bhatia, K., Notario, B., Slattery, E., Giri, C., Stein, G., Roeder, R., and Smulson, M.; Cloning and Expression of cDNA for Human Poly(ADP-Rib) Polymerase, Proc. Natl. Acad. Sci., 84, 1224-1228 (1987).

Cherney, B., McBride, O. W., Chen, D., Alkhatib, H., and Smulson, M.; DNA Sequence, Protein Structure and the Chromosomal Location of the Human Gene for Poly(ADP-Rib) Polymerase, Proc. Natl. Acad. Sci. USA, 84, 8370-8374 (1987).

PAPERS SUBMITTED:

Kasid, U. N., Prieto-Soto, A., Halligan, B., Liu L. F., Dritschilo, A., and Smulson, M. E.; Poly(ADP-ribosylation) of chromatin associated human topoisomerase I: *In Vitro* and *in vivo* analysis. (Submitted), 1988.

Best Available Copy

Bhatia, K., Giri, C., Pommier, Y., Cherney, B., Dritschilo, A., Alkhatib, H., and Smulson, M.; Enhanced rate of DNA strand break repair by hyper expression of recombinant human poly(ADP-Rib) polymerase. (Submitted), 1988.

Bhatia, K., Stein, G. S., Bustin, M., Fornace, A. J., Imaizumi, M., Breitman, T. R., Cherney, B. W., and Smulson, M. E.; Expression of The Poly(ADP-Ribose) Polymerase Gene: The Cell Cycle, Differentiation, and DNA Repair. (Submitted), 1988.

#### RECENT CHAPTERS IN BOOKS:

Smulson, M. Poly (ADP-Ribosylation) of Nucleosomal Chromatin: Electrophoretic and Immunological methods, "Methods in Enzymology 106, 933-943 (1984), Edited by F. Wold. Academic Press, New York.

Smulson, M., and Sugimura, T., "Overview on Poly ADP-Ribosylation, "Methods in Enzymology 106, (1984), Edited by F. Wold. Academic Press, New York.

Smulson, M. E., Hough, D., Kasid, U., Dritschilo, A., and Lubet, R. DNA Strand-Breaks and Poly (ADP-Ribosylated) Mediation of Transcriptionally Active Chromatin and Transforming Gene Stability, in ADP-Ribosylation of Proteins, Edited by F. R. Althaus, H. Hiltz, and S. Shall, Springer-Verlag, Berlin, Heidelberg, 1985.

Smulson, M., Alkhatib, H., Bhatia, K., Cien, D., Cherney, B., Notario, V., Tahourdin, C., Dritschilo, A., Hensley, P., Breitman, T., Stein, G., Pommier, Y., McBride, O. W., Bustin, M., and Giri, C. The Cloning of the cDNA and Gene for Human Poly(ADP-Ribose) Polymerase: Status on the Biological Function(s) Using Recombinant Probes, in Niacin, Nutrition, ADP-Ribosylation and Cancer, Edited by M. Jacobson and E. Jacobson, Springer-Verlag, Berlin, Heidelberg, New York, Tokyo, 1988, In Press.

Best Available Copy