

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0043

TECHNICAL REPORT No. 94

Quantum Mechanics of a Molecular System Adsorbed on a Dielectric Surface

by

H. G. Oh, H. R. Lee, Thomas F. George, C. I. Um, Y. M. Choi and W. H. Kahng

Prepared for Publication

in

Physical Review A

Departments of Chemistry and Physics
State University of New York at Buffalo
Buffalo, New York 14260

March 1989

Reproduction in whole or in part is permitted for any purpose of the
United States Government.

This document has been approved for public release and sale;
its distribution is unlimited.

DTIC
ELECTE
S 14 MAR 1989 D
Q E

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/89/TR-94			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (if applicable)		7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260		7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (if applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract N00014-86-K-0043	
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		10. SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO.		PROJECT NO.	TASK NO.
					WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Quantum Mechanics of a Molecular System Adsorbed on a Dielectric Surface					
12. PERSONAL AUTHOR(S) H.G. Oh, H.R. Lee, Thomas F. George, C. I. Um, Y. M. Choi and W. H. Kahng					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) March 1989	
				15. PAGE COUNT 28	
16. SUPPLEMENTARY NOTATION Prepared for publication in Physical Review A					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	DIELECTRIC SURFACE DAMPED HARMONIC OSCILLATOR		
			ADSORBED MOLECULAR SYSTEM SINUSOIDAL EXTERNAL FORCE		
			QUANTUM MECHANICS FEYNMAN PATH INTEGRAL		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) A molecular system adsorbed on a dielectric surface is modeled as a damped harmonic oscillator driven by a sinusoidal external force. The exact propagator and wavefunction through the Feynman path integral method are obtained. The second quantization of this system is carried out. Expectation values of several physical quantities are evaluated. The amplitude for transitions between harmonic oscillator states and damped driven oscillator states are obtained explicitly, and the result is applied to a two-level system.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT ☒ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT ☐ DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

Quantum Mechanics of a Molecular System Adsorbed on a Dielectric Surface

H. G. Oh *, H. R. Lee and Thomas F. George
Department of Physics and Astronomy
239 Fronczak Hall
State University of New York at Buffalo
Buffalo, New York 14260

C. I. Um, Y. M. Choi and W. H. Kahng
Department of Physics, College of Science
Korea University
Seoul 136-701, Korea

Abstract

A molecular system adsorbed on a dielectric surface is modeled as a damped harmonic oscillator driven by a sinusoidal external force. The exact propagator and wavefunction through the Feynman path integral method are obtained. The second-quantization of this system is carried out. Expectation values of several physical quantities are evaluated. The amplitude for transitions between harmonic oscillator states and damped driven oscillator states are obtained explicitly, and the result is applied to a two-level system.

PACS Nos: 33.10.Cs, 03.65.Fd

*Permanent Address: Department of Physics Education
College of Education
Chonnam National University
Kwangju 500, Korea

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

1. Introduction

The study of time-dependent oscillator systems has received a great deal of attention both in classical and quantum mechanical studies. In the usual quantum-mechanical treatment of these systems, a time-independent Hamiltonian is assumed, which must be obtained through the Legendre transformation of the Lagrangian function which describes the equation of motion for the system.⁽¹⁾ In this context, it is unlikely that all physical oscillator systems have conserved Hamiltonians. The Hamiltonian of a damped driven harmonic oscillator is a typical one with explicit time-dependence. Even though there have been extensive works⁽²⁾ on this system, this apparently simple system has not been exactly solved quantum mechanically, in contrast to the classical case. Furthermore, the physical interpretations of the Hamiltonian itself and of the time-dependent Schrödinger equation obtained by replacing the canonical momentum with $\frac{\hbar}{i} \frac{\partial}{\partial x}$ in the Kaldirola-Kanai-type Hamiltonian⁽³⁾ differ among researchers. Some⁽⁴⁾ claim that this Hamiltonian (accordingly the Schrödinger equation) describes a dissipative quantum system, while others⁽⁵⁾ object to such an interpretation. From the viewpoint of group theory, Cervero et al⁽⁶⁾ interpreted it to represent a variable frequency oscillator with minimum uncertainty.

The Feynman path integral formalism⁽⁷⁾ provides a general approach to quantum systems. Montroll⁽⁸⁾ transformed the original Feynman path integral into a Gaussian integral, and his method was applied by Cheng⁽⁹⁾ to evaluate the propagator for a forced harmonic system. Once the propagator suitable to a given system is obtained, the wavefunction of the system is easily calculated. However, even if the Schrödinger equation is solved through the path integral or some other method, there remains the problem of second-quantization. In general, the Hamiltonian for the damped driven harmonic

oscillator (DDHO) is not conserved and not identical to the total energy of the system, giving rise to much difficulty for developing a second-quantization formalism. However, second-quantization, if possible, is important in connection with the problem of finding invariants of a non-conserved system, and for obtaining the coherent states⁽¹⁰⁾ of DDHO.

On the other hand, the optical properties of molecules are greatly altered when they are adsorbed on or near a solid surface. During the past decade, many theoretical and experimental methods have been developed to investigate surface spectroscopy, surface-enhanced Raman scattering (SERS),⁽¹¹⁾ resonance fluorescence,⁽¹²⁾ photodissociation,⁽¹³⁾ etc. There exist, however, several criticisms and controversies regarding the electromagnetic theory.⁽¹⁴⁾ In order to better understand the surface phenomena mentioned above, one must study at the quantum mechanical level the dynamical behavior and optical response of the molecules adsorbed on a solid (metal or dielectric) surface.

In this work, we present a study of the quantum mechanical properties of a molecule adsorbed on or near a dielectric plane surface. We represent the molecule as a polarizable point particle subject to a time-dependent external electric field. We do not consider the nuclear vibrations in this work. In Sec. 2 the classical equation of motion of the induced molecular dipole is given, and this equation is transformed into that of a damped harmonic oscillator driven by an external force. In Sec. 3 the propagator and wavefunction are obtained through the Feynman path integral method. Energy expectation values and dipole moment matrix elements are expressed in Sec. 4. Sec. 5 is devoted to second quantization and calculation of the transition probability between the harmonic oscillator states and damped driven harmonic oscillator states. In Sec. 6 we discuss the results.

2. Classical Equation of Motion for the Dipole Moment

We consider a molecule adsorbed on a dielectric plane surface with complex dielectric constant ϵ and light beam incident from the vacuum side at an angle θ with respect to the z-axis normal to the surface (xy-plane). The normal modes of the incident field satisfy Maxwell's equations subject to appropriate boundary conditions. The wave vector $\vec{k}$ of the field, which is assumed to be independent of frequency, is decomposed into a two-dimensional component $\vec{k}_{\parallel}$ parallel to the surface and a z-component perpendicular to the surface. The electric field vector is polarized in two ways. The first is the s-polarized wave (s-wave) perpendicular to the incident plane defined by $\vec{k}$ and $\vec{z}$ (a vector along the z-direction). The second is the p-polarized wave (p-wave) in the incident plane. The local field $\vec{E}_\ell(\vec{r}_m, t)$ at the adsorbed molecule position $\vec{r}_m$ is expressed as

$$\vec{E}_\ell(\vec{r}_m, t) = \vec{E}_p(\vec{r}_m, t) + \vec{E}_s(\vec{r}_m, t) \quad (2-1)$$

where t indicates time. $\vec{E}_p(\vec{r}_m, t)$ represents the primary field, which consists of the field incident directly on the molecule and the field reflected from the solid surface when the molecule is absent, and $\vec{E}_p$ is calculated using the Fresnel formula.⁽¹⁵⁾ $\vec{E}_s(\vec{r}_m, t)$ is the secondary field which represents the field emitted by the induced molecular dipole and reflected back to it from the surface. The induced dipole moment $\vec{\mu}(t)$ is the sum of dipole moments induced by the primary and secondary fields. The classical equation of motion for $\vec{\mu}(t)$ is given by

$$\ddot{\vec{\mu}} + \gamma \dot{\vec{\mu}} + \omega_0^2 \vec{\mu} = \omega_0^2 \vec{\alpha} \cdot \vec{E}_\ell \quad (2-2)$$

The dots indicate time derivatives, ω_0 is the oscillation frequency of the dipole charge, γ_0 is a natural damping constant which measures the linewidth in the optical processes, and $\vec{\alpha}$ represent the effective molecular polarizability tensor of second rank. In Eq. (2-2) we neglected the magnetic effects on the dipole motion, and hereafter we set the permeability of the surface dielectric to be unity. At the molecular position (0,0,d) the primary field vector is calculated using the Fresnel formulas as

$$E_p(d,t) = E_0 \exp[-i(kd \cos\theta + \omega't)] \quad (2-3)$$

where the components of E_0 are given as

$$\begin{aligned} E_{0x} &= E_0 \left\{ 1 - f_1(\epsilon(\omega'), \theta) \right\} \cos\theta \\ E_{0y} &= E_0 \left\{ 1 + f_2(\epsilon(\omega'), \theta) \right\} \end{aligned} \quad (2-4)$$

$$E_{0z} = E_0 \left\{ 1 + f_1(\epsilon(\omega'), \theta) \right\} \sin\theta$$

$$\begin{aligned} f_1(\epsilon(\omega'), \theta) &= \frac{\epsilon(\omega') - [\epsilon(\omega') - \sin^2\theta]^{\frac{1}{2}}}{\epsilon(\omega') + [\epsilon(\omega') - \sin^2\theta]^{\frac{1}{2}}} \exp(2ikd\cos\theta) \\ f_2(\epsilon(\omega'), \theta) &= \frac{\cos\theta - [\epsilon(\omega') - \sin^2\theta]^{\frac{1}{2}}}{\cos\theta + [\epsilon(\omega') - \sin^2\theta]^{\frac{1}{2}}} \exp(2ikd\cos\theta) \end{aligned} \quad (2-5)$$

$$\epsilon(\omega') = \epsilon_1(\omega') + i\epsilon_2(\omega') \quad (2-6)$$

Here $k = |\vec{k}|$ is the magnitude of the wavevector, and ω' is the angular frequency of the incident field.

The secondary field at the molecule position is expressed as a function of the induced dipole moment, ⁽¹⁶⁾

$$\vec{E}_s(d, t) = \int_{-\infty}^t dt' \vec{G}_s[d; d; t-t'] \cdot \frac{d\vec{\mu}(t')}{dt'} \quad (2-7)$$

where $\vec{G}_s$ is the scattering part of the dyadic Green's function. ^(17,18)

Equation (2-7) represents the self-polarization effect of the induced dipole. The electric field emitted by the dipole at an earlier time t' interacts with the surface atom (or molecule), causing it to emit lights by polarization.

This emitted field by the surface atom (molecule) interacts with the adsorbed molecule at a later time $t > t'$, adding an extra induced dipole moment, which emits radiation that polarizes the surface atom (molecule) again, etc.

Through these self-polarization processes, the secondary field yields a change in the linewidth and oscillation frequency of the dipole. Efrima et al ⁽¹⁸⁾ used a Fourier transformation method to solve Eq. (2-7), which gives an integro-differential equation, and applied it to the Raman scattering problem of an adsorbed molecular system. The Fourier transform of the dyadic Green's function $\vec{G}_s(d; d; \omega')$ can be calculated only numerically, since it is given in complicate integral form. To avoid such complexity, we incorporate the effects of the secondary field into the linewidth and the oscillation frequency of the dipole moment. In this manner, we obtain a modified classical equation of motion for the dipole moment,

$$\ddot{\vec{\mu}} + \gamma \dot{\vec{\mu}} + \omega_m^2 \vec{\mu} = \omega_0^2 \vec{\alpha}(\omega') \cdot \vec{E}_p(d, t) \quad (2-8)$$

As mentioned above, the self-polarization effects are absorbed within the modified damping constant γ and modified frequency ω_m , whose explicit forms are⁽¹⁹⁾

$$\gamma = \gamma_0 + \gamma_s$$

$$\omega_m^2 = (\omega_0 + \Delta\omega_m)^2$$

$$\gamma_s = \frac{\omega_0^2}{\omega} \hat{n} \cdot \left\{ \vec{\alpha}(\omega') \cdot [\text{Im } \vec{G}_s(d; d; \omega') \cdot \hat{n}] \right\} \quad (2-9)$$

$$\Delta\omega_m = \omega_0 \left[1 - \hat{n} \cdot \left\{ \vec{\alpha}(\omega') \cdot [\text{Re } \vec{G}_s(d; d; \omega') \cdot \hat{n}] \right\} \right]^{1/2} - \omega_0,$$

where $\vec{G}(d; d; \omega')$ is the Fourier transform of $\vec{G}_s(d; d; t-t')$ and $\hat{n} = \vec{\mu}/|\vec{\mu}|$.

Let us now introduce a coordinate system in which the effective polarizability tensor $\vec{\alpha}$ is diagonal,

$$\vec{\alpha}(\omega') = \alpha_x(\omega') \hat{x}\hat{x} + \alpha_y(\omega') \hat{y}\hat{y} + \alpha_z(\omega') \hat{z}\hat{z} \quad (2-10)$$

where $\hat{x}$, $\hat{y}$ and $\hat{z}$ are unit vectors along the coordinate axes. We assume that the molecule is isotropic and its dipole moment is directed toward the positive z-direction. In this case we can set $\hat{n} = \hat{z}$, and thus only the z-component of the primary field, which is given by Eq. (2-3) through (2-6), is required in the equation of motion. Other molecular dipole moment configurations are similar to this case, so that our method does not lose generality. By this manner, Eq. (2-8) can be rewritten as

$$\ddot{\mu} + \gamma \dot{\mu} + \omega_m^2 \mu = f_3(\epsilon(\omega'), \theta) e^{-i\omega' t} \quad , \quad (2-11)$$

where

$$f_3(\epsilon(\omega'), \theta) = E_0 \alpha(\omega') \omega_0^2 \sin \theta \left\{ 1 + f_1(\epsilon(\omega'), \theta) \right\} e^{-ikd \cos \theta} \quad . \quad (2-12)$$

Equation (2-11) can be written in a more familiar form if we take just the real part of the external driving force. After some calculations, we obtain the classical equation of motion for the dipole moment of the adsorbed molecule as

$$\ddot{x} + \gamma \dot{x} + \omega_m^2 x = f(t)/m_0 \quad . \quad (2-13)$$

Here $Qx = \mu$, where Q is the average dipole charge, and m_0 is the mass of the molecule. The form of equation of motion (2-13) is the same as that of the usual damped driven harmonic oscillator, so that f is given as

$$f(t) = g(\omega', \theta) \cos(\omega' t - \phi_0) \quad , \quad (2-14-a)$$

$$g(\omega', \theta) = \frac{2E_0 m_0 \omega_0^2 \sigma(\omega') \sin \theta}{Q g_3(\omega', \theta)} \left\{ g_1^2(\omega', \theta) + g_2^2(\omega', \theta) \right\}^{1/2}$$

$$g_1(\omega', \theta) = \cos \theta \cos(kd \cos \theta) \left[|\epsilon(\omega')|^2 \cos \theta + R(\omega', \theta) \left\{ \epsilon_1(\omega') \cos \eta + \epsilon_2(\omega') \sin \eta \right\} \right] + R(\omega', \theta) \cos \theta \sin(kd \cos \theta) \left\{ \epsilon_1(\omega') \sin \eta - \epsilon_2(\omega') \cos \eta \right\}$$

$$g_2(\omega', \theta) = R(\omega', \theta) \cos \theta \cos(kd \cos \theta) \left\{ \epsilon_2(\omega') \cos \eta - \epsilon_1(\omega') \sin \eta \right\}$$

$$\begin{aligned}
& - R(\omega', \theta) \sin(kd \cos \theta) \left[R(\omega'_1, \theta) + \cos \theta \left\{ \epsilon_1(\omega') \cos \eta + \epsilon_2(\omega') \sin \eta \right\} \right] \\
g_3(\omega', \theta) &= R^2(\omega', \theta) + |\epsilon(\omega')|^2 \cos^2 \theta + 2R(\omega', \theta) \cos \theta \left\{ \epsilon_1(\omega') \cos \eta \right. \\
& \quad \left. + \epsilon_2(\omega') \sin \eta \right\} \quad (2-14-b)
\end{aligned}$$

$$\begin{aligned}
\phi_0 &= \tan^{-1} \left\{ g_2(\epsilon(\omega'), \theta) / g_1(\epsilon(\omega'), \theta) \right\} \\
\eta &= \frac{1}{2} \tan^{-1} \left\{ \epsilon_2(\omega') / [\epsilon_1(\omega') - \sin^2 \theta] \right\} \quad (2-14-c)
\end{aligned}$$

$$R(\omega', \theta) = \left\{ [\epsilon_1(\omega') - \sin^2 \theta]^2 + \epsilon_2(\omega') \right\}^{1/2}.$$

To avoid the complicated procedure of evaluating the wavefunction and other quantities involved, we have taken the real part of the driving force. The classical Lagrangian and Hamiltonian function corresponding to the equation of motion (2-13) are given, respectively, as

$$L(\dot{x}, x, t) = \left[\frac{1}{2} m_0 \dot{x}^2 - \frac{1}{2} m_0 \omega_m^2 x^2 + f(t)x \right] \exp[\gamma t] \quad (2-15)$$

$$H(x, p, t) = \left[\frac{p^2}{2m_0} e^{-2\gamma t} + \frac{1}{2} m_0 \omega_m^2 x^2 - f(t)x \right] \exp(\gamma t), \quad (2-16)$$

where p is the canonical momentum.

3. Propagator and Wavefunction

Recently Um et al⁽²⁰⁾ have treated a quantum damped driven harmonic oscillator with the path integral method.⁽⁷⁾ Following their formalism, we solve the Schrödinger wave equation

$$i\hbar \frac{\partial \psi(x,t)}{\partial t} = \hat{H}\psi(x,t) \quad (3-1)$$

where the Hamiltonian operator $\hat{H}$ is obtained by making the replacement $p \rightarrow -i\hbar \frac{\partial}{\partial x}$ in Eq. (2-16). Adopting a Gaussian form for the propagator K ,

$$K[x(t), t; x(0), 0] = A_0 \exp \left[-a(t)x^2 e^{\gamma t} - b(t)x e^{\gamma t/2} - c(t) \right] \quad (3-2)$$

which satisfies the wave equation

$$i\hbar \frac{\partial K}{\partial t} = \hat{H}K \quad (3-3)$$

we obtain three first-order differential equations for the coefficients in the propagator. These differential equations are readily solved by the boundary condition that K reduces to the propagator for a damped harmonic oscillator⁽²¹⁾ when there is no driving force. Through the above procedure we obtain an explicit form for K ,

$$K[x(t), t; x_0, 0] = \left[\frac{m_0 \omega e^{\gamma t/2}}{2\pi i \sin(\omega t)} \right]^{1/2} \exp \left[A_1(t)x^2 - A_2(t)x \right] \\ \times \exp \left[-A_3(t)x_0^2 + A_4(t)x_0 \right] \quad (3-4)$$

where $x_0 = x(0)$ and $\omega = [\omega_m^2 - \gamma^2/4]^{1/2}$ represents the reduced frequency, which will be assumed to be real throughout the present text. The coefficients in Eq. (3-4) are given as

$$A(\omega', \theta) = g(\omega', \theta) / b_1(\omega')$$

$$b_1(\omega') = \left[(\gamma\omega')^2 + \left\{ \omega^2 - \omega'^2 + \gamma^2/4 \right\}^2 \right]^{1/2}$$

$$\phi_1 = \tan^{-1} \left[\gamma\omega' / (\omega^2 - \omega'^2 + \gamma^2/4) \right]$$

$$A_1(\cdot) = \frac{m_0\omega}{2i\hbar} e^{\gamma t} \left\{ \frac{\gamma}{2\omega} - \cot(\omega t) \right\} \quad (3-5)$$

$$A_2(t) = \frac{A(\omega', \theta)\omega}{i\hbar} e^{\gamma t} \left[\left\{ \frac{\gamma}{2\omega} - \cot(\omega t) \right\} \cos(\omega' t - \phi_0 - \phi_1) - \frac{\omega'}{\omega} \sin(\omega' t - \phi_0 - \phi_1) \right]$$

$$A_3(t) = \frac{m_0\omega}{2i\hbar} \left\{ \frac{\gamma}{2\omega} + \cot(\omega t) \right\}$$

$$A_4(t) = - \frac{A(\omega', \theta)\omega}{i\hbar \sin(\omega t)} e^{\gamma t/2} \cos(\omega' t - \phi_0 - \phi_1) + \frac{m_0\omega x}{i\hbar \sin(\omega t)} e^{\gamma t/2}$$

The wavefunction, $\psi_n(x, t)$, of the dipole moment at time t is obtained by the formula

$$\psi_n(x, t) = \int_{-\infty}^{\infty} dx_0 K[x, t; x_0, 0] \psi_n(x_0, 0) \quad , \quad (3-6)$$

where $\psi_n(x_0, 0)$ is the wavefunction of a damped harmonic oscillator with the reduced frequency ω . Combining Eqs. (3-4) and (3-6), we obtain

$$\psi_n(x, t) = \left[\frac{D(t)}{2^n n! \sqrt{\pi}} \right]^{\frac{1}{2}} \exp \left[-i \left(n + \frac{1}{2} \right) \cot^{-1}(\xi(t)) - B_3(t) \right] \exp \left[-B_1(t)x^2 + B_2(t)x \right] \\ \times H_n[D(t)(x - E(t))] \quad , \quad (3-7)$$

where

$$\xi(t) = \frac{\gamma}{2\omega} + \cot(\omega t)$$

$$D(t) = \left[\frac{m_0 \omega e^{\gamma t}}{\hbar \zeta_1(t)} \right]^{\frac{1}{2}}$$

$$E(t) = \frac{A(\omega', \theta)}{m_0} \cos(\omega' t - \phi_0 - \phi_1)$$

$$B_1(t) = \frac{1}{2} D^2(t) [1 + i\zeta_2(t)]$$

$$B_2(t) = D^2(t) E(t) [1 + i\zeta_3(t)]$$

(3-8)

$$B_3(t) = \frac{1}{2} D^2(t) E^2(t) [1 + i\xi(t)]$$

$$\zeta_1(t) = \sin^2(\omega t) [1 + \xi^2(t)]$$

$$\zeta_2(t) = \xi(t) + \zeta_1(t) \left[\frac{\gamma}{\omega} - \xi(t) \right]$$

$$\zeta_3(t) = \zeta_2(t) - \zeta_1(t) \frac{\omega'}{\omega} \tan(\omega' t - \phi_0 - \phi_1) \quad .$$

From now on, we shall use the notation $\xi(t) = \hat{\xi}$, $\xi_1(t) = \xi_1 \dots D(t) = D$, etc. for simplicity whenever there is no ambiguity. In obtaining the wavefunction we have used the identity

$$\exp\left[\pm i n \cot^{-1} y\right] = \left[1 + y^2\right]^{-\frac{n}{2}} [y \pm i]^n \quad (3-9)$$

We find the following useful relations between the coefficients in the exponential terms of the wavefunction,

$$B_1 + B_1^* = D^2, \quad B_2 + B_2^* = 2D^2 E, \quad B_3 + B_3^* = D^2 E^2 \quad (3-10)$$

which are used in later calculations.

4. Energy Expectation Values and Matrix Elements of the Dipole Moment

Using the Lagrangian of Eq. (2-15), the mechanical energy operator of the system is easily obtained as

$$E_{op} = -\frac{\hbar^2}{2m_0} e^{-2\gamma t} \frac{\partial^2}{\partial x^2} + \frac{1}{2} m_0 \omega_m^2 x^2 - xf(t) \quad (4-1)$$

We note that above energy operator is not identical to the Hamiltonian $\hat{H}$ itself. $\hat{H}$ does not represent the total energy of the system, but rather is the generator of the motion of an energy-dissipative open system. ⁽²²⁾

Equation (4-1) is the same as Eq. (6-1) in Ref. 20 except for the $xf(t)$ term.

It is straightforward to calculate the matrix elements of the energy operator:

$$\begin{aligned}
\langle E_{op} \rangle_{m,n} = & \frac{m_0 \omega_m^2}{2D^2} \left[\sqrt{(n+1)(n+2)} \left(\frac{1}{2} - 2\lambda_1 B_1 \right) \delta_{m,n+2} + \sqrt{2(n+1)} \right. \\
& \times (DE + 2\lambda_1 \lambda_2 B_1 - D\lambda_3) \delta_{m,n+1} + \left\{ (n+\frac{1}{2}) + 2(2n+1)\lambda_1 \lambda_4 B_1 + D^2 E^2 - \lambda_1 \lambda_2^2 - 2D^2 E \lambda_3 \right\} \delta_{m,n} \\
& + \sqrt{2n} (DE - 2\lambda_1 \lambda_2 \lambda_4 - D\lambda_3) \delta_{m,n-1} + \sqrt{n(n-1)} \left(\frac{1}{2} - 2\lambda_1 \lambda_4^2 D^2 \right) \delta_{m,n-2} \Big] \\
& \times \exp \left[i(m-n) \cot^{-1}(\xi(t)) \right] , \tag{4-2}
\end{aligned}$$

where

$$\begin{aligned}
\lambda_1(t) = & \left[\frac{\omega}{\omega_m \zeta_1 D^2} \right]^2 , \quad \lambda_2(t) = D[B_2 - 2B_1 E] \\
\lambda_3(t) = & \frac{A(\omega', \theta) b_1(\omega')}{m_0 \omega_m^2} \cos(\omega' t - \phi_0) , \quad \lambda_4(t) = D^2 - B_1 . \tag{4-3}
\end{aligned}$$

Evidently, only the diagonal element $\langle E_{op} \rangle_{n,n}$ and four off-diagonal elements $\langle E_{op} \rangle_{n\pm 1,n}$ and $\langle E_{op} \rangle_{n\pm 2,n}$ have non-zero values. The matrix elements of the dipole moment are expressed as

$$\langle \mu \rangle_{m,n} = \frac{Q}{2D} \exp \left[i(m-n) \cot^{-1}(\xi) \right] \left[\sqrt{2(n+1)} \delta_{m,n+1} + \sqrt{2n} \delta_{m,n-1} + 2DE \delta_{m,n} \right] . \tag{4-4}$$

This equation contains the diagonal element, which in fact is closely related to the parity problem of the wavefunction, as will be explained later. However, the selection rule for the dipole transition between different states

is $\Delta m = \pm 1$. Combining Eqs. (3-9) and (4-2), we arrive at the following explicit form for the dipole matrix elements:

$$\langle \mu \rangle_{n+1,n} = Q \left[\frac{(n+1)\hbar e \gamma \tau}{2m_0 \omega} \right] \left[e^{i\omega t} + \frac{\gamma}{2\omega} \sin(\omega t) \right] = \langle \mu \rangle_{n,n+1}^*$$

$$\langle \mu \rangle_{n,n} = \frac{QA(\omega', \theta)}{m_0} \cos(\omega' t - \phi_0 - \phi_1) \quad (t \neq 0)$$

(4-5)

$$\langle \mu \rangle_{n,n} = 0 \quad (t = 0)$$

5. Second Quantization and Transition Probability

To obtain a second-quantization formalism for a molecular system adsorbed on a solid surface, we define two functions,

$$u(t) = \frac{\xi(t) + i}{D(t)\sqrt{2[1+\xi^2(t)]}} \quad (5-1)$$

$$\eta(t) = 2i\hbar u(t) B_1(t) ,$$

where $D(t)$ and $\xi(t)$ have already defined in Eq. (3-8) ($\eta(t)$ is not the same as the angle η in Eq. (2-14-b)). Using the relation of Eq. (3-10), we obtain

$$\eta(t) u^*(t) - u(t) \eta^*(t) = i\hbar . \quad (5-2)$$

Thus we define the time-dependent annihilation and creation operators as

$$a(t) = \frac{1}{i\hbar} \left[\eta(t)(x-E(t)) - u(t) \left\{ p + i\hbar[B_2 - 2B_1 E] \right\} \right]$$

$$a^\dagger(t) = \frac{1}{i\hbar} \left[u^*(t) \left\{ p + i\hbar[B_2 - 2B_1 E] \right\} - \eta^*(t)(x - E(t)) \right] , \quad (5-3)$$

where $B_1(t)$, $B_2(t)$ and $E(t)$ are defined in Eq. (3-8). We note that $B_2(t) - 2B_1(t) E(t)$ is pure imaginary. We shall hereafter use the notation of $u(t) = u$ and $\eta(t) = \eta$. It is obvious that the non-Hermitian operators $a^\dagger(t)$ and $a(t)$ satisfy the commutation relation

$$[a(t), a^\dagger(t)] = 1 . \quad (5-4)$$

By the definition of Eq. (5-3), we get

$$x = u^* a(t) + u a^\dagger(t) + E . \quad (5-5)$$

$$p = \eta^* a(t) + \eta a^\dagger(t) - i\hbar[B_2 - 2B_1 E] .$$

Obviously x and p satisfy the relation $[x, p] = i\hbar$. Now we set

$$|m(t)\rangle = \frac{1}{\sqrt{m!}} (a^\dagger(t))^m |0(t)\rangle \quad (5-6)$$

$$a(t) |0(t)\rangle = 0 . \quad (5-7)$$

Combining Eqs. (5-3) and Eq. (5-7), we obtain the time-dependent ground-state wave function

$$\psi_0(x, t) \equiv \langle x | 0(t) \rangle$$

$$= \left[\frac{m_0 \omega e^{\gamma t}}{\pi \hbar \zeta_1(t)} \frac{\xi(t) - i}{\xi(t) + i} \right] \exp \left[-B_1(t)x^2 + B_2(t)x - B_3(t) \right] \quad (5-8)$$

We note that $\psi_0(x,t)$ reduces to $\psi_0(x)$ at $t = 0$, which is the ground-state wavefunction of a simple harmonic oscillator. Combining Eqs. (4-1), (5-4) and (5-5), we find the second-quantized form of the energy Hamiltonian operator, respectively, to be

$$\begin{aligned} E_{op} = & \left[\frac{\eta^2}{2m_0} e^{-2\gamma t} + \frac{1}{2} m_0 \omega_m^2 u^2 \right] a^\dagger(t) a^\dagger(t) + \left[\frac{\eta^{*2}}{2m_0} e^{-2\gamma t} + \frac{1}{2} m_0 \omega_m^2 u^{*2} \right] a(t) a(t) \\ & + \left[\frac{|\eta|^2}{m_0} e^{-2\gamma t} + m_0 \omega_m^2 |u|^2 \right] (a^\dagger(t) a(t) + \frac{1}{2}) + \left[-\frac{i\hbar\eta}{m_0} (B_2 - 2B_1 E) e^{-2\gamma t} - uf \right. \\ & + \left. m_0 \omega_m^2 u E \right] a^\dagger(t) + \left[-\frac{i\hbar\eta^*}{m_0} (B_2 - 2B_1 E) e^{-2\gamma t} - u^* f + m_0 \omega_m^2 u^* E \right] a(t) \\ & - \frac{\hbar^2}{2m_0} (B_2 - 2B_1 E)^2 e^{-2\gamma t} + \frac{1}{2} m_0 \omega_m^2 E^2 - Ef \end{aligned} \quad (5-9)$$

$$\hat{H} = e^{\gamma t} E_{op} \quad (5-10)$$

Equation (5-9) yields the same energy operator expectation values listed in Eq. (4-2). It is now straightforward to obtain the second-quantized form of physical quantities such as position, momentum, dipole moment operator, etc.

Let us find the relations between the time-independent operators $a^\dagger(t)$, $a(t)$ and time-independent operators $a^\dagger, a$. The latter ones are simply defined for a simple harmonic oscillator as

$$a = \sqrt{\frac{m_0 \omega_0}{2\hbar}} \left[x + \frac{ip}{m_0 \omega_0} \right]$$

(5-11)

$$a^\dagger = \sqrt{\frac{m_0 \omega_0}{2\hbar}} \left[x - \frac{ip}{m_0 \omega_0} \right]$$

Substitution of the above definitions into Eq. (5-3) yields

$$a(t) = \beta_1(t)a + \beta_2(t)a^\dagger + \beta_3(t)$$

(5-12)

$$a^\dagger(t) = \beta_2^*(t)a + \beta_1^*(t)a^\dagger + \beta_3^*(t)$$

where the coefficients are given by

$$\beta_1(t) = \frac{\alpha_0 u(t)}{\sqrt{2}} \left[1 + \frac{2B_1(t)}{\alpha_0^2} \right]$$

$$\beta_2(t) = \frac{\alpha_0 u(t)}{\sqrt{2}} \left[-1 + \frac{2B_1(t)}{\alpha_0^2} \right]$$

(5-13)

$$\beta_3 = -u(t)B_2(t)$$

$$\alpha_0 \equiv \left[\frac{m_0 \omega_0}{\hbar} \right]^{1/2}$$

We now find an explicit expression for the transition probability between two states $|m(t)\rangle$ and $|n(0)\rangle$, where $|n(0)\rangle$ is a simple harmonic oscillator state given by

$$|n(0)\rangle = \frac{(a^\dagger)^n}{\sqrt{n!}} |0(0)\rangle$$

(5-14)

Here $|0(0)\rangle$ is the well-known ground state of the simple harmonic oscillator.

Using Eqs. (5-6), (5-12) and (5-14), we get

$$\begin{aligned} \langle m(t) | n(0) \rangle &= \frac{1}{\sqrt{m!n!}} \langle 0(t) | (a(t))^m (a^\dagger)^n | 0(0) \rangle \\ &= \frac{1}{\sqrt{m!n!}} \langle 0(t) | [\beta_1 a + \beta_2 a^\dagger + \beta_3]^m (a^\dagger)^n | 0(0) \rangle, \end{aligned} \quad (5-15)$$

where $|0(t)\rangle$ is the time-dependent ground state of the damped driven harmonic oscillator, given by Eq. (5-8). $\langle m(t) | n(0) \rangle$ represents the probability amplitude of finding the system in a state $|m(t)\rangle$ at time t , given that the system was in a state $|n(0)\rangle$ at time zero. To calculate the right side of Eq. (5-15), we need to rearrange the operators in normal order, which means that all the $a^\dagger$'s appear to the left of the a 's. The normal-ordered arrangement is carried out by introducing a normal-ordering operator⁽²³⁾ N such that

$$[\beta_1 a + \beta_2 a^\dagger + \beta_3]^m (a^\dagger)^n = N \left[\left(\alpha^* + \frac{\partial}{\partial \alpha} \right)^n \left\{ \beta_1 \alpha + \beta_2 \alpha^* + \beta_3 \right\}^m \right], \quad (5-16)$$

where α and α^* are complex c-numbers. With this method, we obtain

$$\begin{aligned} &[\beta_1 a + \beta_2 a^\dagger + \beta_3]^m (a^\dagger)^n \\ &= \sum_{k_1=0}^m \sum_{k_2=0}^{m-k_1} \sum_{k_3=0}^n \binom{m}{k_1} \binom{m-k_1}{k_2} \binom{n}{k_3} \frac{k_1!}{(k_1-k_3)!} \\ &\times (a^\dagger)^{n+k_2-k_3} (a)^{k_1-k_3} \beta_1^{k_1} \beta_2^{k_2} \beta_3^{m-k_1-k_2}. \end{aligned} \quad (5-17)$$

Evidently $a^{k_1-k_3}|0(0)\rangle$ vanishes for all k_3 's except that $k_3 = k_1$. We have used above the notation $\beta_1 = \beta_1(t)$, $\beta_2 = \beta_2(t)$ and $\beta_3 = \beta_3(t)$, which will be used hereafter. $\left[\begin{smallmatrix} m \\ k \end{smallmatrix} \right]$ means binomial distribution.

Substitution of Eq. (5-17) into Eq. (5-15) gives

$$\begin{aligned} \langle m(t) | n(0) \rangle = & \frac{1}{\sqrt{m!n!}} \sum_{k_1}^{[m,n]} \sum_{k_2}^{m-k_1} \left[\begin{smallmatrix} m \\ k_1 \end{smallmatrix} \right] \left[\begin{smallmatrix} m-k_1 \\ k_2 \end{smallmatrix} \right] \left[\begin{smallmatrix} n \\ k_1 \end{smallmatrix} \right] \beta_1^{k_1} \beta_2^{k_2} \beta_3^{m-k_1-k_2} \\ & \times \langle 0(t) | (a^+)^{n-k_1+k_2} | 0(0) \rangle, \end{aligned} \quad (5-18)$$

where $[m,n] = \min(m,n)$ implies the smaller of the two. Using Eqs. (5-8) and (5-14), we find

$$\begin{aligned} \langle 0(t) | (a^+)^{n-k_1+k_2} | 0(0) \rangle = & \left[\frac{\alpha_0}{\sqrt{\pi} 2^{n-k_1+k_2}} \right]^{1/2} \left[\frac{m\omega e^{\gamma t}}{\pi \hbar \zeta_1} \frac{\xi + i}{\xi - i} \right]^{1/2} \exp[-B_3^*] \\ & \times \int_{-\infty}^{\infty} dx \exp \left[-\left\{ B_1^* + \frac{\alpha_0^2}{2} \right\} x^2 + B_2^* x \right] H_{n-k_1+k_2}(\alpha_0 x). \end{aligned} \quad (5-19)$$

To integrate the right side of Eq. (5-19), we introduce a generating function for the Hermite polynomial $He(y)$ such that

$$He(y) = \left[\frac{\partial^l}{\partial s^l} \exp\{-s^2 + 2sy\} \right]_{s=0}. \quad (5-20)$$

Then we have

$$I \equiv \int_{-\infty}^{\infty} dx \exp \left[- \left(B_1^* + \frac{\alpha_0^2}{2} \right) x^2 + B_2^* x \right] H_{n-k_1+k_2}(\alpha_0 x) \\ - \left[\frac{d^\ell}{ds^\ell} I(s) \right]_{s=0} \quad (\ell = n-k_1+k_2) \quad (5-21)$$

where

$$I(s) = \frac{e^{-s^2}}{\alpha_0} \int_{-\infty}^{\infty} dy \exp \left[- \frac{\alpha_0^2 + 2B_1^*}{2\alpha_0^2} y^2 + \frac{2\alpha_0 s + B_2^*}{\alpha_0} y \right], \quad (5-22)$$

which is easily integrated. Substituting the result of integration into Eq. (5-21) and performing some calculations, we obtain the following result:

$$I = \left[\frac{2\pi}{\alpha_0^2 + 2B_1^*} \right]^{\frac{1}{2}} \exp \left[\frac{(B_2^*)^2}{2(\alpha_0^2 + 2B_1^*)} \right] \sum_{k_3=0}^{n-k_1+k_2} \binom{n-k_1+k_2}{k_3} \frac{k_3!}{(k_3/2)!} \left[\frac{2B_1^* - \alpha_0^2}{2B_1^* + \alpha_0^2} \right]^{k_3/2} \\ \times \left[\frac{2\alpha_0 B_2^*}{\alpha_0^2 + 2B_1^*} \right]^{n-k_1+k_2-k_3} \quad (k_3: \text{even}) \quad (5-23)$$

By Eqs. (5-18), (5-19) and (5-23), we obtain the probability amplitude as

$$\langle m(t) | n(0) \rangle = \frac{1}{\sqrt{2^n m! n!}} \left[\frac{2\alpha_0}{\alpha_0^2 + 2B_1^*} \right]^{\frac{1}{2}} \left[\frac{m_0 \omega e^{\gamma t}}{\hbar \zeta_1} \frac{\xi + i}{\xi - i} \right]^{\frac{1}{2}} \exp \left[-B_3^*(t) + \frac{(B_2^*)^2}{2(\alpha_0^2 + 2B_1^*)} \right]$$

$$\begin{aligned}
& \times \sum_{k_1=0}^{[m,n]} \sum_{k_2=0}^{m-k_1} \sum_{k_3=0}^{n-k_1+k_2} \begin{Bmatrix} m \\ k_1 \end{Bmatrix} \begin{Bmatrix} m-k_1 \\ k_2 \end{Bmatrix} \begin{Bmatrix} n \\ k_1 \end{Bmatrix} \begin{Bmatrix} n-k_1+k_2 \\ k_3 \end{Bmatrix} \frac{(k_3)!}{(k_3/2)!} \frac{\beta_1^{k_1} \beta_2^{k_2} \beta_3^{m-k_1-k_2}}{2^{(k_2-k_1)/2}} \\
& \times \left[\frac{2B_1^* - \alpha_0^2}{2B_1^* + \alpha_0^2} \right]^{k_3/2} \left[\frac{2\alpha_0 B_2^*}{2B_1^* + \alpha_0^2} \right]^{n-k_1+k_2-k_3} \quad (k_3: \text{even}) \quad (5-24)
\end{aligned}$$

The transition probability between the DDHO state $|m(t)\rangle$ and the simple harmonic oscillator state $|n(0)\rangle$ is given by

$$P_{m,n}(t) = |\langle m(t) | n(0) \rangle|^2 \quad (5-25)$$

6. Results and Discussion

We have modeled a molecule adsorbed on a dielectric solid surface as a damped harmonic oscillator driven by a time-dependent electric field, consisting of a primary and a secondary field. The Hamiltonian of this modeled system is not identical to the mechanical energy operator, showing a general characteristic of the Kalidola-Kanai-type Hamiltonian. The propagator of Eq. (3-4) has the same structure as that of Gerry⁽²⁴⁾ when there exists no driving force, and is similar to that of Dodonov et al.⁽⁴⁾ The wavefunction of Eq. (3-7) has no definite parity, which can be easily seen from the matrix elements of x and x^2 ,

$$\langle m(t) | x | n(t) \rangle = \sqrt{n+1} u(t) \delta_{m,n+1} + \sqrt{n} u^*(t) \delta_{m,n-1} + E(t) \delta_{m,n} \quad (6-1)$$

$$\langle m(t) | x^2 | n(t) \rangle = \sqrt{(n+1)(n+2)} u^2(t) \delta_{m,n+2} + \sqrt{n(n-1)} u^{*2}(t) \delta_{m,n-2}$$

$$\begin{aligned}
& + \left[(2n+1) |u(t)|^2 + E^2(t) \right] \delta_{m,n} \\
& + 2\sqrt{(n+1)} u(t) E(t) \delta_{m,n+1} + 2\sqrt{n} u^*(t) E(t) \delta_{m,n-1} \quad (6-2)
\end{aligned}$$

If $\psi_m(x,t)$ and $\psi_n(x,t)$ have definite parity and both have the same (even or odd) parity, $\langle m(t) | x | n(t) \rangle$ must be always zero, while if they have opposite parities, $\langle m(t) | x^2 | n(t) \rangle$ must vanish. Equations (6-1) and (6-2), however, do not satisfy this rule. This property of the wavefunctions does not result from the decay characteristic of the system, but from the interaction between the system and the driving force, because the wavefunction of a damped (not driven) harmonic oscillator has a definite parity.⁽²¹⁾ Due to the absence of definite parity of the wavefunction, the diagonal element of the dipole moment takes the form given in Eq. (4-5), in contrast to a simple harmonic oscillator. The real part of the off-diagonal elements of the dipole moment is the same as that of the classical case.⁽²⁵⁾

Because of the time-dependence $|\psi_n(x,t)|^2$, the expectation values of E_{op} (or $\hat{H}$) and other physical quantities do not stay constant in time. The diagonal parts of E_{op} and $\hat{H}$ are approximated as,

$$\langle E_{op} \rangle_{n,n} \approx (n + \frac{1}{2}) \hbar \omega e^{-\gamma t} + \frac{g^2(\omega, \theta)}{2m_0[\omega_m^2 - \omega'^2 + \gamma\omega']} \quad (6-3)$$

$$\langle \hat{H} \rangle_{n,n} = e^{\gamma t} \langle E_{op} \rangle_{n,n} \quad (6-4)$$

The first term in Eq. (6-3) shows the decay of the quantum state, and the second term represents the energy absorption from the external field; these processes are related to the lineshape in optical phenomena. When there

exists no driving force ($g(\omega', \theta) = 0$), Eqs. (6-3) and (6-4) reduce to those of a damped harmonic oscillator.⁽²⁶⁾ Hence, we can say that for a damped harmonic oscillator the expectation values of $\hat{H}$ stays constant in time and those of E_{op} do not, while for a damped driven oscillator the expectation of both operators vary in time. Of course, for the simple harmonic case, the two operators are identical and their expectation values remain stationary.

The method adopted in Sec. 5 to calculate $P_{m,n}(t)$ is different from that of others,^(20,27) and Eq. (5-24) is of a new form. In practice, however, it is a formidable task to get the transition probability, $P_{m,n}$, from Eq. (5-24). Even for the simplest two-level case such that $|m(t)\rangle = |1(\cdot)\rangle$ and $|n(0)\rangle = |0(0)\rangle$, the calculation is quite tedious. Hence, we simply write the result for this case:

$$\begin{aligned}
 P_{1,0} &= |\langle 1(t) | 0(0) \rangle|^2 \\
 &= 2\alpha_0^2 E^2(t) e^{\gamma t/2} \left[1 - e^{\gamma t} + 2e^{-2\gamma t} + (1 + e^{\gamma t}) \zeta_3^2(t) \right] \left[1 + e^{-\gamma t} \right]^{-3} \\
 &\times \exp \left[- \frac{\alpha_0^2 E(t)^2 \left\{ 1 + e^{\gamma t} \zeta_3^2(t) \right\}}{1 + e^{-\gamma t}} \right], \quad (6-5)
 \end{aligned}$$

where we have used the approximation $D \approx \alpha_0 \exp(\gamma t/2)$, since in a real system $\omega \approx \omega_0$ and $\gamma \ll \omega$.

Here we note that the transitions should occur on a time scale shorter than the system damping time⁽²³⁾ γ^{-1} , i.e., $\gamma t \ll 1$. Thus Eq. (6-5) can be approximated as

$$P_{1,0}(t) = I(\omega', t) \exp[-I(\omega', t)] \quad (6-6)$$

$$I(\omega', \tau) = \frac{\alpha_0^2 g^2(\omega', \theta) \left[\cos^2(\omega' \tau - \phi_0 - \phi_1) + \left(\frac{\omega'}{\omega}\right)^2 \sin^2(\omega' \tau - \phi_0 - \phi_1) \right]}{2m_0^2 \left[\left\{ \omega^2 - \omega'^2 + \gamma^2/4 \right\}^2 + (\gamma\omega')^2 \right]} \quad (6-7)$$

The transition probability depends strongly on the external driving force and the energy transfer mechanism. Around the resonance frequencies $\omega' \approx \omega^2 + \gamma^2/4 = \omega_m^2$, Eq. (6-9) becomes

$$I_R(\omega') = \frac{\alpha_0^2 g^2(\omega', \theta)}{2m_0^2} \left[\left\{ \omega^2 - \omega'^2 + \gamma^2/4 \right\}^2 + (\gamma\omega')^2 \right]^{-1} \quad (6-8)$$

where this form is related to the lineshape in optical phenomena. Application of the results in this work to optical phenomena on a dielectric surface, to finding the coherent states⁽²⁸⁾ and to the time-dependent invariant⁽²⁹⁾ problem will be left for further study.

Acknowledgments

This research was supported by the Office of Naval Research, the National Science Foundation under Grant CHE-8620274, and the Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract F49620-86-C-0009. Dr. Oh acknowledges support by the Korea Science and Engineering Foundation for an overseas fellowship. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon.

References

1. H. Goldstein, Classical Mechanics 2nd Ed. (Addison-Wesley, Reading, Massachusetts, 1980), Chapt. 1 and 8.
2. H. Dekker, Phys. Rev. 80 (1981) and references listed there.
3. (a) P. Kaldirola, Nuovo Cimento 18, 393 (1941);
(b) E. Kanai, Prog. Theor. Phys. 3, 440 (1948).
4. V. V. Dodonov and V. I. Mankov, Phys. Rev. A 20, 550 (1979).
5. (a) I. R. Senitzky, Phys. Rev. 119, 670 (1960);
(b) D. Greenberger, J. Math. Phys. 20, 962 (1979).
6. J. M. Cervero and J. Villarroel, J. Phys. A: Math. Gen. 17, 2963 (1984).
7. R. P. Feynman and A. R. Hibbs, Quantum Mechanics and Path Integrals (New York, McGraw Hill, New York, 1965), Chapt. 8.
8. E. W. Montroll, Commun. Pure. Appl. Math. 5, 415 (1952).
9. B. K. Cheng, J. Math. Phys. 25, 1804 (1984).
10. R. J. Glauber, Phys. Rev. 131, 2766 (1963).
11. (a) J. C. Tsang and J. R. Kirtley, Solid State Commun. 30, 617 (1979);
(b) I. A. Creighton, Surf. Sci. 124, 209 (1983).
12. H. F. Arnoldus and T. F. George, J. Phys. B: At. Mol. Opt. Phys. 21, 431 (1988).
13. P. T. Leung and T. F. George, J. Chem. Phys. 85, 4728 (1986).
14. (a) G. S. Agarwal, Phys. Rev. A 11, 230, 243, 253 (1975);
(b) M. Moskovits, Rev. Mod. Phys. 57, 788 (1985).
15. (a) M. Born and E. Wolf, Principles of Optics, 6th Ed. (Pergamon, New York, 1980), Chapt. I, # 1.5.
(b) J. D. Jackson, Classical Electrodynamics, 2nd Ed. (Wiley, New York, 1975), Chapt. 7, # 3.

- 2
16. (a) S. Efrima and H. Metiu, Chem. Phys. Lett. 60, 59 (1978);
(b) S. Efrima and H. Metiu, J. Chem. Phys. 70, 1602 (1979).
 17. Chen-To Tai, Dyadic Green's Function in Electrodynamical Theory (Intext Educational Publishers, Scranton, Pennsylvania, 1971), Chapt. 4.
 18. S. Efrima and H. Metiu, J. Chem. Phys. 70, 1939, 2297 (1979).
 19. J. I. Gersten and A. Nitzan, Surf. Sci. 158, 165 (1985).
 20. C. I. Um, K. H. Yeon and W. H. Kahng, J. Phys. A: Math. Gen. 20, 611 (1987).
 21. K. H. Yeon, C. I. Um and T. F. George, Phys. Rev. B 36, 5287 (1987).
 22. J. R. Brinati and S. S. Mizrahi, J. Math. Phys. 21, 2154 (1980).
 23. W. H. Louisell, Quantum Statistical Properties of Radiation (Wiley, New York, 1973), Chaps. 3 & 6.
 24. C. C. Gerry, J. Math. Phys. 25, 1820 (1984).
 25. W. Demtröder, Laser Spectroscopy, Springer Series in Chemical Physics 5 (Springer-Verlag, Berlin, 1981), Chapt. 3.
 26. R. W. Hasse, J. Math. Phys. 16, 2005 (1975).
 27. (a) L. F. Landovitz, A. M. Levin, E. Ozizmir and W. M. Schrieber, J. Chem. Phys. 78, 6133 (1983).
(b) L. F. Landovitz, A. M. Levine and W. M. Schreiber, J. Math. Phys. 21, 2159 (1980).
 28. (a) R. J. Glauber, Phys. Rev. 131, 2766 (1963);
(b) S. Howard and S. K. Roy, Am. J. Phys. 55, 1109 (1987).
 29. (a) H. R. Lewis, Jr. and W. B. Riesenfeld, J. Math. Phys. 10, 1458 (1969).
(b) R. S. Kaushal and J. J. Korsch, J. Math. Phys. 22, 1904 (1981).

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. C. B. Harris
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. H. Weaver
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. F. Kutzler
Department of Chemistry
Box 5055
Tennessee Technological University
Cookeville, Tennessee 38501

Dr. A. Reisman
Microelectronics Center of North Carolina
Research Triangle Park, North Carolina
27709

Dr. D. DiLella
Chemistry Department
George Washington University
Washington D.C. 20052

Dr. M. Grunze
Laboratory for Surface Science and
Technology
University of Maine
Orono, Maine 04469

Dr. R. Reeves
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. J. Butler
Naval Research Laboratory
Code 6115
Washington D.C. 20375-5000

Dr. Steven M. George
Stanford University
Department of Chemistry
Stanford, CA 94305

Dr. L. Interante
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Mark Johnson
Yale University
Department of Chemistry
New Haven, CT 06511-8118

Dr. Irvin Heard
Chemistry and Physics Department
Lincoln University
Lincoln University, Pennsylvania 19352

Dr. W. Knauer
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. K.J. Klaubunde
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. Murday
Naval Research Laboratory
Code 6170
Washington, D.C. 20375-5000

Dr. J. B. Hudson
Materials Division
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Theodore E. Madey
Surface Chemistry Section
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

Dr. J. E. Demuth
IBM Corporation
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. M. G. Lagally
Department of Metallurgical
and Mining Engineering
University of Wisconsin
Madison, Wisconsin 53706

Dr. R. P. Van Duyne
Chemistry Department
Northwestern University
Evanston, Illinois 60637

Dr. J. M. White
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. D. E. Harrison
Department of Physics
Naval Postgraduate School
Monterey, California 93940

Dr. R. L. Park
Director, Center of Materials
Research
University of Maryland
College Park, Maryland 20742

Dr. W. T. Peria
Electrical Engineering Department
University of Minnesota
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson
Department of Metallurgy and
Materials Science
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. S. Sibener
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Arnold Green
Quantum Surface Dynamics Branch
Code 3817
Naval Weapons Center
China Lake, California 93555

Dr. A. Wold
Department of Chemistry
Brown University
Providence, Rhode Island 02912

Dr. S. L. Bernasek
Department of Chemistry
Princeton University
Princeton, New Jersey 08544

Dr. W. Kohn
Department of Physics
University of California, San Diego
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Richard Colton
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Dan Pierce
National Bureau of Standards
Optical Physics Division
Washington, D.C. 20234

Dr. R. Stanley Williams
Department of Chemistry
University of California
Los Angeles, California 90024

Dr. R. P. Messmer
Materials Characterization Lab.
General Electric Company
Schenectady, New York 22217

Dr. Robert Gomer
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Ronald Lee
R301
Naval Surface Weapons Center
White Oak
Silver Spring, Maryland 20910

Dr. Paul Schoen
Code 6190
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. John T. Yates
Department of Chemistry
University of Pittsburgh
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene
Code 5230
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. L. Kesmodel
Department of Physics
Indiana University
Bloomington, Indiana 47403

Dr. K. C. Janda
University of Pittsburgh
Chemistry Building
Pittsburg, PA 15260

Dr. E. A. Irene
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Adam Heller
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton SO9 5NH
UNITED KINGDOM

Dr. H. Tachikawa
Chemistry Department
Jackson State University
Jackson, Mississippi 39217

Dr. John W. Wilkins
Cornell University
Laboratory of Atomic and
Solid State Physics
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis
Department of Physics
University of California
Irvine, California 92664

Dr. D. Ramaker
Chemistry Department
George Washington University
Washington, D.C. 20052

Dr. J. C. Hemminger
Chemistry Department
University of California
Irvine, California 92717

Dr. T. F. George
Chemistry Department
University of Rochester
Rochester, New York 14627

Dr. G. Rubloff
IBM
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. Horia Metiu
Chemistry Department
University of California
Santa Barbara, California 93106

Dr. W. Goddard
Department of Chemistry and Chemical
Engineering
California Institute of Technology
Pasadena, California 91125

Dr. P. Hansma
Department of Physics
University of California
Santa Barbara, California 93106

Dr. J. Baldeschwieler
Department of Chemistry and
Chemical Engineering
California Institute of Technology
Pasadena, California 91125

Dr. J. T. Keiser
Department of Chemistry
University of Richmond
Richmond, Virginia 23173

Dr. R. W. Plummer
Department of Physics
University of Pennsylvania
Philadelphia, Pennsylvania 19104

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 41106

Dr. N. Winograd
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. Roald Hoffmann
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. A. Steckl
Department of Electrical and
Systems Engineering
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. G.H. Morrison
Department of Chemistry
Cornell University
Ithaca, New York 14853