

2

DTIC

AD-A208 165

**CHEMICAL
RESEARCH,
-DEVELOPMENT &
ENGINEERING
CENTER**

CRDEC-TR-052

**THE ACUTE TOXICITY
OF ISOPROPYLAMINE
AND 2-METHYLCYCLOHEXANOL**

**Mark V. Haley
Wayne G. Landis**

RESEARCH DIRECTORATE

April 1989

DTIC
ELECTE
MAY 19 1989
S E D

This document has been approved
for public release and sale; its
distribution is unlimited.

**U.S. ARMY
ARMAMENT
MUNITIONS
CHEMICAL COMMAND**



Aberdeen Proving Ground, Maryland 21010-5423

89 5 19 142

Disclaimer

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorizing documents.

Distribution Statement

Approved for public release; distribution is unlimited.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE				
4. PERFORMING ORGANIZATION REPORT NUMBER(S) CRDEC-TR-052			5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION CRDEC	6b. OFFICE SYMBOL (If applicable) SMCCR-RST-E	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) Aberdeen Proving Ground, MD 21010-5423		7b. ADDRESS (City, State, and ZIP Code)		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION CRDEC	8b. OFFICE SYMBOL (If applicable) SMCCR-RST	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c. ADDRESS (City, State, and ZIP Code) Aberdeen Proving Ground, MD 21010-5423		10. SOURCE OF FUNDING NUMBERS		
		PROGRAM ELEMENT NO.	PROJECT NO. 1C162622	TASK NO. A554
		WORK UNIT ACCESSION NO.		
11. TITLE (Include Security Classification) The Acute Toxicity of Isopropylamine and 2-Methylcyclohexanol				
12. PERSONAL AUTHOR(S) Haley, Mark V., and Landis, Wayne G.				
13a. TYPE OF REPORT Technical	13b. TIME COVERED FROM 88 Apr TO 88 Oct	14. DATE OF REPORT (Year, Month, Day) 1989 April		15. PAGE COUNT 13
16. SUPPLEMENTARY NOTATION				
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP		
06	06.09.11		Isopropylamine 2-MCH	
08	01		IPA Daphnia magna	
			2-methylcyclohexanol (continued on reverse)	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The toxicity of isopropylamine (IPA) and 2-methylcyclohexanol (2-MCH) was determined. Acute 48-hr bioassays were performed using the water flea, <u>Daphnia magna</u> . Ninety-six hour, growth inhibition assays were conducted using freshwater green eucaryotic algae, <u>Selenastrum capricornutum</u> , and <u>Ankistrodesmus falcatus</u> . Isopropylamine was more toxic to the test organisms than 2-MCH. The 48-hr EC50 of IPA to daphnia was 91.5 mg/L. When buffered with 10% HCl, the 48-hr EC50 of IPA increased to 148.8 mg/L. The 96-hr IC50 of buffered IPA to <u>S. capricornutum</u> was 118.4 mg/L. The 48-hr EC50 of 2-MCH to daphnia was 267.0 mg/L. The 96-hr IC50 of 2-MCH to algae was 395.0 mg/L. (M)				
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT ☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT. ☐ DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL SANDRA J. JOHNSON			22b. TELEPHONE (Include Area Code) (301) 671-2914	22c. OFFICE SYMBOL SMCCR-SPS-1

PREFACE

The work described in this report was authorized under Project No. 1C162622A554, Chemical Munitions. This work was started in April 1988 and completed in October 1988.

The use of trade names or manufacturers' names in this report does not constitute an official endorsement of any commercial products. This report may not be cited for purposes of advertisement.

Reproduction of this document in whole or in part is prohibited except with permission of the Commander, U.S. Army Chemical Research, Development and Engineering Center, ATTN: SMCCR-SPS-T, Aberdeen Proving Ground, Maryland 21010-5423. However, the Defense Technical Information Center and the National Technical Information Service are authorized to reproduce the document for U.S. Government purposes.

This report has been approved for release to the public.

Accession For	
1. MAJ	☒
2. TM	☐
Unpublished	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



Blank

CONTENTS

	Page
1. INTRODUCTION	7
2. METHODS AND MATERIALS	7
2.1 Daphnia Assays	7
2.2 Algal Growth Inhibition Assays	9
3. RESULTS	10
4. DISCUSSION	10
5. CONCLUSIONS	11
LITERATURE CITED	13

Blank

THE ACUTE TOXICITY OF ISOPROPYLAMINE AND 2-METHYLCYCLOHEXANOL

1. INTRODUCTION

To provide adequate protection to the environment, a data base must be established for all chemicals. A data base provides toxicity information that can be used in estimating environmental impact. Some of the first steps in establishing the aquatic toxicity of a compound are the daphnia and algae bioassays. These bioassays are short term, widely used, and have well-documented test procedures.

In this study, the acute toxicity of isopropylamine (IPA) and 2-methylcyclohexanol (2-MCH) is examined. Acute 48-hr bioassays were performed using the water flea, Daphnia magna, and 96-hr, growth inhibition assays were conducted using the freshwater green eucaryotic algae, Selenastrum capricornutum (S. capricornutum) and Ankistrodesmus falcatus (A. falcatus).

Table 1 shows some of the physical and chemical properties of the test compounds. Isopropylamine is used in the synthesis of pharmaceuticals, pesticides, and dyes.¹ 2-Methylcyclohexanol is used in lacquers, films, various solvents, and soaps.²

2. METHODS AND MATERIALS

Isopropylamine (lot #1295-23) was obtained from Dr. Eugene Olajos, Research Directorate, U.S. Army Chemical Research, Development and Engineering Center (CRDEC). Isopropylamine caused the pH value to shift when added to the media. Therefore, duplicate tests were run using buffered (pH 7.2, using 10% HCl) IPA solutions.


2-Methylcyclohexanol was obtained from Mr. Charles Crouse, also of the Research Directorate, CRDEC. All tests conformed to current U.S. Environmental Protection Agency (EPA)³ and American Society for Testing and Material (ASTM)⁴ guidelines.

2.1 Daphnia Assays.

D. magna were obtained from Dr. Freda Taub at the University of Washington (Seattle, WA). Daphnia were reared in the laboratory as described by Goulden et al.⁵ Daphnid stock cultures were fed a mixture of A. falcatus, S. capricornutum, and Chlamydomonas reinhardtii 90. The culture media was prepared from municipal drinking water that was hardened to 132 ppm total CaCO_3 .⁴ The pH was adjusted to 7.2-7.5. Ten neonates less than 24 hr old were placed into 250-mL glass beakers filled with 100 mL of the test solution. The test beakers were placed into an incubator with a light-dark cycle of 16:8 hr with 315 ft candles of light at 20 °C. Two replicates were used in each test. Daphnia were gently touched with a pastier pipet at 24 and 48 hr. If the daphnia could not swim actively for 15 s, immobilization (mortality) was recorded. The effective concentrations at which 50% of the organisms are immobilized (EC50) were computed using the probit analysis prepared by Stephan.⁶


Table 1. Some Physical and Chemical Properties of the Test Compounds^{1,2}

Isopropylamine



Synonyms:	2-aminopropane, monoisopropylamine, 2-propanamine
Description:	colorless liquid, ammonia order, basic
Mol wt:	59.11
Boiling pt:	33-34 °C
Density:	0.694
Flash pt:	-26 °C
Solubility:	miscible with water
Toxicity:	oral rat LD50 = 0.82 g/kg

2-Methylcyclohexanol



Synonyms:	0-methylcyclohexanol, 2-methylcyclohexyl, alcohol, hexahydrocresol and hexahydromethylphenol
Description:	colorless liquid or pale yellow, menthol odor
Mol wt:	114.19
Boiling pt:	173.0-175.3 °C
Freezing pt:	-50 °C
Density:	0.913
Flash pt:	67.7 °C
Solubility:	3-4% in water
Toxicity:	oral rat LD50 = 2.0 g/kg

The percent mortality values were plotted against concentration. A least-square regression line was drawn, and the EC50 was tabulated graphically. The graphically determined EC50 was used in verifying the probit analysis.

2.2 Algal Growth Inhibition Assays.

S. capricornutum and *A. falcatus* were also obtained from Dr. Freda Taub, University of Washington (Seattle, WA). Stock cultures of algae were maintained on 1.5% Difco-Bacto agar slants. Test algae were grown in a semiflow-through culture apparatus on T82MV(7) and taken during log phase growth for inoculation into the test flasks. Five-hundred milliliter Erlenmeyer flasks with ground glass stoppers were used as test chambers. One-hundred milliliters of media was placed in each test chamber and inoculated with approximately 4.0×10^4 algal cells per milliliter. The algae were placed in an incubator under the conditions described above for the daphnia 48-hr assays. Using a Newbauer Counting Chamber, cell densities were determined every 24 hr for 5 consecutive days. The area under the growth curve (A) was calculated using the following equation:

$$A = \frac{(N_0 + N_1) - 2N_0}{2} \times (t_1) + \frac{(N_1 + N_2) - 2N_0}{2} \times (t_2 - t_1) + \frac{(N_n - 1 + N_n) - 2N_0}{2} \times (t_n - t_n - 1) \quad (1)$$

where

N_0 = number of cells at t_0

N_1 = number of cells at t_1

N_n = number of cells at t_n

t_0 = time zero

t_1 = time of first measurement

t_n = time of the n^{th} measurement

The percent inhibition was calculated using the area under the growth curve. The following equation was used to calculate the percent inhibition (%In):

$$\%In = \frac{A_c - A_t}{A_c} \times 100 \quad (2)$$

where A_c = area of control growth curve, and A_t = area of treatment growth curve.

The percent inhibition values were plotted against the concentration. A least-square regression line was drawn, and the concentration at which algal growth is reduced to 50% of the control (IC₅₀) was determined. Analysis of Variance was run on the replicates to determine if any of the groups were significantly different. Dunnett's test was conducted to determine which treatment groups were different from the control.

3. RESULTS

Isopropylamine caused extreme pH-value shifts when added to the media (Figure 1); the dissolved oxygen remained unchanged. The unbuffered 48-hr EC₅₀ of IPA to daphnia was 91.5 mg/L. The buffered 48-hr EC₅₀ of IPA to daphnia was 148.8 mg/L.




Figure 1. pH Encountered During the Unbuffered Assays.

Unbuffered solutions of IPA caused a 100% decrease in algal growth (*S. capricornutum*) at concentrations down to 62.5 mg/L. When buffered, the 96-hr IC₅₀ of IPA to *S. capricornutum* was 118.4 mg/L. Figure 2 shows algal growth curves subjected to buffered and nonbuffered solutions of IPA. Algal growth, in the buffered and nonbuffered IPA tests, was significantly different from the control at $p \leq 0.05$.

2-Methylcyclohexanol was the less toxic of the two test compounds. The pH value and dissolved oxygen remained unchanged. The 48-hr daphnia EC₅₀ = 267.0 mg/L. The 96-hr IC₅₀ of 2-MCH to *A. falcatus* was 395.0 mg/L. At 100 mg/L of 2-MCH, algal growth was unchanged. At 250, 500, 750, and 900 mg/L of 2-MCH, algal growth was significantly reduced at $p \leq 0.05$.

4. DISCUSSION

Isopropylamine was more toxic than 2-MCH. The amine group in IPA caused a pH value shift when added to the media. Buffering the IPA solutions reduced the EC₅₀ and IC₅₀ from 38 to 47%, depending on test organism. However,

the IPA solutions were still ranked 4 (on a scale of 0-9 with 9 being the most toxic) on the scoring criteria for aquatic toxicity.⁸ If released into an aquatic ecosystem, the impact of IPA would depend on the buffering capacity of that system.

2-Methylcyclohexanol was ranked 2 on the scoring criteria for aquatic toxicity and exhibited no adverse effects on pH or dissolved oxygen.

5. CONCLUSIONS

Isopropylamine is more toxic than 2-MCH with a scoring criteria for aquatic toxicity of 4 and 2, respectively. Buffering solutions of IPA showed toxicity reduction. The buffering capacity of the ecosystem will play a strong roll in reducing the toxicity of IPA if an accidental spill occurs. A safety factor of 20 is commonly used when trying to estimate a not-to-exceed limit to prevent chronic effects. Using the most sensitive organism tested, the not-to-exceed level for buffered IPA is 5.9 mg/L. The not-to-exceed level for 2-MCH is 13.3 mg/L.




Figure 2. Growth Curves of *S. capricornutum* Exposed to Buffered and Nonbuffered IPA

LITERATURE CITED

1. Toxic and Hazardous Industrial Chemicals Safety Manual, For Handling and Disposal with Toxicity and Hazard Data, p 333, 1975.
2. Material Safety Data Sheet, Isopropyl Amine.
3. Users Guide: Procedures for Conducting Daphnia magna Toxicity Bioassays, EPA-660/8-87/011, U.S. Environmental Protection Agency, Washington DC, March 1987.
4. Guide for Conducting Acute Toxicity Tests with Fishes, Macro-invertebrates and Amphibians, Standard E729, American Society for Testing and Materials, Philadelphia, PA, 1986.
5. Goulden, C.E., Conotto, R.M., Hendrickson, J.A, Jr., Hornig, L.L., and Johnson, K.L., Procedures and Recommendations for the Culture and Use of Daphnia magna in Bioassay Studies, ASTM Special Technical Report Publication 766, p 139-160, American Society for Testing and Materials, Philadelphia, PA, 1982.
6. Stephen, C.E., Probit Analysis, U.S. Environmental Protection Agency, Duluth, MN.
7. Taube, F.B., "Generic Freshwater Microcosm Test," U.S. Environmental Protection Agency Federal Register, Vol. 52, No. 187 (1987).
8. O'Bryan, T., and Ross, R., Chemical Scoring System for Hazard and Exposure Assessments, Draft Report, Office of Toxic Substances, U.S. Environmental Protection Agency, Washington, DC.