

DOCUMENTATION PAGE

AD-A208 281

1a. REPO
UN
2a. SECUR

2b. DECLASSIFICATION / DOWNGRADING SCHEDULE

4. PERFORMING ORGANIZATION REPORT NUMBER(S)
SD30-5

6a. NAME OF PERFORMING ORGANIZATION
Armed Forces Radiobiology
Research Institute

6b. OFFICE SYMBOL
(If applicable)
AFRRI

6c. ADDRESS (City, State, and ZIP Code)
Defense Nuclear Agency
Bethesda, Maryland 20814-5145

8a. NAME OF FUNDING / SPONSORING
ORGANIZATION
Defense Nuclear Agency

8b. OFFICE SYMBOL
(If applicable)
DNA

8c. ADDRESS (City, State, and ZIP Code)
Washington, DC 20305

1b. RESTRICTIVE MARKINGS

3. DISTRIBUTION / AVAILABILITY OF REPORT
Approved for public release; distribution
unlimited.

5. MONITORING ORGANIZATION REPORT NUMBER(S)

7a. NAME OF MONITORING ORGANIZATION

7b. ADDRESS (City, State, and ZIP Code)

9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

10. SOURCE OF FUNDING NUMBERS

PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT ACCESSION NO.
NWED QAXM			

11. TITLE (Include Security Classification)
Changes in canine neutrophil function(s) following cellular isolation by
percoll gradient centrifugation or isotonic lysis

12. PERSONAL AUTHOR(S)
Gruber, P.F. and L'Alessandro, M.M.

13a. TYPE OF REPORT
Reprints

13b. TIME COVERED
FROM TO

14. DATE OF REPORT (Year, Month, Day)
1989 January

15. PAGE COUNT

16. SUPPLEMENTARY NOTATION

17. COSATI CODES

FIELD	GROUP	SUB-GROUP

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

DTIC
ELECTE
MAY 12 1989
A

20. DISTRIBUTION / AVAILABILITY OF ABSTRACT

☐ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT. ☐ DTIC USERS

21. ABSTRACT SECURITY CLASSIFICATION

UNCLASSIFIED

22a. NAME OF RESPONSIBLE INDIVIDUAL
Gloria Ruggiero

22b. TELEPHONE (Include Area Code)
(301)295-3536

22c. OFFICE SYMBOL
ISDP



CHANGES IN CANINE NEUTROPHIL FUNCTION(S) FOLLOWING
CELLULAR ISOLATION BY PERCOLL GRADIENT CENTRIFUGATION OR
ISOTONIC LYSIS^{1,2}

Dale F. Gruber* and Michele M. D'Alesandro
Experimental Hematology Department
Armed Forces Radiobiology Research Institute
Bethesda MD 20814-5145

ABSTRACT

Analysis of cellular effector function(s) often requires their isolation from other cellular types. Cell separatory techniques could mask, or select out, clinically important functional lesions. We examined differences in canine peripheral blood neutrophil functions, i.e. migration and H₂O₂ production, following two commonly used cell separation techniques: isotonic lysis or density gradient (Percoll) centrifugation. Separation methodology was observed to have a significant impact on both metabolic and mobility functions. In comparison to isotonic lysis, Percoll separation caused near 100% increases in random migration, near 40% decreases in chemotaxis and 70% increases in H₂O₂ production.

INTRODUCTION

Many functional and biochemical studies require large numbers of purified cells. Investigators have long sought to develop in vitro minimally manipulative methodologies allowing suitable cellular separation and recovery. Numerous new techniques have evolved some of which have included; cell sorting, electrophoresis and various affinity methods. Investigators have also attempted to make use of cellular density differences. The application of density separating techniques has resulted in three general approaches; (i) neutral density sedimentation, (ii) discontinuous-gradient, and (iii) isopycnic-gradient separations. Two of the more

frequently used separation mediums are Ficoll/Hypaque (a copolymer of sucrose and epichlorohydrin) and Percoll (colloidal silica coated with polyvinyl pyrrolidone). For many investigators Percoll has become the separation medium of choice because of its lower viscosity (1), greater cell yields (2), noncytotoxic character (3,4), and because it is not actively phagocytized (3). Since Percoll's inception, many papers making use of its separation qualities have been published. In this report, we demonstrate differences in canine neutrophil effector functions after separation by whole-blood isotonic lysis or Percoll density gradient methods.

MATERIALS AND METHODS

Dogs: Male and female littermates (<2 years old, 10-12 kg Hra Beagles) were used. Dogs were quarantined on arrival and screened for evidence of disease before being released to experiments. Animals were kennelled in an AAALAC-accredited facility in runs, and provided commercial dog chow and tap water ad libitum.

Cell Separation and Isolation: Peripheral blood samples in preservative-free heparin (10 U/ml) were drawn from the lateral saphenous vein, centrifuged (400 x g, 10 minutes) to remove plasma, diluted 2-fold in saline and divided into two aliquots. One aliquot was isototically lysed with 0.83% NH₄Cl (4°C, 5 minutes), washed in saline and resuspended in saline supplemented with 0.2% heat-inactivated fetal bovine serum (56°C, 30 minutes). The second aliquot was layered unto a 60% (p=1.077)/70% (p=1.080) discontinuous Percoll gradient and centrifuged (700 x g, 30 minutes, room temperature). Neutrophils sedimented to the 60%/70% interface whereas mononuclear cells (i.e., monocytes, lymphocytes) remained on top of the 60% Percoll. The neutrophil layer was aspirated from the gradient and washed to remove adherent Percoll. Contaminating red blood cells in the Percoll isolate were isototically lysed as described above, washed and resuspended in supplemented saline. On the basis of vital dye exclusion cellular viabilities were routinely >95%.

Cellular Migration: Neutrophil migration was evaluated by quantitating cells that migrated through a 10 um-thick polycarbonate membrane. A 48-well micro-chemotaxis chamber assembly (Neuro Probe Inc., Bethesda, MD) was used as previously described (5). Lower wells contained a 1:100 dilution of chemoattractant (zymosan activated plasma (6)) or buffer. Upper wells contained 50 ul of

cellular suspension which had been corrected to 2×10^6 neutrophils/ml. Chambers were incubated at 37°C in a humidified environment and gassed with 5% CO_2 and air for 1 hour. Membranes were fixed in 100% methanol for 1 minute before being stained with Dif-Quick differential stain (Fisher Scientific, Silver Spring, MD). Seven high power (100X) microscopic fields across the diameter of the well were examined and the mean cellular migration/field/hour was determined.

Quantitation of Intracellular H2O2 Production:

Neutrophil H2O2 production was assayed as described by Bass et al (7). Neutrophils (1×10^6 cells/ml) were incubated with 5 μM dichlorofluorescein diacetate (DCFH-DA) for 10 minutes at 37°C . DCFH-DA traverses the cell membrane and is converted by cellular esterases into intracellularly trapped nonfluorescent DCFH. Cytosolic H2O2 endogenously present in resting cells or produced following activation, oxidizes DCFH to DCF, a fluorescent analogue. The relative fluorescence intensity of the cell is, therefore, an indirect measure of cellular H2O2 production. Cells were analyzed on a FACS analyzer (Becton Dickinson, Mountain View, CA). Green fluorescence was monitored between 515 and 545 nm after excitation with a mercury arc lamp using a 485/22 nm excitation filter. Cells were stimulated with phorbol myristate acetate (100 ng/ml, 15 minutes, 37°C) and analyzed. Neutrophils were distinguished from other cellular types on the basis of coulter volumes(s) and right angle light scatter properties.

RESULTS

The leukocyte population isolated from peripheral blood by isotonic lysis was >85% neutrophils based on differential analysis. The neutrophil population at the 60%/70% Percoll interface was >95% neutrophils. Less than 5% of the total neutrophil population remained with the mononuclear cells above the 60% Percoll layer.

No differences due to sex were apparent within the parameters examined in this report. Male: compared to separation by isotonic lysis, random migration of Percoll separated neutrophils was increased >100% (Figure 1), H2O2 production was increased by 79% (Figure 2), and chemotactic migration decreased by 37% (Figure 3). Female: Random migration of Percoll separated neutrophils increased by 95% (Figure 1), H2O2 production increased by 66% (Figure 2), and chemotactic migration (Figure 3) decreased by 43% compared with cells isolated by whole blood lysis.




Figure 1. Random migration of canine neutrophils isolated by isotonic lysis alone or Percoll gradient separation followed by isotonic lysis. Each data point represents the mean $\pm$ SEM number of migrating cells per high-power field from a minimum of six animals.




Figure 2. H₂O₂ production following isotonic lysis alone or Percoll gradient separation followed by isotonic lysis. Each data point represents the mean $\pm$ SEM percent change in production (x100) following PMA stimulation from a minimum of six animals.




Figure 3. Chemotaxis of canine neutrophils to zymosan-activated normal dog plasma (10e-2 M) following isotonic lysis alone or Percoll gradient separation followed by isotonic lysis. Each data point represents the mean $\pm$ SEM number of migrating cells per high-power field from a minimum of six animals.

DISCUSSION

Clinically relevant immunological dysfunction(s) may be the result of subtle cellular alterations. Unfortunately studies of specific cellular effector function(s) often require the isolation of the cell(s) of interest from other contaminating elements, cellular and otherwise. We demonstrate here that the isolation method significantly affects the metabolic and motility effector function(s) of canine neutrophils. Previous studies have reported that isolation through Ficoll-metrizoate density gradients altered membrane events so as to produce an oxidative burst (8). Based on the data presented here, separation-induced anomalies can be extended to include metabolic and motility functions as well. Percoll separation followed by isotonic lysis, in comparison to lysis alone, increased the basal motility of neutrophils evidenced by significant increases in levels of random migration. Previous reports (9,10), in conjunction with the functional differences reported here, suggest that cellular anomalies may be inherent to the manner of separation and must be considered before making judgements on apparent effector cell dysfunction(s).

FOOTNOTES

1. Supported by the Armed Forces Radiobiology Research Institute, Defense Nuclear Agency, under research work unit 00130. Views presented in this paper are those of the authors; no endorsement by the Defense Nuclear Agency has been given or should be inferred.
2. Research was conducted according to the principles enunciated in the 'Guide for the Care and Use of Laboratory Animals' prepared by the Institute of Laboratory Research, National Research Council.

REFERENCES

1. Kunnick, J.T., Ostberg, L., Stegagno, M., Kimura, A.K., Orr, A., and Sjoberg, D., A rapid method for the separation of functional lymphoid cell populations of human and animal origin on PVP-silica (Percoll) density gradients, *Scand. J. Immunol.*, 10:563, 1979.
2. Dettman, G.L., and Wilber, S.M., Colloidal silica-aluminum modified-PVP density gradient centrifugation: centrifuge tube wall, cell

- adherence, aggregation, separation properties and comparison to BSA and Ficoll, *J. Immunol. Methods*, 27:205, 1979.
3. Penttoft, H., Laurent, T.C., Laas, T., and Kagedal, L., Density gradients prepared from colloidal silica particles coated with polyvinylpyrrolidone (Percoll), *Anal. Biochem.*, 88:271, 1978.
 4. Gmelig-Meyling, F., and Waldmann, T.A., Separation of human blood monocytes and lymphocytes on a continuous Percoll gradient, *J. Immunol. Methods*, 33:1, 1980.
 5. Harvath, L., Falk, W. and Leonard, E.J., Rapid quantitation of neutrophil chemotaxis: use of a polyvinylpyrrolidone-free polycarbonate membrane in a multiwell assembly, *J. Immunol. Methods*, 37:39, 1980.
 6. Redl, H., Flynn, P., Lamche, H., Schiessen, A., Schlag, G., and Hammerschmidt, D.E., Aggregation, chemotaxis and chemiluminescence of canine granulocytes. Studies utilizing improved cell preparation techniques, *Inflammation* 7:67, 1983.
 7. Bass, D.L., Parce, J.W., Dechatelet, L.R., Szejda, P., Seeds, M.C., and Thomas, M., Flow cytometric studies of oxidative product formation by neutrophils: a graded response to membrane stimulation, *J. Immunol.*, 130:1910, 1983.
 8. Berkow, R.L., Tzeng, D.Y., Williams, L.V., and Baehner, R.L., The comparative responses of human polymorphonuclear leukocytes obtained by counterflow centrifugal elutriation and Ficoll-Hypaque density centrifugation. I. Resting volume, stimulus-induced superoxide production, and primary and secondary granule release, *J. Lab. Clin. Med.*, 102:811, 1984.
 9. Seeds, M.C., Parce, J.W., Szejda, P. and Bass, D.A., Independent stimulation of membrane potential changes and the oxidative metabolic burst in polymorphonuclear leukocytes, *Blood*, 65:233, 1985.
 10. Poston, L., Jones, R.B. and Hilton, P.F., Sodium transport on polymorphonuclear leukocytes: effect of isolation by the Ficoll/Triosil method, *Clin. Sci.*, 62:563, 1982.



4120