

UN
SECURITY

AD-A208 292

DOCUMENTATION PAGE

1a. REPC
UN

2a. SECURITY CLASSIFICATION

2b. DECLASSIFICATION / DOWNGRADING SCHEDULE

4. PERFORMING ORGANIZATION REPORT NUMBER(S)

SR89-1

6a. NAME OF PERFORMING ORGANIZATION
Armed Forces Radiobiology
Research Institute

6b. OFFICE SYMBOL
(If applicable)
AFRRI

1b. RESTRICTIVE MARKINGS

3. DISTRIBUTION / AVAILABILITY OF REPORT

Approved for public release; distribution
unlimited.

5. MONITORING ORGANIZATION REPORT NUMBER(S)

6c. ADDRESS (City, State, and ZIP Code)

Defense Nuclear Agency
Bethesda, Maryland 20814-5145

7b. ADDRESS (City, State, and ZIP Code)

8a. NAME OF FUNDING / SPONSORING
ORGANIZATION

Defense Nuclear Agency

8b. OFFICE SYMBOL
(If applicable)

DNA

9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

8c. ADDRESS (City, State, and ZIP Code)

Washington, DC 20305

10. SOURCE OF FUNDING NUMBERS

PROGRAM
ELEMENT NO.
NWED
QAXM

PROJECT
NO.

TASK
NO.

WORK UNIT
ACCESSION NO.

11. TITLE (Include Security Classification)

Effects of ethiofos (WR-1071) and malathion on monkey visual discrimination performance

12. PERSONAL AUTHOR(S)

Bogo et al.

13a. TYPE OF REPORT

Reprints

13b. TIME COVERED

FROM TO

14. DATE OF REPORT (Year, Month, Day)

1989 January

15. PAGE COUNT

16. SUPPLEMENTARY NOTATION

17. COSATI CODES

FIELD	GROUP	SUB-GROUP

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

S
ELECT
MAY 12 1989
A

20. DISTRIBUTION / AVAILABILITY OF ABSTRACT

☐ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT. ☐ DTIC USERS

21. ABSTRACT SECURITY CLASSIFICATION

UNCLASSIFIED

22a. NAME OF RESPONSIBLE INDIVIDUAL

Gloria Ruggiero

22b. TELEPHONE (Include Area Code)

(301)295-3536

22c. OFFICE SYMBOL

ISDP

Symposium Editors: J. F. WEISS and M. G. SIMIC

Brief Communication



EFFECTS OF ETHIOFOS (WR-2721) AND RADIATION ON MONKEY VISUAL DISCRIMINATION PERFORMANCE

V. BOGO*, C. G. FRANZ*, A. J. JACOBS†, J. F. WEISS† and R. W. YOUNG‡

*Behavioral Sciences and †Radiation Biochemistry Departments, Armed Forces Radiobiology Research Institute, Bethesda, Maryland 20814-5145, U.S.A.

‡Radiation Policy Division, Radiation Sciences Directorate, Defense Nuclear Agency, Washington, DC 20305-1000, U.S.A.

WR-2721 (ethiofos) is a promising protector against radiation-induced lethality and may be useful in cancer radiotherapy (Davidson *et al.*, 1980). However, ethiofos also produces nausea, vomiting, diarrhea, and hypotension, which implies severe behavioral consequences (Bogo *et al.*, 1985). We studied the effects of ethiofos on behavior in monkeys and its ability to mitigate early transient incapacitation (ETI). ETI is the abrupt cessation of performance (for at least 1 min) following rapidly delivered, high doses of radiation and usually occurs 5–10 min after irradiation and lasts for 1–15 min (Bogo, 1988a).

Male rhesus (*Macaca mulatta*) monkeys ($N = 24$) were obtained from the Joint Services Primate Breeding Colony, Naval Aerospace Medical Research Laboratory, Pensacola, Florida, and quarantined 60 d for evaluation, tuberculosis testing, and stabilization. Monkeys were maintained in conventional stainless steel, wall-mounted cages in rooms maintained at $72 \pm 2^\circ\text{F}$, with $50\% \pm 10\%$ relative humidity using 100% conditioned fresh air at 12 air changes per hr. They were provided commercial monkey chow and hyperchlorinated tap water *ad libitum*, and were kept on a 12 hr light/12 hr dark, full-spectrum light cycle with no twilight.

Monkeys seated in plastic chairs were exposed to a TRIGA reactor pulse (50 msec) of neutron/gamma radiation (ratio 0.4). The doses were 14, 16, 22, and 24 Gy. A 22 Gy pulse was used in the ethiofos/radiation condition because it was known to be 90% effective in producing ETI in monkeys performing the speed stress visual discrimination task (SSVDT) (Bogo *et al.*, 1987). Since the ETI dose was well above the $\text{LD}_{50/30}$ of 6 Gy for a monkey, ethiofos was not expected to offer any protection from lethality. Ethiofos was given i.v. (saphenous vein) at least 30 min before irradiation. The 200 mg/kg dose was diluted in 60 cc saline and injected with an infusion pump over 5 min.

Performance was assessed with either a visual discrimination task (VDT) or an SSVDT. In these shock avoidance tasks, trained monkeys discriminated between a circle and a square (always the correct choice) randomly presented every 10 s on backlit press-plates. Maximum response time in the VDT was 5.0 s, and 0.7 s or less in the SSVDT. Performance was assessed 15 min before and after ethiofos, during radiation exposure, and for 90 min after irradiation. The test measures were (a) ETI 90 min following radiation, (b) percent correct choice following ethiofos and/or radiation, (c) emesis after ethiofos and/or radiation, and (d) time to death.

In the repeat-measure study, the same six monkeys (mean weight 8.3 kg) were tested on the SSVDT as follows: (a) sham ethiofos (saline)/sham radiation (baseline control), (b) ethiofos/sham radiation, and (c) ethiofos/radiation. A three week interval was allowed between tests. At the time of the study the number of large, trained monkeys was limited, and no concurrent radiation-only control group existed to illustrate ETI. Radiation-only controls were obtained from the Armed Forces Radiobiology Research Institute's Primate Behavioral Data Base in smaller but similarly handled monkeys (weight 3.7 kg) (Franz *et al.*, 1981). The 14 and 16 Gy irradiated monkeys performed the SSVDT, and the 24 Gy irradiated monkeys performed the VDT. All monkeys were monitored before and after exposure to record incidence of emesis and were




FIG. 1. Baseline, post-ethiofos, and post-ethiofos radiation performance on the SSVDT. Circles show the number of subjects with ETL. Significant effects, compared with controls are shown with asterisks ($p < 0.05$). $N = 6$.




FIG. 2. Radiation control conditions for Fig. 1. Performance on the SSVDT after 14 and 16 Gy radiation, and performance on the VDT after 24 Gy radiation. Circles show the number of subjects with ETL. $N = 6$ group.

monitored until death. Percent correct performance was analyzed by two-way ANOVAs. Chi-square analysis was used to compare incidence of ETL and emesis.

In the repeat-measure conditions, performance during the post-ethiofos test period was significantly below the same-subject baseline, ranging from 25% to 30% below baseline at 45 and 85 min posttreatment, respectively (Fig. 1). Performance of the ethiofos/radiation condition was significantly below both the same-subject baseline and ethiofos conditions after treatment. The decrements ranged from 35% below baseline 10 min after irradiation to 80% below baseline 45 min after exposure. Both the ethiofos and ethiofos/radiation conditions were overall significantly below the same-subject baseline.

In the radiation-only conditions, performance dropped an average of 56%, 10–15 min postexposure (Fig. 2). Performance in the 14 and 16 Gy conditions recovered to 15% below control after 20 min postexposure, while recovery was more gradual in the 24 Gy condition, to 25% below control at 70 min. All radiation-only conditions in Fig. 2 were significantly below the baseline condition in Fig. 1.

The incidence of ETI was significantly greater in the ethiofos/radiation condition (83%) than ethiofos alone (33%). The incidence of emesis did not differ significantly between the combined and ethiofos-alone conditions (42%). The time until death did not differ significantly between the monkeys irradiated with 24 Gy alone and those treated with ethiofos and irradiated with 22 Gy, i.e. average time to death was 24 hr.

Ethiofos produced a decrement in the SSVDT and offered no protection against radiation-induced ETI; performance was worse in the combined condition than in either the ethiofos or radiation conditions alone. These findings are similar to those previously reported in rats performing a motor function task (Bogo *et al.*, 1985). Since ethiofos is behaviorally toxic in two animal models, future work will focus on either reducing its behavioral toxicity or searching for other effective radioprotectors with less toxic effects on behavior (Bogo, 1988b).

Acknowledgements—We wish to thank J. Hartison and P. Mannon for training subjects and collecting data. This research was supported by the Armed Forces Radiobiology Research Institute, Defense Nuclear Agency (DNA), under work unit 00096. Views presented in this paper are those of the authors; no endorsement by DNA has been given or should be inferred. Research was conducted according to the principles enunciated in the *Guide for the Care and Use of Laboratory Animals* prepared by the Institute of Laboratory Animal Resources, National Research Council.

REFERENCES

- BOGO, V. (1988a) Early behavioral toxicity produced by acute ionizing radiation. *Comments on Toxicology* (in press).
 BOGO, V. (1988b) Behavioral radioprotection. *Pharmac. Ther.* **39**: 73–78, this volume.
 BOGO, V., JACOBS, A. J. and WEISS, J. F. (1985) Behavioral toxicity and efficacy of WR-2721 as a radioprotectant. *Radiat. Res.* **104**: 182–190.
 BOGO, V., FRANZ, C. G. and YOUNG, R. W. (1987) Effects of radiation on monkey visual discrimination performance. *Proceedings of the Eighth International Congress on Radiation Research*, p. 259. FIELDEN, E. M., FOWLER, J. F., HENDRY, J. H. and SCOTT, D. (eds) Taylor and Francis, London.
 DAVIDSON, D. E., GRISAN, M. M. and SWEENEY, T. R. (1980) Biological characteristics of improved radioprotectors. In: *Radiation Sensitizers. Their Use in the Clinical Management of Cancer*, pp. 309–320. BRADY, L. W. (ed.) Masson Publishing, New York.
 FRANZ, C. G., YOUNG, R. W. and MITCHELL, W. E. (1981) *Behavioral Studies Following Ionizing Radiation Exposures: A Data Base*, Technical Report TR81-4. Armed Forces Radiobiology Research Institute, Bethesda, Maryland.



Distribution Justification	
By _____	
Distribution/	
Availability Codes	
date	Avail and/or Special
A-1	20