

AD-A208 310

2

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0640

R&T Code 413g008

Technical Report No. 1

"Synthesis and Structural Characterization of Pt-Si Dimers
with Unusually Short Cross-Ring Silicon-Silicon Interactions"

by

Eugene A. Zarate, Claire A. Tessier-Youngs, and Wiley J. Youngs

Published

in the

Journal of the American Chemical Society

Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 44106

May 5, 1989

Reproduction in whole, or in part, is permitted for any purpose of the United States Government.

* This document has been approved for public release and sale: its distribution is unlimited.

DTIC
ELECTE
MAY 31 1989
S a H D

89 5 30 039

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) No. 1		5. MONITORING ORGANIZATION REPORT NUMBER(S) N00014-86-K-0640	
6a. NAME OF PERFORMING ORGANIZATION WILEY J. YOUNGS CASE WESTERN RESERVE UNIVERSITY	6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION OFFICE OF NAV RESEARCH Dr. HAROLD GUARD	
6c. ADDRESS (City, State, and ZIP Code) DEPARTMENT OF CHEMISTRY CASE WESTERN RESERVE UNIVERSITY CLEVELAND, OHIO 44106		7b. ADDRESS (City, State, and ZIP Code) CODE 1113 800 N. QUINCY STREET ARLINGTON, VA 22217	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION ONR	8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code) (see 7b)		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO	PROJECT NO.
		TASK NO.	WORK UNIT ACCESSION NO
11. TITLE (Include Security Classification) "Synthesis and Structural Characterization of Pt-Si Dimers with Unusually Short Cross-Ring Silicon-Silicon Interactions"			
12. PERSONAL AUTHOR(S) Eugene A. Zarate, Claire A. Tessier-Youngs, and Wiley J. Youngs			
13a. TYPE OF REPORT TECHNICAL	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, Month, Day) May 5, 1989	15. PAGE COUNT
16. SUPPLEMENTARY NOTATION Published in the Journal of the American Chemical Society			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)			
SEE ATTACHED			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT ☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT ☐ DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL		22b. TELEPHONE (Include Area Code)	22c. OFFICE SYMBOL

reprints, platinum
silicon compounds,
silicon. (Aug 1988)

Synthesis and Structural Characterization of Pt-Si Dimers with Unusually Short Cross-Ring Silicon-Silicon Interactions


Eugene A. Zarate, Claire A. Tessier-Youngs,* and Wiley J. Youngs*

Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 44106

Received November 5, 1987

Polysilanes have use in applications where photosensitivity is required and as precursors to β -SiC.¹ Polysilanes are usually prepared from dichloroorganosilanes and sodium at $\approx 100^\circ\text{C}$.² Several reports of transition-metal-catalyzed formation of polysilanes from R_2SiH_2 and RSiH_3 have appeared in which product distributions have been the main focus.³ The mechanism of this

Scheme 1^a



* **1a**, X = Y = H; **1b**, X = Y = Cl; **1c**, X = H, Y = Cl.

catalytic process is poorly understood.^{2b,4,5} Dimeric silyl- or hydride-bridged complexes of titanium and zirconium are the only reported metal-containing species isolated from catalytic silane polymerizations.⁶ Herein we report the synthesis and structural characterization of a series of platinum-silyl dimers with unusually short cross-ring Si-Si distances. These dimers are catalysts or catalyst precursors in silane oligomerization reactions.

The preparation of cocrystallized mixtures of three platinum-silyl dimers $\{(\text{Et}_3\text{P})_2\text{Pt}(\text{SiPhX})(\text{SiPhY})\text{Pt}(\text{PEt}_3)_2\}$ (**1a**, X = Y = H; **1b**, X = Y = Cl; and **1c**, X = H and Y = Cl) from two very different reaction paths is shown in Scheme 1. The reaction of $\text{PtCl}_2(\text{PEt}_3)_2$ with LiSiHPh_2 in THF (path a, Scheme 1) yields a small quantity of the cocrystallized mixture **1a-c** as air stable yellow crystals suitable for X-ray diffraction studies. The complexes **1a-c** are the major platinum-containing products produced by the reaction of $\text{PtCl}_2(\text{PEt}_3)_2$ with SiH_3Ph in THF with the addition of a slight excess of sodium powder (path b, Scheme 1).⁷




Figure 1. ORTEP labeling diagram of **1a-c** and **1b**. The thermal ellipsoids are drawn at the 20% probability level. Hydrogens are omitted for clarity.

- (1) For a recent review, see: West, R. J. *Organomet. Chem.* **1986**, 300, 327-346.
- (2) (a) Ojima, I.; Inaba, S.-I.; Kogure, T.; Nagai, Y. *J. Organomet. Chem.* **1973**, 55, C7-C8. (b) Yamamoto, K.; Okinoshima, H.; Kumada, M. *J. Organomet. Chem.* **1971**, 27, C31-C32; **1970**, 23, C7-C8. (c) Corey, J. Y.; Chang, L. S.; Corey, E. R. *Organometallics* **1987**, 6, 1595-1596. (d) Brown-Wensley, K. A. *Organometallics* **1987**, 6, 1590-1591. (e) Lappert, M. F.; Maskell, R. K. *J. Organomet. Chem.* **1984**, 264, 217-228.

- (3) (a) Curtis, M. D.; Epstein, R. S. *Adv. Organomet. Chem.* **1981**, 19, 213-255. (b) Petz, W. *Chem. Rev.* **1986**, 86, 1019-1047. (c) Zybail, C.; Müller, G. *Angew. Chem., Int. Ed. Engl.* **1987**, 669-670. (d) Straus, D. A.; Tilley, T. D.; Rheingold, A. L. *J. Am. Chem. Soc.* **1987**, 109, 5872-5873.
- (4) (a) Aitken, C. T.; Harrod, J. F.; Samuel, E. *J. Am. Chem. Soc.* **1986**, 108, 4059-4066. (b) Aitken, C. T.; Harrod, J. F.; Samuel, E. *Can. J. Chem.* **1986**, 64, 1677-1679. (c) Aitken, C. T.; Harrod, J. F.; Samuel, E. *J. Organomet. Chem.* **1985**, 279, C11-C13. (d) Harrod, J. F.; Yun, S. S. *Organometallics* **1987**, 6, 1381-1387.
- (5) $\text{PtCl}_2(\text{PEt}_3)_2$ was prepared from $\text{PtCl}_2(\text{NC-Ph})_2$ and PEt_3 . Its properties were as reported by Jensen (Jensen, K. A. Z. *Anorg. Allg. Chem.* **1936**, 229, 225-281).
- (6) Gilman, H.; Steudel, W. *Chem. Ind.* **1959**, 1094.
- (7) All reactions were conducted under an argon atmosphere. The product from path a was obtained by removing the THF under vacuum followed by extraction with and recrystallization from hexane. The reaction mixture of path b showed the evolution of gas, presumably H_2 . Filtration of the mixture and removal of the solvent leaves a dark brown oil which when quickly washed with acetone yields the yellow solid **1a-c**. Low isolated yields (10-20%) of pure **1a-c** from path b are due to reaction of acetone with **1a-c** and to difficulties in completely removing traces of the liquid oligomeric silanes from **1a-c** via subsequent recrystallizations from hexane.

Crystals of pure **1b** can be obtained by treating toluene solutions of **1a-c** with carbon dioxide followed by recrystallization from benzene. Apparently the Si-H bonds of **1a** and **1c** react with carbon dioxide to yield unidentified complexes, whereas **1b** is unreactive.

Both paths a and b in Scheme 1 are complicated by partial chlorination of the Si-H bonds. This is not unexpected because Noble metals catalyze chlorination of Si-H bonds using a variety of chloride sources.⁸ Path a proceeds with the net loss of a phenyl substituent. We have not determined whether this can be attributed to a metal-catalyzed disproportionation of the substituents^{2a,d,3a,9,10} or to rearrangement of the silyllithium reagent.¹¹

The addition of catalytic quantities of **1a-c** to phenylsilane yields di- and trisilanes, with some scrambling of the phenyl and hydride substituents.¹² The di- and trisilanes are also produced by the addition of $\text{PtCl}_2(\text{PEt}_3)_2$ to excess SiH_3Ph and are byproducts of path b.¹²

Field desorption mass spectrometry of **1a-c** clearly shows that all three dimers are present in the crystalline compound. The X-ray study of the product from path a was performed.¹³ Alternating refinements to convergence holding the thermal parameters of the chlorine atom fixed while the occupancy factor was freed and vice versa shows that a silyl chloride is present in place of a silyl hydride approximately 65% of the time in this particular crystalline mixture.¹³ Pure **1b** has also been crystallographically characterized.¹⁴ All products were further characterized by IR and NMR spectroscopy.^{15,16}

An ORTEP drawing of the structure of **1a-c** and pure **1b** is shown in Figure 1. The silicon atoms and H or Cl and phenyl substituents lie in a plane which is perpendicular to the Pt-Pt vector and the plane defined by the platinum-silicon ring. The Si-Si separations in **1a-c** (2.575 (15)–2.602 (4) Å) are within the range of known single Si-Si bonds (2.33–2.70 Å)¹⁷ and are the shortest cross-ring Si-Si distances in any metal-silicon four-membered ring.^{18–20} The Pt-Si bonds (2.355 (7)–2.383 (8) Å) are long when compared to other Pt-Si bonds in environments of similar steric demands.^{10,18} The acute deviation of the angle at the "square-planar" platinum atoms (Si-Pt-Si = 65.9 (3)–66.5 (1)°) and the increase in the angles of the "tetrahedral" silicon atoms (Pt-Si-Pt = 113.6 (1)–114.2 (3)°) suggest that the Si-Si interaction forms

an integral part of the bonding scheme. The acute Si-Pt-Si angle is not a result of van der Waals repulsions between the platinum atoms because the Pt-Pt distances range from 3.973 (1)–3.997 (2) Å, whereas platinum has a van der Waals radius of 1.7–1.8 Å.²² The large cross-ring Si-Si separations (3.852–4.225 Å),²¹ acute M-Si-M angles (69.6–75.0°), and short M-M distances (2.708–3.183 Å) of most other metal-silicon dimers are in sharp contrast to those of **1a-c**.^{10,18–20} Only $[\text{Cp}_2\text{TiSiH}_2]_2$ has a similar cross-ring Si-Si distance (2.69 Å)²⁰ and M-Si-M angle (103°).²⁰

The π -coordination of the disilene $\text{Ph}_2\text{Si}=\text{SiPh}_2$ to two Pt-(PEt₃)₂ moieties provides a bonding picture of the complexes **1a-c** which is consistent with their geometry. This bonding picture of **1a-c** has been supported by a recent theoretical investigation.²³ The π -coordination of a bulky disilene to a single platinum has been reported, but crystallographic data are not available.²⁴ As a consequence of π -coordination, the Si-Si distance of **1a-c** is lengthened by 19–22% from that in free disilenes (2.13–2.16 Å).²⁵ This percent lengthening can be compared to the 12% or 27% lengthening of the C-C distance in $\text{CH}_2=\text{CH}_2$ on π -coordination to two zirconium centers in $\text{Zr}_2\text{X}_6(\text{PEt}_3)_4(\text{CH}_2=\text{CH}_2)$ (X = Br or Cl, respectively).²⁶ In **1a-c** and $\text{Zr}_2\text{X}_6(\text{PEt}_3)_4(\text{CH}_2\text{CH}_2)$ the resultant Si-Si (2.58–2.60 Å) or C-C (1.56 and 1.69 Å) separations are longer than the "typical" single bond distances (Si-Si = 2.34 Å and C-C = 1.54 Å).²² The bonding descriptions of **1a-c** and the zirconium-ethylene complexes²⁶ are extensions of the Dewar-Chatt-Duncanson (DCD) model, which recently has also been applied to the cross-ring Si-Si interactions in $[\text{R}_2\text{SiX}]_2$ (X = CH₂, NH, O, PH, or S).²⁷ The chemistry of **1a-c** shows similarities to that of π -coordinated diphosphene complexes because the synthesis of $\text{M}(\eta^2\text{-RP}=\text{PR})\text{L}_n$ can involve monophosphorus reagents^{28,29} and similar DCD bonding schemes have been developed.²⁹ As a consequence of the bonding scheme the silicon atoms of **1a-c** are pentacoordinated in a highly distorted trigonal bipyramidal geometry.^{30,31} Placing the platinum atoms in the axial positions is consistent with the relatively long Pt-Si bonds,³⁰ and this results in positioning the hydrides in their favored

(8) (a) Hazeldine, R. N.; Parish, R. V.; Parry, D. J. *J. Chem. Soc. A* **1969**, 683–690. (b) Citron, J. D.; Lyons, J. E.; Sommer, L. H. *J. Org. Chem.* **1969**, *34*, 638–640. (c) Citron, J. D. *J. Org. Chem.* **1969**, *34*, 1977. (d) Sommer, L. H.; Citron, J. D. *J. Org. Chem.* **1967**, *32*, 2470. (e) Watanabe, H.; Okawa, T.; Maraoka, T.; Nagai, Y. *Chem. Lett.* **1981**, 1321–1322. (f) Watanabe, H.; Kato, M.; Okawa, T.; Koguyou, Y.; Nagai, Y.; Goto, M.; *Appl. Organomet. Chem.* **1987**, *1*, 157–169.

(9) Benkeser, R. A.; Yeh, M.-H. *J. Organomet. Chem.* **1984**, *264*, 239–244.

(10) Aylett, B. J. *Adv. Inorg. Chem. Radiochem.* **1982**, *25*, 1–133.

(11) Wittenberg, D.; Gilman, H. *Quart. Rev.* **1959**, *13*, 116–145.

(12) Determined by FD-MS, IR, and NMR data.

(13) (a) Crystallographic data for **1a-c**: space group $P2_1/n$, $a = 10.985$ (2) Å, $b = 11.803$ (2) Å, $c = 17.636$ (3) Å, $\beta = 99.62$ (2)°, $V = 2254.4$ (8) Å³, and $Z = 2$. Intensity data were collected from $3.0 < 2\theta < 50.0^\circ$. Refinement on the basis of 1946 reflections ($I \geq 3\sigma$) resulted in a final $R = 0.068$ and $R_w = 0.070$. (b) UCLA Crystallographic Computing Package, December 2, 1985. Numerical absorption correction was applied.

(14) Crystallographic data for **1b**: space group $P2_1/n$, $a = 10.962$ (2) Å, $b = 11.833$ (2) Å, $c = 17.609$ (4) Å, $\beta = 99.62$ (2)°, $V = 2252.2$ (8) Å³, and $Z = 2$. Intensity data were collected from $3.0 < 2\theta < 52.0^\circ$. Refinement on the basis of 4116 reflections ($I \geq 3\sigma$) resulted in a final $R = 0.054$ and $R_w = 0.043$. See ref 13b.

(15) Selected spectral data for the mixture **1a-c**: IR (Nujol mull) $\nu_{\text{Si-H}} = 1955$ (m) and 1980 (m) cm^{-1} ; ^1H NMR (C_6D_6) 0.85 (m, CH_3), 1.72–1.83 (m, CH_2), 4.37 (m, SiH), 7.25, 7.71, and 7.94 (t, d, and d, respectively, C_6H_5). Except for the absence of the Si-H signals, the IR and NMR data for **1b** are virtually identical with those of **1a-c**.

(16) Hydride bridges on the metal-silicon bonds are a common feature in the structures of metal-silicon four-membered rings.¹⁰ We find no evidence of such in the crystallographic or spectra data of **1a-c** or **1b**.

(17) (a) Wiberg, N.; Schuster, H.; Simon, A.; Peters, K. *Angew. Chem., Int. Ed. Engl.* **1986**, *25*, 79–80. (b) Watanabe, H.; Kato, M.; Okawa, T.; Nagai, Y.; Goto, M. *J. Organomet. Chem.* **1984**, *271*, 225–233. (c) Weidenbruch, M. *Comments Inorg. Chem.* **1986**, *5*, 247–261.

(18) Auburn, M.; Ciriano, M.; Howard, J. A. K.; Murray, M.; Pugh, N. J.; Spencer, J. L.; Stone, F. G. A.; Woodward, P. *J. Chem. Soc., Dalton Trans.* **1980**, 659–666.

(19) (a) Bennet, M. J.; Simpson, K. A. *J. Am. Chem. Soc.* **1971**, *93*, 7156. (b) Crozat, M. M.; Watkins, S. F. *J. Chem. Soc., Dalton Trans.* **1972**, 2512–2515. (c) Cowie, M.; Bennett, M. J. *Inorg. Chem.* **1977**, *16*, 2321–2325 and 2325–2329. (d) Simon, G. L.; Dahl, I. F. *J. Am. Chem. Soc.* **1973**, *95*, 783.

(20) (a) Hencken, G.; Weiss, E. *Chem. Ber.* **1973**, *106*, 1747–1751. (b) $[\text{Cp}_2\text{TiHSi}(\text{Ph})\text{H}]_2$ is unlike any other metal-silicon dimer (Si-Si = 3.820 Å, Ti-Ti = 3.890 Å, $\angle\text{Ti-Si-Ti} = 90.40^\circ$ and 90.88° , and $\angle\text{Si-Ti-Si} = 87.91^\circ$ and 89.18°).²¹

(21) Cross-ring Si-Si separations and other parameters were calculated by us with data provided in the original reference.

(22) Huheey, J. E. *Inorganic Chemistry*, 3rd. ed.; Harper & Row: New York, 1983, pp 258–259 (van der Waals radii) and pp A32–A40 (bond lengths).

(23) Anderson, A. B.; Shiller, P.; Zarate, E. A.; Tessier-Youngs, C. A.; Youngs, W. J., submitted to *J. Am. Chem. Soc.*

(24) West, R. *Angew. Chem., Int. Ed. Engl.* **1987**, *26*, 1201–1211.

(25) (a) Fink, M. J.; Michalczyk, M. J.; Hallen, K. J.; West, R.; Michl, J. *Organometallics* **1984**, *3*, 793–800. (b) Masamune, S.; Muarkami, S.; Snow, J. T.; Williams, D. J. *Organometallics* **1984**, *3*, 333–334.

(26) (a) Cotton, F. A.; Kibala, P. A. *Polyhedron* **1987**, *6*, 645–646. (b) For a related complex, see: Burns, C. J.; Andersen, R. A. *J. Am. Chem. Soc.* **1987**, *109*, 915–917.

(27) (a) Grev, R. S.; Schaefer, H. F. *J. Am. Chem. Soc.* **1987**, *109*, 6577–6585. (b) Fink, M. J.; Haller, K. J.; West, R.; Michl, J. *J. Am. Chem. Soc.* **1984**, *106*, 822–823. (c) Yokelson, H. B.; Millevolte, A. J.; Adams, B. R.; West, R. *J. Am. Chem. Soc.* **1987**, *109*, 4116–4118.

(28) (a) Schäfer, H.; Binder, D.; Fenske, D. *Angew. Chem., Int. Ed. Engl.* **1985**, *24*, 522–524. (b) Schäfer, H. *Z. Naturforsch., B: Anorg. Chem., Org. Chem.* **1979**, *34B*, 1358–1361. (c) Hey, E.; Engelhardt, L. M.; Raston, C. L.; White, A. H. *Angew. Chem., Int. Ed. Engl.* **1987**, *26*, 81–82. (d) Caminade, A.-M.; Majoral, J.-P.; Sanchez, M.; Mathieu, R.; Attali, S.; Grand, A. *Organometallics* **1987**, *6*, 1459–1465.

(29) (a) Cowley, A. H.; Norman, N. C. *Prog. Inorg. Chem.* **1986**, *34*, 1–63. (b) Cowley, A. H. *Polyhedron* **1984**, *3*, 389–432.

(30) Holmes, R. R. *Prog. Inorg. Chem.* **1984**, *32*, 119–235.

(31) For examples of pentacoordinated silicon in transition-metal complexes see ref 10, 18, 21a, and 22b and the following: (a) Schubert, U.; Alt, H. G. *Organometallics* **1987**, *6*, 469–472. (b) Smith, R. A.; Bennett, M. J. *Acta Crystallogr., Sect. B: Struct. Crystallogr. Cryst. Chem.* **1977**, *B33*, 1118–1122. (c) Schubert, U.; Scholz, G.; Müller, J.; Ackermann, K.; Wörle, B.; Stansfield, R. F. D. *J. Organomet. Chem.* **1986**, *306*, 303–326. (d) Carré, F.; Colomer, E.; Corriu, R. J. P.; Vioux, A. *Organometallics* **1984**, *3*, 1272–1278.

equatorial sites.³² The inclusion of the Si-Si bond in two fused three-membered rings can help explain its length.³³

Acknowledgment. We thank Dr. William Kroenke of BFGoodrich Co. for mass spectra, Professor Robert West for providing information prior to publication, and the Office of Naval Research for support.

Registry No. **1a**, 113999-66-9; **1b**, 113999-67-0; **1c**, 113999-68-1; $\text{PtCl}_2(\text{PEt}_3)_2$, 15390-95-1; LiSiHPh_2 , 17950-77-5; PhSiH_3 , 694-53-1.

Supplementary Material Available: Tables of data collection and refinement details, atomic positional parameters, thermal parameters, and bond lengths and angles for the cocrystallized dimers **1a-c** and dichloride dimer **1b** (10 pages); observed and calculated structure factors for **1a-c** and **1b** (27 pages). Ordering information is given on any current masthead page.

(32) Breliere, C.; Carré, F.; Corriu, R. J. P.; Poirier, M.; Royo, G. *Organometallics* **1986**, *5*, 388.

(33) (a) Dabisch, T.; Schoeller, W. W. *J. Chem. Soc., Chem. Commun.* **1986**, 896-898. (b) Schoeller, W. W.; Dabisch, T.; Busch, T. *Inorg. Chem.* **1987**, *26*, 4383-4389. (c) Schleyer, P. v. R.; Sax, A. F.; Kalcher, J.; Janoschek, R. *Angew. Chem., Int. Ed. Engl.* **1987**, *26*, 364-366. (d) Collins, S.; Butler, R.; Rauk, A. J. *Am. Chem. Soc.* **1987**, *109*, 2564-2569. (e) Nagase, S.; Kudo, T. *Organometallics* **1987**, *6*, 2456-2458. (f) Schleyer, P. v. R.; Janoschek, R. *Angew. Chem., Int. Ed. Engl.* **1987**, *26*, 1267-1268.



Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	20