

Unclassified
SECURITY

1
T DOCUMENTATION PAGE

1a. REPORT NUMBER AD-A208 875		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION Unclassified		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; Distribution unlimited.	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) GL-TR-89-0132		7a. NAME OF MONITORING ORGANIZATION	
6a. NAME OF PERFORMING ORGANIZATION Geophysics Laboratory	6b. OFFICE SYMBOL (if applicable) PHP	7b. ADDRESS (City, State, and ZIP Code)	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8b. OFFICE SYMBOL (if applicable)		10. SOURCE OF FUNDING NUMBERS	
8c. ADDRESS (City, State, and ZIP Code)		PROGRAM ELEMENT NO. 62101F	PROJECT NO. 7601
		TASK NO. 22	WORK UNIT ACCESSION 01
11. TITLE (Include Security Classification) The IMF Bz and Solar Wind Speed Dependence for Precipitating Ion Hemispheric Energy Flux			
12. PERSONAL AUTHOR(S) D.H. Brautigam, M.S. Gussenhoven, D.A. Hardy			
13a. TYPE OF REPORT Reprint	13b. TIME COVERED FROM TO	14. DATE OF REPORT (Year, Month, Day) 1989 May 19	15. PAGE COUNT 4
16. SUPPLEMENTARY NOTATION Reprinted from Adv. Space Res. Vol. 8, No. 9-10, pp (9)65-(9)68, 1988			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
		Auroral Ions, Energetic Particles, Interplanetary Magnetic Field, Solar Wind Speed, Solar Wind Coupling, Auroral Morphology	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) A new statistical study of the total power input to high latitude regions from precipitating ions, making primary data sorts by the IMF Bz and solar wind speed, shows that a) the minimum power input occurs for low, positive values of Bz and low values of Vsw, b) power input increases proportionately more with increasing Vsw than with increasing Bz negative.			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT ☐ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT ☐ DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL M.S. Gussenhoven		22b. TELEPHONE (Include Area Code) (617) 377-3212	22c. OFFICE SYMBOL PHP

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted
All other editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

Unclassified

89 6 05 072

THE IMF Bz AND SOLAR WIND SPEED DEPENDENCE FOR PRECIPITATING ION HEMISPHERIC ENERGY FLUX

D. H. Brautigam, M. S. Gussenhoven and D. A. Hardy

Air Force Geophysical Laboratory, Hanscom AFB Bedford, MA 01731, U.S.A.

ABSTRACT

A new statistical study of the total power input to high latitude regions from precipitating ions, making primary data sorts by the IMF Bz and solar wind speed, shows that a) the minimum power input occurs for low, positive values of Bz and low values of Vsw, b) the power input increases proportionately more with increasing Vsw than with increasing Bz negative.

INTRODUCTION

One measure of the effectiveness of energy transfer from the solar wind to the magnetosphere is the total energy flux of particles precipitating into the high latitude regions. Previous statistical studies have determined the power input of precipitating electrons as a function of the geomagnetic activity index Kp [1,2,3,4] and for the interplanetary magnetic field (IMF) Bz component either positive or negative [4]. Power input from precipitating ions as a function of Kp was determined by Hardy et al. [5]. These studies show that the total precipitating electron energy flux is directly proportional to Kp, and that electrons contribute about seven times the energy flux of ions. Foster showed that more electron energy flux flows into the auroral regions during periods of Bz south than for Bz north [4].

Efforts to quantify the solar wind-magnetosphere electrical coupling strength, by relating the cross-polar-cap potential difference to combinations of the solar wind speed and the direction of the IMF have been reviewed by Reiff and Luhmann [6]. A direct dependence of the potential difference has been shown for Kp, solar wind speed, and magnitude of the southward component of the IMF. However, the coupling is not so clearly defined for times during which Bz is northward.

Here we present results showing the variation of ion energy flux input to high latitudes as a function of the IMF Bz and the solar wind speed, Vsw. By making separations of the precipitating ion data for strong and weak Bz north and south, as well as a low and high Vsw, we have found that the minimum power input occurs for weak Bz north and low Vsw, and that the dependence on Vsw is strong.

INSTRUMENTATION, DATA COVERAGE AND STATISTICAL PROCEDURES

The data used in this study were taken onboard two United States Defense Meteorological Satellite Program satellites, DMSP/F6 and DMSP/F7. The satellites are in sun-synchronous, circular, polar orbits at 840 km altitude, and are three-axis stabilized. DMSP/F6 orbits lie in the 0640-1840 meridian plane, and DMSP/F7 in the 1030-2230 meridian plane. On each satellite, electrostatic analyzers of identical design measure electrons and ions in the energy range from 30 eV to 30 keV in twenty logarithmically spaced intervals [7]. For each species a complete spectrum is returned each second. The detectors' apertures are fixed with respect to the satellite, pointing upward. At high latitudes they measure precipitating particles.

Due to the offset between geographic and geomagnetic poles, the sun-synchronous satellites, each day, sweep out a wide swath of values in corrected geomagnetic latitude (MLAT) and magnetic local time (MLT). Thus, the combined satellite, combined hemispheric data coverage is nearly complete for this coordinate system in the auroral regions. There are minor data gaps in the postnoon and postmidnight local time sector for active periods when the auroral oval has expanded to low latitudes.

We use data collected for three years on the DMSP/F6 satellite and two years on the DMSP/F7 satellite (5 satellite-years of data) and spanning the time period January, 1983 to March, 1986, to create statistical maps of ion precipitation in MLAT-MLT coordinates. The spatial grid into which the data are binned has an MLAT resolution of 2 degrees for 50-60 and 80-90 degrees MLAT, and 1 degree for 60-80 degrees MLAT, and an MLT resolution of one half hour.

Additionally, the data are sorted by the IMF component, B_z , in solar magnetospheric coordinates, and the solar wind speed, V_{sw} . Each ion spectrum in the data base is associated with a one-hour averaged value of B_z and V_{sw} , taken from the listing provided by NSSDC, and offset by a one-hour time lag. There are 4 IMF B_z bins: B_z large and negative (-10 to -3 nT); B_z small and negative (-3 to 0 nT); B_z small and positive (0 to +3 nT) and B_z large and positive (+3 to +10 nT). The average B_z values for each bin are -4.6, -1.3, +1.3, and +4.6 nT, respectively. There are 2 V_{sw} bins: low (from 200 to 400 km/s) and high (from 400 to 800 km/s), with average V_{sw} values of 364 and 518 km/s, respectively. Ultimately, 17 million ion spectra are separated into the eight primary B_z - V_{sw} maps defined in MLAT-MLT coordinates. The statistics for the low V_{sw} cases range from .5 to 1.8 million spectra/-map, while the range for the high V_{sw} cases is from 1.4 to 4.9 million spectra/map.

For each MLAT-MLT bin an average ion spectrum is calculated. Data gaps in the spatial coverage are filled by interpolating between neighboring data points. Since a significant portion of the high energy tail of ion spectra in the evening diffuse auroral region lies beyond the 30 keV cutoff of our instrument we extrapolated the average spectra to 100 keV. This results in a correction of the hemispheric energy flux which is dependent on activity level and ranges from an increase of 14% for the quietest case (low V_{sw} , weak B_z north) to 75% for the most active case (high V_{sw} , strong B_z north). The same qualitative trend of energy flux versus B_z/V_{sw} occurs for both the unextrapolated and the extrapolated cases. However, we believe that the extrapolated values provide a necessary correction to the otherwise artificially depressed values due to the detectors' upper limit of 30 keV. See Hardy *et al.* [5] for details on data reduction techniques (cross satellite normalization, smoothing, and extrapolating).

For each spatial bin we calculate moments of the distribution function from the average spectrum, and, in particular, the energy flux. We then integrate over the full spatial map (i.e., above 50° MLAT), using the assumption that the distributions are isotropic, to obtain the total precipitating ion energy flux input for each of the eight average interplanetary conditions.

RESULTS AND DISCUSSION

The total hemispheric precipitating ion energy flux, in gigawatts, is tabulated in Table 1 and plotted in Figure 1b for the eight B_z - V_{sw} conditions.

TABLE 1 Hemispheric Energy Flux (G Watts)

V_{sw} / B_z	-4.6 nT	-1.3 nT	+1.3 nT	+4.6 nT
364 km/s	4.67 (3.02)	2.02 (1.59)	1.36 (1.19)	1.85 (1.56)
518 km/s	5.74 (3.27)	2.96 (1.87)	2.27 (1.53)	2.91 (2.14)

Note: Energy flux values from unextrapolated spectra are in parentheses.

For a constant value of V_{sw} , the hemispheric energy flux maximizes for strong B_z south, minimizes for weak B_z north, and has nearly equal intermediate values for weak B_z south and strong B_z north. The variation in energy flux for each B_z interval, due to the V_{sw} difference alone, is a constant 1 GW (for extrapolated values). This offset is significant both because of its relatively large magnitude and because of its independence of B_z . If we define the ground state of the magnetosphere as one in which the precipitating particle energy flux is minimized, then the ground state occurs during periods of weak B_z north and low V_{sw} . It is not the limit of extrapolating low solar wind velocity states to higher and higher positive values of B_z .

In Figures 1a and 1b we summarize the results of the major studies of precipitating particle energy flux input using low altitude satellite data.

Figure 1a gives the total precipitating energy flux into the auroral zone as a function of K_p for electrons (filled symbols) and ions (open symbols). The agreement between the various electron studies is excellent. For both electrons and ions the log of the energy input increases in a linear fashion with increasing K_p for K_p greater than one. The minimum value is .5 GW for electrons and .7 GW for ions. Both values are somewhat depressed below a projected linear value. For this and other values of K_p , the electrons contribute between 6 - 8 times the energy flux of ions.

Figure 1b shows the results of the ion study discussed above, and those of Foster /4/ for which electron data were separated into two B_z bins, $> +3$ nT and < -3 nT. For comparison with Foster, the ion energy flux for the high and low V_{sw} cases are averaged for the corresponding values of B_z . The electron energy flux input is about 6 times that for the ions, in agreement with the Kp studies. Assuming that the same scaling of energy flux between electrons and ions holds for all values of B_z , we infer that as a function of B_z the minimum electron power input should be about 11 GW and would also occur under weakly positive B_z conditions.

Comparing the minimum energy input values as a function of Kp and as a function of B_z - V_{sw} , we note that the energy flux input for Kp = 0, 0+ is lower than that for low V_{sw} and weakly positive B_z . Possible explanations for this include: 1) the corresponding B_z/V_{sw} bin was too broad; 2) conditions additional to the instantaneous value of the solar wind speed and the z-component of the IMF are required to specify the condition of weakest precipitating particle energy flux input.




Fig. 1. Precipitating electron and ion hemispheric energy flux as a function of (a) Kp and (b) IMF B_z .

REFERENCES

1. D.D. Wallis and E.E. Budzinski, Empirical Models of Height-Integrated Conductivities, J. Geophys. Res. 86, 125 (1981)
2. R.W. Spiro, P.H. Reiff, and L.J. Maher, Jr., Precipitating Electron Energy Flux and Auroral Zone Conductances - An Empirical Model, J. Geophys. Res. 86, 8215 (1982)
3. D.A. Hardy, M.S. Gussenhoven, and E. Holeman, A Statistical Model of Auroral Electron Precipitation, J. Geophys. Res. 86, 4229 (1985)
4. J.C. Foster, J.M. Holt, R.G. Musgrove, and D.S. Evans, Solar Wind Dependencies of High-Latitude Convection and Precipitation, in: Solar Wind-Magnetosphere Coupling, ed. Y. Kamide and J.A. Slavin, Terra Scientific Publishing Company, Tokyo 1986, p. 477.
5. D.A. Hardy, M.S. Gussenhoven, D.H. Brautigam, A Statistical Model of Auroral Ion Precipitation, J. Geophys. Res., accepted for publication, 1988.
6. P.H. Reiff and J.G. Luhmann, Solar Wind Control of the Polar-Cap Voltage, Solar Wind-Magnetosphere Coupling, ed. Y. Kamide and J.A. Slavin, Terra Scientific Publishing Company, Tokyo 1986, p.453.
7. D.A. Hardy, L.K. Schmitt, M.S. Gussenhoven, F.J. Marshall, H.C. Yeh, T.L. Schumaker, A. Huber and J. Pantazis, Precipitating Electron and Ion Detectors (SSJ/4) for the Block 5D/Flight 6-10 DMSP Satellites: Calibration and Data Presentation, AFGL-TR-84-0314, Air Force Geophysics Laboratory, Hanscom AFB, MA (November 1984)



Accession For	
NTIS CRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	20