

U
SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

AD-A208 897

1b RESTRICTIVE MARKINGS

NA

3 DISTRIBUTION / AVAILABILITY OF REPORT

Distribution Unlimited

4. PERFORMING ORGANIZATION REPORT NUMBER(S)

Univ. of Puerto Rico #

5 MONITORING ORGANIZATION REPORT NUMBER(S)

NA

6a. NAME OF PERFORMING ORGANIZATION

Univ. of Puerto Rico

6b OFFICE SYMBOL
(If applicable)

NA

7a. NAME OF MONITORING ORGANIZATION

Office of Naval Research

6c. ADDRESS (City, State, and ZIP Code)

Department of Marine Sciences
Box 5000
Mayaguez, P.R. 00709

7b. ADDRESS (City, State, and ZIP Code)

800 N. Quincy St.
Arlington, Va. 22217-5000

8a. NAME OF FUNDING / SPONSORING
ORGANIZATION

Office of Naval Research

8b OFFICE SYMBOL
(If applicable)

ONR

9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

N00014-88-K-0131

8c. ADDRESS (City, State, and ZIP Code)

800 N. Quincy St.
Arlington, Va. 22217-5000

10. SOURCE OF FUNDING NUMBERS

PROGRAM
ELEMENT NO
61153N

PROJECT
NO
RR04106

TASK
NO
441h-008

WORK UNIT
ACCESSION NO

11. TITLE (Include Security Classification)

(U) Molecular Specificity of Adsorption of Biofilm Macromolecules and
Microbial Biofouling on Artificial Surfaces in the Sea

12. PERSONAL AUTHOR(S)

Tosteson, Thomas Robert and Yamamura, Yasuhiro (Ponce School of Medicine)

13a. TYPE OF REPORT

Progress

13b. TIME COVERED

FROM 2/89 TO 6/89

14. DATE OF REPORT (Year, Month, Day)

June 1, 1989

15. PAGE COUNT

3

16. SUPPLEMENTARY NOTATION

Immunoglobulin G

17. COSATI CODES

FIELD	GROUP	SUB-GROUP
06	03	

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

Adhesion Enhancing (AE) Macromolecules, AE Mono-
clonal Antibodies, Macromolecular Biofouling, Sea
Surfaces, AE Biofouling Antigens,

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

Hyperimmune mouse (IgG) made against microbial adhesion enhancement
macromolecules (MAEM) reacts specifically with plates coated with immune
chicken IgG. Very high titers (100,000) of hyperimmune mouse sera suggest
production of antibody(ies) against MAEM antigens, however, the ELISA assay
could not discriminate specific from apparent, nonspecific reactions.
D-galactose and elevated salt concentrations did not affect this binding.
The chicken and mouse may be making antibodies to different portions of
the same MAEM or to different MAEM antigens. To clarify this: (1) MAEM
antigens from the same source are being isolated independently by toad
and chicken immunoaffinity columns. (2) Adhesion enhancement activity
of the respective recovered antigens is being assessed. (3) MAEM antigens
recovered from the toad immunoaffinity columns are being examined for
their quantitative recovery and adhesion activity when they are reacted

20. DISTRIBUTION / AVAILABILITY OF ABSTRACT

☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT ☐ DTIC USERS

21. ABSTRACT SECURITY CLASSIFICATION

(U)

22a. NAME OF RESPONSIBLE INDIVIDUAL

Dr. Michael Marron

22b. TELEPHONE (Include Area Code)

202/696-4760

22c. OFFICE SYMBOL

ONR

DD Form 1473, JUN 86

Previous editions are obsolete

SECURITY CLASSIFICATION OF THIS PAGE

S/N 0102-LF-014-6603

A 89

6 12 134

#18. Macromolecular Sea Surfaces

- #19 ^{cont'd} with the immunoaffinity column made with chicken antibodies.
#4) Antigens recovered by the two types of columns will be used to produce hyperimmune mouse sera for reexamination of the ELISA specificity. Inhibitory effects of these sera on the adhesion advancement activity of MAEM antigens will be used to assess similarities among adsorbed MAEM antigens. (AW) 



Distribution List for Annual and Final Reports

1. Put a cover page (Form DD 1473) on your report and attach a copy of the distribution list. Mail one copy of the report to each person on the contractor subset list attached on which your name appears. The other subset list is for your information only. Please don't forget to attach this distribution list to your report - otherwise the folks below think they have mistakenly received the copy meant for the Molecular Biology Program and forward it to us.

2. Mail two copies to (include a DTIC Form 50 with these two copies too)
 Administrator
 Defense Technical Information Center
 Building 5, Cameron Station
 Alexandria, VA 22314

3. Mail one copy to each of the following:

(a) Dr. Michael Marron ONR Code 1141 Molecular Biology Program 800 N. Quincy Street Arlington, VA 22217-5000	(e) Director Chemical and Biological Sci Div Army Research Office P. O. Box 12211 Research Triangle Park, NC 27709
(b) Administrative Contracting Officer ONR Resident Representative (address varies - see copy of your grant)	(f) Life Sciences Directorate Air Force Office of Scientific Research Bolling Air Force Base Washington, DC 20332
(c) Director, Applied Research Directorate ONR Code 12 800 N. Quincy Street Arlington, VA 22217-5000	(g) Director Naval Research Laboratory Technical Information Div, Code 2627 Washington, DC 20375
(d) Director Office of Naval Technology Code 22 800 N. Quincy Street Arlington, VA 22217-5000	



Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification _____	
By _____	
Distribution/ _____	
Availability Codes _____	
Dist _____	Special _____

A-1

DATE: 1 JUNE, 1989

PROGRESS REPORT ON CONTRACT NO0014-88-K-0131

TITLE: The Molecular Specificity of Adsorption of Biofilm
Macromolecules and Accumulation of Microbial Biofouling
on Artificial Surfaces in the Sea.

PRINCIPAL INVESTIGATORS: T.R. Tosteson and Y. Yamamura. Department of Marine Sciences, University of Puerto Rico-Mayaguez and Microbiology Department, Ponce School of Medicine, Ponce, Puerto Rico.

RESEARCH OBJECTIVES: The diversity and variability of macromolecular components that mediate initial microbial attachment to surfaces in ambient seawater is being determined employing immunological techniques. Antisera against microbial adhesion enhancing macromolecules (MAEM) have been raised in chickens and the immunoglobulin-G (IgG) fraction utilized for isolation of crude MAEM from samples of cell free coastal seawater and marine microbial culture media by immunoaffinity chromatography. The objective of the present study is to produce monoclonal antibodies (MAbs) against MAEM and use these to chromatographically isolate and purify individual MAEM from mixtures of such components. Variability and diversity of MAEM produced by biofouling microorganisms, those found soluble in ambient seawater and on biofouled surfaces will be assessed employing a spectrum of MAEM MAbs. The specificity of the interactions of the various MAEM with glass and metallic surfaces will be assessed. Alterations in the physical characteristics of the test surfaces and changes in their susceptibility to microbial biofouling will be correlated with these interactions.

PROGRESS TO DATE: Microbial adhesion enhancing macromolecules (MAEM) have been isolated by immunoaffinity chromatography from samples of cell free coastal seawater and from the laboratory culture growth media of marine microorganisms recovered from biofouled surfaces. MAEM were recovered from these samples by immunoaffinity chromatography employing polyclonal antibodies against these macromolecules. Recovered antigens were pooled for use in the production of monoclonal antibodies (MAbs).

A major objective of this project was to standardize an immune peroxidase ELISA method suitable for mass screening of hybridomas produced against MAEM antigens. Hens were hyperimmunized by repeated subcutaneous inoculation with PS antigens and their eggs collected. Chicken IgG was isolated using polyethylene glycol and subsequently purified by repeated precipitation with ammonium sulfate. ST plates were individually coated with either normal or immune chicken IgG. Hyperimmune mouse IgG (but not normal mouse IgG) reacted (bound) specifically with immune chicken IgG coated plates, in the presence or absence of MAEM antigen. Thus, the ELISA method did not appear to be specific. The assay yielded a titer of approximately 100,000 for the hyperimmune mouse serum over that of the normal control. It is important to note that the hyperimmune mouse serum, which was a pool from four mice, consistently showed this property, despite the presence of D-galactose and other simple sugars, as well as elevated salt concentrations.

The success of our approach to this project heavily relies on the specificity of the ELISA assay. The very high titer of hyperimmune mouse serum suggests that immunization has produced antibody(ies) against MAEM antigens, however, the assay could not discriminate specific from apparent nonspecific

reactions. The specific interaction of immune chicken IgG and hyperimmune mouse sera suggests that either these two animals are making antibodies to different portions of the same MAEM or, more likely to different molecules found in the mixture of MAEM antigens originally recovered from seawater using polyclonal antibodies. MAEM antigens were originally recovered from seawater samples using immunoaffinity columns composed of toad antibodies. These antibodies were raised against the hydroxylapatite (HTP) purified fraction of high molecular weight components found in ambient seawater that specifically enhanced microbial adhesion to surfaces (Tosteson, T.R. *et al.*, Journal of Colloid and Interface Science 104:60-71, 1985). Antigens recovered from the toad antibody column were used to raise antibodies in the chicken in order to construct an additional, larger immunoaffinity column composed of chicken antibodies. Both of these columns, combined in series were used to accumulate MAEM antigens that were initially used in this project. The complex specificity found in the interaction of immune chicken IgG and hyperimmune mouse sera may result from differences in the responses of the toad, chicken and mouse to MAEM. Experiments are now in progress to resolve this problem.

The second project objective was to make and isolate hybridomas that produce MABs against individual PS antigens. To date, 28 hybridoma clones have been shown to produce reactive IgG, detectable using the immune chicken IgG coated plate method described above. These clones are stored frozen in liquid nitrogen, awaiting further refinement of the ELIZA method.

WORKPLAN (remainder of Year 2): The following studies are planned to resolve the ELIZA assay specificity problem: (1) MAEM antigens will be independently recovered from the same source (coastal seawater or microbial culture media) by toad and chicken immunoaffinity columns. The quantity and adhesion activity of the MAEM antigens recovered from the respective columns will be assessed. The quantity and activity of antigens initially recovered from the toad column will be analyzed using the immunoaffinity column made with chicken antibodies. (2) Antigens recovered by both types of columns will be used to produce hyperimmune mouse sera for reexamination of the ELIZA specificity. (3) Inhibitory effects of these sera on MAEM adhesion activity will be examined to compare recovered MAEM antigens. Similar inhibition studies will be conducted with the supernatant solutions recovered from hybridoma clones that produce reactive IgG (i.e. detectable by immune chicken IgG coated plates). (4) HTP purified MAEM will be used directly to produce hyperimmune mouse sera.

INVENTIONS: None.

PUBLICATIONS AND PRESENTATIONS: Molecular Specificity of Microbial Adhesion Enhancing Macromolecules. T.R. Tosteson and Yasuhiro Yamamura. Presented by Y. Yamamura at the Marine Biosurfaces Contractors Meeting, May 20-22, 1989. Hopkins Marine Station, Pacific Grove, California.

TRAINING ACTIVITIES: None.

AWARDS AND FELLOWSHIPS: None.