

2

AD-A210 079

OFFICE OF NAVAL RESEARCH

Contract N00014-87-K-0541

R & T Code 4133020

Technical Report No. 12

The Structure of the Mercury/Dimethylsulfoxide Interface
in the Presence of Tetraalkylammonium Perchlorates

by

W.R. Fawcett, R.F. Unterleitner, and A.J. Motheo

Prepared for Presentation

at

The Electrochemical Society Meeting
Los Angeles, CA , May 1989

May 1, 1989

DTIC
ELECTE
JUN 20 1989
D

Reproduction in whole or in part is permitted
for any purpose of the United States Government

"This document has been approved for public release
and sale; its distribution is unlimited"

89 6 20 013

REPORT DOCUMENTATION PAGE		1. REPORT NO. 12	2.	3. Recipient's Accession No.
4. Title and Subtitle The Structure of the Mercury/Dimethylsulfoxide Interface in the Presence of Tetraalkylammonium Perchlorates.			5. Report Date May 1, 1989	
7. Author(s) W.R. Fawcett, R.F. Unterleitner and A.J. Motheo			8. Performing Organization Report No.	
9. Performing Organization Name and Address Department of Chemistry University of California Davis, CA 95616			10. Project/Task/Work Unit No.	
11. Sponsoring Organization Name and Address Office of Naval Research 800 N. Quincy Arlington, VA 22217-5000			12. Type of Report & Period Covered	
13. Supplementary Notes Prepared for Presentation at: The Electrochemical Society Meeting, Los Angeles, CA, May 1989				
14. Abstract (Limit 200 words) Differential capacity against potential data were obtained in dimethylsulfoxide solutions containing tetraalkylammonium perchlorates. It is shown that perchlorate anion is weakly adsorbed in the vicinity of the potential of zero charge on the basis of the dependence of the experimental capacity on electrolyte concentration for tetrabutylammonium perchlorate solutions. The influence of cation size on inner layer capacity at more negative potentials, where specific adsorption is absent, is discussed and compared with that found in other aprotic solvents. <i>MOLECULAR STRUCTURE</i>				
17. Document Analysis a. Descriptors b. Identifiers/Open-Ended Terms c. COSATI Field/Group			Accession For NTIS GRA&I ☒ DTIC TAB ☐ Unannounced ☐ Justification ☐	
			By Distribution/ Availability Codes Avail and/or Special	
			Dist A-1	
18. Availability Statement			19. Security Class (This Report) Unclassified.	21. No. of Pages
			20. Security Class (This Page)	22. Price

75-Word Abstract Form

Extended Abstract must be submitted with the 75-Word Abstract by December 1, 1988

Los Angeles, California—May 7-12, 1989

Submit to: The Electrochemical Society, Inc.
10 South Main Street, Pennington, NJ 08534-2896
With a copy to the Organizing Chairman

Abstract No.
(to be assigned by the Society)

Schedule for General Session
(Title of Symposium)

Sponsored by Physical Electrochemistry
(Division/Group)

Title of paper The Structure of the Mercury/Dimethylsulfoxide Interface in the
Presence of Tetraalkylammonium Perchlorates

Authors (Underline name of author presenting paper.) W. Ronald Fawcett, Rolf F. Unterleitner,
and Artur J. Motheo

Business affiliation and address Department of Chemistry
University of California

..... Davis, CA 95616 (916) 752-1105
(State or Country) (ZIP Code) (Telephone No.)

(Type abstract in this area—double-spaced.)

Differential capacity against potential data were obtained in dimethylsulfoxide solutions containing tetraalkylammonium perchlorates. It is shown that perchlorate anion is weakly adsorbed in the vicinity of the potential of zero charge on the basis of the dependence of the experimental capacity on electrolyte concentration for tetrabutylammonium perchlorate solutions. The influence of cation size on inner layer capacity at more negative potentials, where specific adsorption is absent, is discussed and compared with that found in other aprotic solvents.

Do you require any audiovisual equipment?

- ☐ 35 mm (2 x 2 in.) slide projector
☒ Overhead projector
☐ Specify other (subject to availability and cost)

Has the information in this abstract been presented verbally, submitted for publication, or published?

☐ Yes ☐ No

If the answer is yes, please provide the reference (except in the case of invited review presentations).

Is a full length paper on this work to be submitted for Society Journal publication? ☐ Yes ☐ No

Papers presented before a Society technical meeting become the property of the Society and may not be published elsewhere without written permission of the Society. Papers presented at Society technical meetings must be authored by a member or sponsored by an active member.

W.R. Fawcett and A.J. Motheo

Insert name of Society member author or sponsor

The Structure of the Mercury/Dimethylsulfoxide Interface in the Presence of Tetraalkylammonium Perchlorate.

W. Ronald Fawcett, Rolf F. Unterleimer, and Arthur J. Motheo*

Department of Chemistry, University of California, Davis, CA 95616

The structure of the mercury/non-aqueous solution interface has been studied in the presence of a number of aprotic solvents [1, 2]. In the case of dimethylsulfoxide (DMSO), Payne [3] studied the behavior of alkali metal salts with a variety of anions and observed a strong resemblance to the corresponding aqueous systems. On the other hand, the properties of the mercury/DMSO solution interface have not been investigated in the presence of tetraalkylammonium salts in spite of the popularity of these electrolytes in non-aqueous electrochemistry. In this paper, we report a study of double layer structure in the presence of tetraethylammonium perchlorate (TEAP), and the corresponding tetrapropylammonium (TPAP) and tetrabutylammonium salts (TBAP), assess the role of perchlorate anion adsorption, and compare the inner layer capacity curves with those found in other aprotic solvents such as acetonitrile [4] and dimethylformamide [5].

Differential capacity against potential data were obtained using an a.c. impedance bridge with potentiostatic control of the mercury electrode. The potential of zero charge was determined using a streaming mercury electrode. In the case of TBAP, capacity data were collected at ten concentrations in the range 0.013 to 0.10 M. The potential of the Hg electrode was measured with respect to a Ag/Ag⁺ reference electrode. The tetraalkylammonium salts and the DMSO were purified using well established procedures.

Differential capacity against potential data for the Hg/TBAP system in DMSO are shown in Fig. 1, the potential scale being referenced to the potential of zero charge (p.z.c.) Diffuse layer minima coincident with the p.z.c. are clearly defined at lower salt concentrations. According to the Gouy-Chapman-Stern model of the double layer, the reciprocal of the double layer capacity at the p.z.c., $1/C_0$ should be linear in the reciprocal of the square root of the electrolyte concentration $1/c_3^{1/2}$ if specific adsorption is absent, the relationship being [4]

$$1/C_0 = 1/C_{i0} + 1/f \Theta c_3^{1/2} \quad (1)$$

C_{i0} is the inner layer capacity at the p.z.c., $\Theta^2 = 2 RT\epsilon\epsilon_0$ where ϵ is the dielectric constant of DMSO, and $f = F/RT$. Although the relationship is linear (Fig. 2), the slope is not equal to $1/f \Theta$ suggesting that weak specific adsorption is present. The presence of specific adsorption can also be tested on the basis of plots of $1/C$ against $1/C_d$ at constant electrode charge density

where C_d is the diffuse layer capacity estimated on the basis of Gouy-Chapman theory (see Fig. 3). According to these plots specific adsorption disappears at negative charge densities indicating that the weakly adsorbing species is the perchlorate ion.

As expected, the shape of the capacity-potential curves depends on the nature of the tetraalkylammonium cation at negative charge densities. This dependence is discussed in terms of the relative influence of the cation on the dielectric constant of the inner layer and on its thickness. Finally, the data in DMSO are compared with those obtained earlier in acetonitrile [4] and dimethylformamide [5].

Acknowledgements

This work was supported by the Office of Naval Research, Washington. A.J.M. thanks the Conselho Nacional de Desenvolvimento Científico e Tecnológico, Brazil for a maintenance grant.

References

1. R. Payne, Adv. Electrochem. Electrochem. Eng., **7**, 1 (1970).
2. W.R. Fawcett, Israel J. Chem., **18**, 3 (1979).
3. R. Payne, J. Am. Chem. Soc., **89**, 489 (1967).
4. W.R. Fawcett and R.O. Loutfy, Can. J. Chem., **51**, 230 (1973).
5. W.R. Fawcett, B.M. Ikeda, and J.B. Sellan, Can. J. Chem., **57**, 2268 (1979).

*Permanent address - Instituto de Física e Química de São Carlos, USP, São Carlos, SP 13560, Brazil.




Fig. 1 Differential capacity against potential for the Hg/DMSO interface with varying concentrations of TBAP: 0.013(1), 0.032(2), 0.056(3) and 0.10 M(4).




Fig. 2 Reciprocal capacity at the p.z.c. $1/C_0$ plotted against the reciprocal of the square root of salt concentration for the Hg/DMSO interface with TBAP as electrolyte.




Fig. 3 Reciprocal capacity $1/C$ against the reciprocal of the diffuse layer capacity $1/C_d$ for constant electrode charge density as indicated. The lines are drawn with unit slope.