

AD-A210 132

THE FILE COPY

②

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0043

TECHNICAL REPORT No. 104

Squeezing of Atomic Variables in the One-Photon and Two-Photon
Jaynes-Cummings Model

by

Xiao-shen Li, D. L. Lin and Thomas F. George

Prepared for Publication

in

Physical Review A

Departments of Chemistry and Physics
State University of New York at Buffalo
Buffalo, New York 14260

June 1989

Reproduction in whole or in part is permitted for any purpose of the
United States Government.

This document has been approved for public release and sale;
its distribution is unlimited.

DTIC
ELECTE
JUL 17 1989
S B D
Cb

89

7

17

102

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/89/TR-104			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION			
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260		7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract N00014-86-K-0043			
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		10. SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Squeezing of Atomic Variables in the One-Photon and Two-Photon Jaynes-Cummings Model					
12. PERSONAL AUTHOR(S) Xiao-shen Li, D. L. Lin and Thomas F. George					
13a. TYPE OF REPORT	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, Month, Day) June 1989	15. PAGE COUNT 17		
16. SUPPLEMENTARY NOTATION Prepared for publication in Physical Review A					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	JAYNES-CUMMINGS MODEL > ONE-PHOTON PROCESS,		
			ATOMIC DIPOLE MOMENT, TWO-PHOTON PROCESS.		
			SQUEEZING QUANTUM MECHANICAL STUDY		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Squeezing of the atomic dipole moment in the one-photon and two-photon Jaynes-Cummings model is investigated with different initial conditions. Almost perfect squeezing is found in certain cases.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT ☒ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT. ☐ DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE (Include Area Code) (202) 696-4410	22c. OFFICE SYMBOL	

Physical Review A, in press

Squeezing of atomic variables in the one-photon and two-photon
Jaynes-Cummings model

Xiao-shen Li
Center for Theoretical Physics
Chinese Center of Advanced Science and Technology (World Laboratory)
P. O. Box 8730
Beijing 100080, People's Republic of China

and

Shanghai Institute of Metallurgy
Chinese Academy of Sciences
Shanghai, People's Republic of China

D. L. Lin and Thomas F. George
Department of Physics & Astronomy
239 Fronczak Hall
State University of New York at Buffalo
Buffalo, New York 14260

Abstract

Squeezing of the atomic dipole moment in the one-photon and two-photon
Jaynes-Cummings model is investigated with different initial conditions.
Almost perfect squeezing is found in certain cases.

PACS Numbers: 32.80.-t, 42.5.Kb

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

I. Introduction

It is by now well known that squeezed light has potential applications in low-noise communications and high-precision measurements involving light.^{1,2} The Jaynes-Cummings (JC) model^{3,4} is one of the possible nonlinear optical models capable of creating squeezed states.⁵ A number of calculations of field squeezing have been made in recent years. In the standard JC model, the maximum field squeezing was found to be 20%,⁵ but in a recent paper it was argued that this squeezing could be 100%.⁶ When the initial coherent state of the field is replaced by the vacuum state, the maximum squeezing is increased to 25%.⁷ More squeezing is found in many other cases of the generalized JC model. The maximum squeezing is 42% for two-level two-photon,⁸ 57% for two-level three-photon,⁹ 31% for three-level one-mode,¹⁰ 36% for four-level one-mode,¹¹ and 52% for ten-level one-mode¹² cases. However, little attention has been paid to the squeezing of the atomic variables. The squeezing of the atomic dipole operators for a two-level atom driven by a classical field¹³ has been investigated. A general relation has been established between the coherent states of the SU(2) and SU(1,1) Lie algebra and possible reduction of fluctuations in angular momentum.¹⁴ For the special case of a vacuum initial field mode, the relation between the atomic variable squeezing and the field squeezing has been examined.¹⁵ Only recently, a quantum mechanical study devoted to the squeezing of the atomic dipole moment has been carried out,¹⁶ where it is found that a maximum squeezing of 60% can be reached.

In this note we calculate the squeezing of the atomic dipole moment in the one-photon and two-photon JC model with different initial conditions. It is found that the atomic variables generally possess stronger squeezing than the field under the same conditions. In particular circumstances, a nearly pure squeezing state can be achieved. In the standard JC model, a one-mode

cavity field with frequency Ω couples to a two-level atom through the well-known Hamiltonian in the rotating-wave approximation,

$$H = \frac{1}{2} \hbar \omega \sigma_3 + \hbar \Omega a^\dagger a + \hbar \lambda (a^\dagger \sigma^- + a \sigma^+) \quad , \quad (1)$$

where $a^\dagger(a)$ and Ω are the creation (annihilation) operator and frequency of the cavity field, respectively, and λ is the coupling constant. The two-level atom with transition frequency ω is described by the Pauli raising and lowering operators $\sigma^\pm$ and the inversion operator $\sigma_3 = \sigma^+ \sigma^- - \sigma^- \sigma^+$. The dispersive and absorptive components of the slowly-varying atomic dipole¹⁷ can be written as

$$\sigma_1 = \frac{1}{2} (\sigma^+ e^{-i\omega t} + \sigma^- e^{i\omega t}) \quad (2)$$

and

$$\sigma_2 = \frac{1}{2i} (\sigma^+ e^{-i\omega t} - \sigma^- e^{i\omega t}) \quad , \quad (3)$$

respectively. They obey the commutation relation

$$[\sigma_1, \sigma_2] = \frac{i}{2} \sigma_3 \quad (4)$$

and the corresponding uncertainty relation

$$(\Delta \sigma_1)^2 (\Delta \sigma_2)^2 \geq \frac{1}{16} \langle \sigma_3 \rangle^2 \quad . \quad (5)$$

The atomic state is said to be squeezed when σ_1 or σ_2 satisfies the relation

$$(\Delta\sigma_i)^2 < \frac{1}{4} |\langle\sigma_3\rangle|, \quad i = 1, 2. \quad (6)$$

Since

$$(\Delta\sigma_1)^2 = \frac{1}{4} - (\text{Re}\langle\sigma^-\rangle e^{i\omega t})^2 \quad (7a)$$

$$(\Delta\sigma_2)^2 = \frac{1}{4} - (\text{Im}\langle\sigma^-\rangle e^{i\omega t})^2, \quad (7b)$$

the condition described by Eq. (6) can be rewritten as

$$S_1 = \frac{1 - 4(\text{Re}\langle\sigma^-\rangle e^{i\omega t})^2}{|\langle\sigma_3\rangle|} < 1 \quad (8a)$$

or

$$S_2 = \frac{1 - 4(\text{Im}\langle\sigma^-\rangle e^{i\omega t})^2}{|\langle\sigma_3\rangle|} < 1 \quad (8b)$$

for squeezing in the dispersive or absorptive component of the dipole. In what follows we shall investigate this squeezing effect by using different initial conditions for the atom and the cavity field.

II. Squeezing in the JC model

We denote by $|+\rangle$ and $|-\rangle$ the excited and ground states of the atom and by $|n\rangle$ the Fock states of the field. First we consider the initial condition

that the atom is in the excited state and the field in a two-photon coherent state or squeezed state,^{1,2,18}

$$|\alpha, r\rangle = \sum_n F(n) |n\rangle \quad (9)$$

$$F(n) = (n! \cosh r)^{-1/2} \left(\frac{\beta}{2 \cosh r} \right)^n Y^{-n} H_n(Y) \exp\left[-\frac{\beta^2}{2}(1 - \tanh r)\right] \quad (10)$$

where $\beta = \alpha e^r$ and $Y = \beta(2 \cosh r \sinh r)^{-1/2}$, and we have chosen the squeezing parameter r to be real.¹⁹ The initial mean photon number is given by $\bar{n} = |\alpha|^2 + \sinh^2 r$, so that the initial state vector for the atomic-field system can be written as

$$|\psi(0)\rangle = \sum_n F(n) |+, n\rangle \quad (11)$$

At a time $t > 0$, the state vector in the interaction picture is found from the JC model Hamiltonian (1) to be

$$|\psi(t)\rangle = \sum_n F(n) [A(n, t) |+, n\rangle + B(n+1, t) |-, n+1\rangle] \quad (12)$$

where

$$A(n, t) = e^{-i\Delta t/2} (\cos \mu t + i \frac{\Delta}{2} \sin \mu t / \mu) \quad (13)$$

$$B(n+1, t) = -i \frac{Y}{\mu} e^{-i\Delta t/2} \sin \mu t \quad (14)$$

with $V = \lambda\sqrt{n+1}$, $\mu^2 = V^2 + \Delta^2/4$ and $\Delta = \Omega - \omega$. From Eqs. (12)-(14), we obtain the expectation values of the atomic operators:

$$\begin{aligned} \langle \sigma^- \rangle e^{i\omega t} &= \langle \psi(t) | \sigma^- e^{i\omega t} | \psi(t) \rangle \\ &= \sum_{n=0}^{\infty} F^*(n) F(n+1) B^*(n+1) A(n+1) \end{aligned} \quad (15)$$

$$\langle \sigma_3 \rangle = \sum_{n=0}^{\infty} |F(n)|^2 [|A(n)|^2 - |B(n+1)|^2] \quad (16)$$

The time evolutions of S_1 and S_2 are calculated for the case of off-resonant excitation with medium strength. The units employed throughout this paper are λ for energy and λ^{-1} for time. The results are presented in Fig. 1. It is observed that σ_1 and σ_2 are squeezed alternatively. Roughly speaking, the squeezing parameters S_1 and S_2 oscillate at about the same frequency but out of phase. A more careful analysis reveals that both are composed of two oscillations with different frequencies. The maximum squeezing in this case is approximately 25% for both S_1 and S_2 .

For the case of on-resonance excitation, $\langle \sigma_1 \rangle$ is zero but its fluctuation is still nonvanishing because of quantum mechanical effects. Figure 2 shows the time evolution of S_2 for the on-resonance excitation by a squeezed and a coherent light field. A comparison of (a) and (b) in Fig. 2 shows that squeezing occurs only for a short time after the interaction is turned on when the initial field is in a coherent state. The maximum squeezings in these cases are, however, about the same, namely, 65% and 68% for the coherent and squeezed excitations, respectively. The role played by

the detuning can be clearly seen by comparing Figs. 1(b) and 2(a). Evidently, a larger detuning implies weaker coupling between the atom and the field and hence less squeezing. Although the results shown are obtained for the same $\bar{n}$, we have found from our numerical study that in general a larger exciting strength or larger $\bar{n}$ results in stronger squeezing effects.

Next we consider a vacuum initial field in the JC model. The atom is injected into the field in a coherent superposition of excited and ground states,⁷

$$|\psi(0)\rangle = e^{i\phi} \cos\theta |-,0\rangle + \sin\theta |+,0\rangle . \quad (17)$$

From the Hamiltonian (1) and Eq. (17), the state vector at time $t > 0$ can be written as

$$|\psi(t)\rangle = C_0^- |-,0\rangle + C_1^- |-,1\rangle + C_0^+ |+,0\rangle , \quad (18)$$

where

$$C_1^-(t) = -i \sin\theta \sin\lambda t \quad (19)$$

$$C_0^+(t) = \sin\theta \cos\lambda t \quad (20)$$

$$C_0^-(t) = e^{i\phi} \cos\theta . \quad (21)$$

Using the same procedure as outlined above, we can write

$$\langle \sigma^- \rangle e^{i\omega t} = \langle \psi(t) | \sigma^- e^{i\omega t} | \psi(t) \rangle = C_0^{-*} C_0^+ - \frac{1}{2} e^{-i\phi} \sin 2\theta \cos \lambda t \quad (22)$$

and

$$\langle \sigma_3 \rangle = 1 - 2\sin^2 \theta \cos^2 \lambda t \quad . \quad (23)$$

Inserting Eqs. (22) and (23) into (8), we find the squeezing parameters as

$$S_1 = (1 - \sin^2 2\theta \cos^2 \phi \cos^2 \lambda t) / |1 - 2\sin^2 \theta \cos^2 \lambda t| \quad , \quad (24)$$

$$S_2 = (1 - \sin^2 2\theta \sin^2 \phi \cos^2 \lambda t) / |1 - 2\sin^2 \theta \cos^2 \lambda t| \quad . \quad (25)$$

From these two equations one can see that when the atom starts in a purely excited state, i.e., $\theta = \frac{\pi}{2}$, there is no squeezing effect. It is apparent that maximum squeezing occurs when $\lambda t = k\pi$ ($k = 0, 1, 2, \dots$) and $\phi = k\pi$ (for σ_1) or $(k + \frac{1}{2})\pi$ (for σ_2). We have, under these conditions,

$$S_1 = (1 - \sin^2 2\theta) / |1 - 2\sin^2 \theta| = |\cos 2\theta| \quad , \quad (26)$$

where $\theta = \frac{1}{2}(k + \frac{1}{2})\pi$. Equation (20) yields

$$S_1 = 0 \quad . \quad (27)$$

This means that the maximum squeezing of 100% is possible.

III. Squeezing in two-photon processes

A simple model for atom-radiation interaction that can be solved analytically has recently been proposed.⁷ It involves two-photon Raman

coupling between two atomic states $|+\rangle$ and $|-\rangle$ degenerate in energy. The Hamiltonian for this model is therefore

$$H = \hbar\Omega a^\dagger a + \hbar\lambda a^\dagger a(\sigma^- + \sigma^+) \quad , \quad (28)$$

where we have set the atomic transition frequency to be zero because of the degeneracy. It has been shown that the system has perfect revival.⁷ Here we shall study the squeezing of the atomic dipole operator in this model.

We assume that the atom starts in the initial state $|+\rangle$ and the field is in a coherent state $|\alpha\rangle$,

$$|\alpha\rangle = \sum_n G(n) |n\rangle \quad (29)$$

$$G(n) = \frac{\alpha^n}{\sqrt{n!}} e^{-\bar{n}/2} \quad , \quad (30)$$

where $\bar{n} = |\alpha|^2$ is the initial mean photon number of the field. The initial state vector then takes the form

$$|\psi(0)\rangle = \sum_n G(n) |+, n\rangle \quad . \quad (31)$$

At any time $t > 0$, the state vector can be written as

$$|\psi(t)\rangle = \sum_{n=0}^{\infty} G(n) [D^+(n) |+, n\rangle + D^-(n) |-, n\rangle] \quad , \quad (32)$$

where

$$D^+(n,t) = \cos(\lambda nt) \quad (33a)$$

$$D^-(n,t) = -i \sin(\lambda nt) \quad (33b)$$

From Eqs. (32) and (33), one can easily find the mean value of the atomic dipole moment as

$$\langle \sigma^- \rangle = \frac{i}{2} \sum_{n=0}^{\infty} e^{-\bar{n}} \frac{\bar{n}^n}{n!} \sin(2\lambda nt) = \frac{i}{2} \exp[-2\bar{n} \sin^2(\lambda t)] \sin[\bar{n}(\sin 2\lambda t)] \quad (34)$$

and the inversion

$$\langle \sigma_3 \rangle = \exp(-2\bar{n} \sin^2 \lambda t) \cos(\bar{n} \sin 2\lambda t) \quad (35)$$

It is clear that the mean value of the dispersive component for the atomic dipole fluctuates around zero. Hence there is no squeezing in this component. As for the squeezing in the other component σ_2 , we find from Eqs. (8), (34) and (35)

$$S_2 = \frac{1 - \exp(-4\bar{n} \sin^2 \lambda t) \sin^2(\bar{n} \sin 2\lambda t)}{\exp(-2\bar{n} \sin^2 \lambda t) |\cos(\bar{n} \sin 2\lambda t)|} \quad (36)$$

To find the maximum magnitude of squeezing, we may consider the case where $\bar{n} \gg 1$ but $\bar{n}(\lambda t)^2 \ll 1$. Thus we have

$$S_2 \approx \frac{1 - [1 - 4\bar{n}(\lambda t)^2] \sin^2(2\bar{n}\lambda t)}{|\cos(2\bar{n}\lambda t)|}$$

$$= \frac{1 - (1 - \bar{n}^{-1}x) \sin^2 x}{|\cos x|}, \quad (37)$$

where $x = 2\bar{n}\lambda t$. When x is in the vicinity of $\frac{\pi}{2}$, i.e., $x = \frac{\pi}{2} - \epsilon$ with $\frac{1}{\bar{n}} \ll \epsilon \ll 1$, we may expand $\sin x$ and $\cos x$ in a Taylor series around $\frac{\pi}{2}$ to the second term. Hence, for $\lambda t \approx \frac{\pi}{4} \bar{n}^{-1}$ we have

$$S_2 \approx \frac{\pi^2}{4} (\bar{n}\epsilon)^{-1}. \quad (38)$$

Equation (38) implies that for strong excitation, S_2 is very small and the maximum magnitude of squeezing is nearly 100%.

Acknowledgments

This research was supported in part by the National Science Foundation under Grant CHE-8620274, the Office of Naval Research and the Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract F49620-86-C-0009. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon.

References

1. H. P. Yuen, Phys. Rev. A 13, 2226 (1976).
2. D. F. Walls, Nature 306, 141 (1983).
3. E. T. Jaynes and F. W. Cummings, Proc. IEEE 51, 89 (1963).
4. J. H. Eberly, N. B. Narozhny, J. J. Sanchez-Mondragon, Phys. Rev. Lett. 44, 1323 (1980).
5. P. Meystre and M. S. Zubairy, Phys. Lett. 89A, 390 (1982).
6. J. R. Kukliński and J. L. Madajczyk, Phys. Rev. A 37, 3175 (1988).
7. P. L. Knight, Phys. Scr. T12, 51 (1986).
8. G. Compagno, J. S. Peng and F. Persico, Opt. Commun. 57, 415 (1986).
9. A. S. Shumovsky, F. Le Kien and E. I. Abskenderov, Phys. Lett. 124A, 351 (1987).
10. S. Y. Zhu, Z. D. Liu and X. S. Li, Phys. Lett. 128A, 89 (1988).
11. Z. D. Liu, S. Y. Zhu and X. S. Li, J. Mod. Opt., in press.
12. F. L. Li, X. S. Li and D. L. Lin, unpublished.
13. D. F. Walls and P. Zoller, Phys. Rev. Lett. 47, 709 (1981).
14. K. Wodkiewicz and J. H. Eberly, J. Opt. Soc. Am. B 2, 458 (1985).
15. K. Wódkiewicz, P. L. Knight, S. J. Buckle and S. M. Barnett, Phys. Rev. A 35, 2567 (1987).
16. X. S. Li, D. L. Lin, T. F. George and Z. D. Liu, Phys. Rev. A, in press.
17. N. B. Narozhny, J. J. Sanchez-Mondragon and J. H. Eberly, Phys. Rev. A 23, 236 (1981).
18. R. Loudon and P. L. Knight, J. Mod. Opt. 34, 709 (1987).
19. G. J. Milburn and D. F. Walls, Phys. Rev. A 27, 392 (1983).

Figure Captions

1. Time evolution of the squeezing parameters (a) S_1 and (b) S_2 calculated for the case of off-resonance excitation. The parameters used are $\bar{n} = 100$, $r = 1$ and $\Delta = 50$.
2. Time evolution of S_2 for the case of on-resonance excitation by a (a) squeezed ($r = 1$) and (b) coherent ($r = 0$) field with $\bar{n} = 100$.







Fig. 1(a)



(b)

Fig. 2



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. C. B. Harris
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. H. Weaver
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. F. Kutzler
Department of Chemistry
Box 5055
Tennessee Technological University
Cookeville, Tennessee 38501

Dr. A. Reisman
Microelectronics Center of North Carolina
Research Triangle Park, North Carolina
27709

Dr. D. DiLella
Chemistry Department
George Washington University
Washington D.C. 20052

Dr. M. Grunze
Laboratory for Surface Science and
Technology
University of Maine
Orono, Maine 04469

Dr. R. Reeves
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. J. Butler
Naval Research Laboratory
Code 6115
Washington D.C. 20375-5000

Dr. Steven M. George
Stanford University
Department of Chemistry
Stanford, CA 94305

Dr. L. Interante
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Mark Johnson
Yale University
Department of Chemistry
New Haven, CT 06511-8118

Dr. Irvin Heard
Chemistry and Physics Department
Lincoln University
Lincoln University, Pennsylvania 19352

Dr. W. Knauer
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. K.J. Klaubunde
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. Murday
Naval Research Laboratory
Code 6170
Washington, D.C. 20375-5000

Dr. J. B. Hudson
Materials Division
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Theodore E. Madey
Surface Chemistry Section
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

Dr. J. E. Demuth
IBM Corporation
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. M. G. Lagally
Department of Metallurgical
and Mining Engineering
University of Wisconsin
Madison, Wisconsin 53706

Dr. R. P. Van Duyne
Chemistry Department
Northwestern University
Evanston, Illinois 60637

Dr. J. M. White
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. D. E. Harrison
Department of Physics
Naval Postgraduate School
Monterey, California 93940

Dr. R. L. Park
Director, Center of Materials
Research
University of Maryland
College Park, Maryland 20742

Dr. W. T. Peria
Electrical Engineering Department
University of Minnesota
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson
Department of Metallurgy and
Materials Science
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. S. Sibener
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Arnold Green
Quantum Surface Dynamics Branch
Code 3817
Naval Weapons Center
China Lake, California 93555

Dr. A. Wold
Department of Chemistry
Brown University
Providence, Rhode Island 02912

Dr. S. L. Bernasek
Department of Chemistry
Princeton University
Princeton, New Jersey 08544

Dr. W. Kohn
Department of Physics
University of California, San Diego
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Richard Colton
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Dan Pierce
National Bureau of Standards
Optical Physics Division
Washington, D.C. 20234

Dr. R. Stanley Williams
Department of Chemistry
University of California
Los Angeles, California 90024

Dr. R. P. Messmer
Materials Characterization Lab.
General Electric Company
Schenectady, New York 22217

Dr. Robert Gomer
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Ronald Lee
R301
Naval Surface Weapons Center
White Oak
Silver Spring, Maryland 20910

Dr. Paul Schoen
Code 6190
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. John T. Yates
Department of Chemistry
University of Pittsburgh
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene
Code 5230
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. L. Kesmodel
Department of Physics
Indiana University
Bloomington, Indiana 47403

Dr. K. C. Janda
University of Pittsburgh
Chemistry Building
Pittsburg, PA 15260

Dr. E. A. Irene
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Adam Heller
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton SO9 5NH
UNITED KINGDOM

Dr. H. Tachikawa
Chemistry Department
Jackson State University
Jackson, Mississippi 39217

Dr. John W. Wilkins
Cornell University
Laboratory of Atomic and
Solid State Physics
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis
Department of Physics
University of California
Irvine, California 92664

Dr. D. Ramaker
Chemistry Department
George Washington University
Washington, D.C. 20052

Dr. J. C. Hemminger
Chemistry Department
University of California
Irvine, California 92717

Dr. T. F. George
Chemistry Department
University of Rochester
Rochester, New York 14627

Dr. G. Rubloff
IBM
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. Horia Metiu
Chemistry Department
University of California
Santa Barbara, California 93106

Dr. W. Goddard
Department of Chemistry and Chemical
Engineering
California Institute of Technology
Pasadena, California 91125

Dr. P. Hansma
Department of Physics
University of California
Santa Barbara, California 93106

Dr. J. Baldeschwieler
Department of Chemistry and
Chemical Engineering
California Institute of Technology
Pasadena, California 91125

Dr. J. T. Keiser
Department of Chemistry
University of Richmond
Richmond, Virginia 23173

Dr. R. W. Plummer
Department of Physics
University of Pennsylvania
Philadelphia, Pennsylvania 19104

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 41106

Dr. N. Winograd
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. Roald Hoffmann
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. A. Steckl
Department of Electrical and
Systems Engineering
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. G.H. Morrison
Department of Chemistry
Cornell University
Ithaca, New York 14853