

AD-A211 065

2

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE				Form Approved OMB No 0704-0188	
1 REPORT SECURITY CLASSIFICATION U			1b RESTRICTIVE MARKINGS NA		
2 SECURITY CLASSIFICATION AUTHORITY NA			3 DISTRIBUTION/AVAILABILITY OF REPORT Distribution/Limited		
4 DECLASSIFICATION/DOWNGRADING SCHEDULE NA					
PERFORMING ORGANIZATION REPORT NUMBER(S) Chemistry University of Colorado			5 MONITORING ORGANIZATION REPORT NUMBER(S) NA		
5 NAME OF PERFORMING ORGANIZATION University of Colorado		6b OFFICE SYMBOL (If applicable) NA	7a NAME OF MONITORING ORGANIZATION Office of Naval Research		
6 ADDRESS (City, State, and ZIP Code) Boulder, Colorado 80309			7b ADDRESS (City, State, and ZIP Code) 800 N. Quincy Street Arlington, VA 22217-5000		
8a NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b OFFICE SYMBOL (If applicable) ONR	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N000 14-88-K-0353		
8c ADDRESS (City, State, and ZIP Code) 800 N. Quincy Street Arlington, VA 22217-5000			10 SOURCE OF FUNDING NUMBERS		
PROGRAM ELEMENT NO 61153N	PROJECT NO RR04106	TASK NO 4412047	WORK UNIT ACCESSION NO -		
11 TITLE (Include Security Classification) Chemical and Biochemical Properties of the Iron Mineral Core of Mammalian Ferritin.					
12 PERSONAL AUTHOR(S) Watt, Gerald D.					
13a TYPE OF REPORT Annual		13b TIME COVERED FROM 6/1/88 TO 6/1/89		14 DATE OF REPORT (Year, Month, Day) 6/5/89	
15 PAGE COUNT 3					
16 SUPPLEMENTARY NOTATION					
17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Ferritin, electron transfer, redox reactions biomineral		
FIELD 06	GROUP 03	SUB-GROUP -			
19 ABSTRACT (Continue on reverse if necessary and identify by block number) Both apo and holo ferritin bind Fe^{+} as well as other metal ions (Cu^{2+} , Zn^{2+} and Mn^{2+}) under anaerobic conditions as a function of pH. Apo ferritin binds 8 Fe^{2+} at protein sites whereas holo binds large numbers of Fe^{2+} on its mineral core surface. Holo ferritin undergoes reduction at its $FeOOH$ mineral core forming a Fe^{2+} mineral phase. Apo ferritin also undergoes redox reactions presumable at some amino acid cite. Electron transfer reactions occur readily in the $FeOOH$ core indicating the mineral core has relatively high electrical conductivity.					
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT ☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT ☐ DTIC USERS			21 ABSTRACT SECURITY CLASSIFICATION U		
22a NAME OF RESPONSIBLE INDIVIDUAL Michael Marron			22b TELEPHONE (Include Area Code) 202/696-4760		22c OFFICE SYMBOL ONR

DTIC
ELECTE
AUG 09 1989

39 8 07 105

ANNUAL REPORT: Contract N00014-88-K-0953
PRINCIPAL INVESTIGATOR: Gerald D. Watt
CONTRACTOR: The University of Colorado
CONTRACT TITLE: Chemical and Biochemical Properties of
the Iron Mineral Core of Mammalian
Ferritin
START DATE: 6/1/88

RESEARCH OBJECTIVES: To evaluate the chemical composition of the iron core of mammalian ferritin with regard to: 1) binding of metal ions; 2) electrochemical energy storage and 3) electron transfer reactions.

PROGRESS (YEAR 1)

Metal Ion Binding. Holo mammalian ferritin contains up to 4500 iron atoms within its hollow protein interior (80A in diameter) in the form of an FeOOH mineral core. Removal of this iron core by reduction and Fe(II) chelation forms apo ferritin with an empty, hollow interior. We have shown that both apo and holo ferritin avidly bind Fe²⁺ under anaerobic conditions in a pH dependent binding process. For holo ferritin, the binding occurs at the FeOOH mineral surface while for apo ferritin, Fe²⁺ binds at 8 protein sites. Cu²⁺, Zn²⁺ and Mn²⁺ also strongly bind to both holo and apo ferritin. These metals are readily removed by chelating agents. Immobilization of both ferritin forms on acrylamide or sepharose supports readily occurs with complete retention of this metal ion binding ability.

Redox Chemistry. The mineral core of mammalian ferritin undergoes complete reduction to a corresponding Fe²⁺ core (of unknown composition and structure at present). The reduction potential is -310 mv at pH 8 but is strongly pH dependent, suggesting the following overall reaction: $\text{FeOOH} + e + 2\text{H}^+ = \text{FeOH}^+ + \text{H}_2\text{O}$. The mineral core of ferritin is thus capable of storing large numbers of low potential electrons as well as protons in its isolated interior. Partial reduction of the ferritin core produces discrete Fe(II) and Fe(III) within the core mineral.

or	
41	☒
d	☐
	☐
TV Codes	



A-1

Electron Transfer. Addition of labeled $^{57}\text{Fe}^{2+}$ to anaerobic holo ferritin produces $^{57}\text{Fe}^{3+}$ bound to the mineral core, demonstrating electron transfer to the bulk core and indicating that the mineral core is conductive. Addition of Fe^{2+} specific chelators removes $^{57}\text{Fe}^{2+}$, demonstrating the reversibility of this electron transfer reaction. Attempts to determine how redox reagents external to the sequestered mineral core are able to transfer electrons to the interior through the intervening 20-30A protein shell are underway. Three hypotheses are being investigated: 1) Fe^{2+} - Fe^{3+} mediated electron transfer; 2) electron tunneling and 3) protein mediated electron transfer from internal amino acid residues. Evidence for the latter process has been obtained recently.

WORK PLAN (YEAR 2): Having demonstrated the presence of a reduced (Fe^{2+}) core, attempts to reconvert it into the original FeOOH state will be undertaken. This reactivity will demonstrate whether the redox reactions are reversible. We will also establish whether other metal ions (Zn , Cu etc.) are capable of forming aggregated mineral phases within the ferritin interior. Attempts to form magnetite and other iron aggregates will also be undertaken. Finally, we will investigate in more detail the nature of the electron transfer process through the ferritin protein shell.

INVENTIONS: None

PUBLICATIONS:

1. Redox Reactivity of Bacterial and Mammalian Ferritin: Is Reductant Entry into the Ferritin Interior a Necessary Step for Iron Release?

Proc. Natl. Acad. Sci. USA. (1989) 85, 7457-7461.

2. Redox Reactions Associated with Iron Release from Mammalian Ferritin.

Biochemistry (1989) 28,1650-1655.

3. Fe^{2+} Binding to Apo and Holo Mammalian Ferritin

In Press. Biochemistry

TRAINING ACTIVITIES: The following undergraduate research students (all US citizens) were supported during the past year: Steve Hilty (Occidental, College); Chris Van Buren (University of Colorado) and Chi Hua Chiu (University of Colorado).