

AD-A212 713


1A AND 13C NUCLEAR MAGNETIC RESONANCE OF
DIHYDROIMIDAZO-PYRIDINE AND IMID. (U) CHEMICAL RESEARCH
DEVELOPMENT AND ENGINEERING CENTER ABERDEEN.
S MUNAVALLI ET AL. JUL 89 CRDEC-TR-059


1/1

UNCLASSIFIED

F/G 7/3

NL





CHEMICAL
RESEARCH,
DEVELOPMENT &
ENGINEERING
CENTER

CRDEC-TR-059

^1H AND ^{13}C NUCLEAR MAGNETIC RESONANCE
OF DIHYDROIMIDAZO-PYRIDINE AND
IMIDAZO-[1,2-a]-PYRIDINE DERIVATIVES

AD-A212 713

Shekhar Munavalli, Ph.D.
MUNITIONS DIRECTORATE

Fu-Lian Hsu, Ph.D.
August J. Muller, Ph.D.
Everett O. Lewis, Ph.D.
William T. Beaudry, Ph.D.
Linda L. Szafraniec


RESEARCH DIRECTORATE

DTIC
ELECTE
SEP 21 1989
S B D

July 1989

DISTRIBUTION STATEMENT A
Approved for public release
Distribution Unlimited

U.S. ARMY
ARMAMENT
MUNITIONS
CHEMICAL COMMAND



Disclaimer

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorizing documents.

Distribution Statement

Approved for public release; distribution is unlimited.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b RESTRICTIVE MARKINGS	
2a SECURITY CLASSIFICATION AUTHORITY			3 DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.	
2b DECLASSIFICATION / DOWNGRADING SCHEDULE				
4 PERFORMING ORGANIZATION REPORT NUMBER(S) CRDEC-TR-059			5 MONITORING ORGANIZATION REPORT NUMBER(S)	
6a NAME OF PERFORMING ORGANIZATION CRDEC	6b OFFICE SYMBOL (If applicable) SMCCR-MUC	7a NAME OF MONITORING ORGANIZATION		
6c ADDRESS (City, State, and ZIP Code) Aberdeen Proving Ground, MD 21010-5423		7b ADDRESS (City, State, and ZIP Code)		
8a NAME OF FUNDING / SPONSORING ORGANIZATION CRDEC	8b OFFICE SYMBOL (If applicable) SMCCR-MUC	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c ADDRESS (City, State, and ZIP Code) Aberdeen Proving Ground, MD 21010-5423		10 SOURCE OF FUNDING NUMBERS		
		PROGRAM ELEMENT NO	PROJECT NO 1-32-85-000	TASK NO A-372
		WORK UNIT ACCESSION NO		
11 TITLE (Include Security Classification) 14 and ¹³ C Nuclear Magnetic Resonance of Dihydroimidazo-Pyridine and Imidazo-[1,2-a]-Pyridine Derivatives				
12 PERSONAL AUTHOR(S) (see reverse)				
13a TYPE OF REPORT Technical	13b TIME COVERED FROM 86 Sep to 88 Jun	14 DATE OF REPORT (Year, Month, Day) 1989 July	15 PAGE COUNT 26	
16 SUPPLEMENTARY NOTATION				
17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD 20	GROUP 05	SUB-GROUP	1H NMR Dihydroimidazo-pyridines ¹³ C NMR 2D NMR Imidazo-pyridines Chemical shifts Quaternization effects	
19 ABSTRACT (Continue on reverse if necessary and identify by block number) Replacement of the aromatic hydrogens of the pyridinium moiety of the dihydroimidazo-pyridine and imidazo(1,2-a)-pyridine derivatives with electron-donating and electron-withdrawing substituents causes displacement of the chemical shifts of aromatic protons. Quaternization of the nitrogen atoms causes molecular perturbations and affects the ¹ H and ¹³ C nuclear magnetic resonance chemical shifts. The sign of the C and C _β displacements can be used to detect the site of quaternization.				
20 DISTRIBUTION / AVAILABILITY OF ABSTRACT ☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT ☐ DTIC USERS			21 ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a NAME OF RESPONSIBLE INDIVIDUAL SANDRA J. JOHNSON			22b TELEPHONE (Include Area Code) (301) 671-2914	22c OFFICE SYMBOL SMCCR-SPS-T

UNCLASSIFIED

(Original Patterns) (continued)

Alm, John, Ph.D., Hsu, Fu-Lian, Ph.D., Muller, August J., Ph.D.,
Lewicki, Robert G., Ph.D., Beaudry, William T., Ph.D., Szafraniec, Linda L.

PREFACE

The work described in this report was authorized under Project No. 1-32-85-000-A-372. This work was started in September 1986 and completed in June 1988.

The use of trade names or manufacturers' names in this report does not constitute an official endorsement of any commercial products. This report may not be cited for purposes of advertisement.

Reproduction of this document in whole or in part is prohibited except with permission of the Commander, U. S. Army Chemical Research, Development and Engineering Center, ATTN: SMCCR-SPS-T, Aberdeen Proving Ground, Maryland 21010-5423. However, the Defense Technical Information Center and the National Technical Information Service are authorized to reproduce the document for U. S. Government purposes.

This report has been approved for release to the public.

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Blank

CONTENTS

	Page
1. INTRODUCTION	7
2. EXPERIMENTAL PROCEDURES	9
2.1 Compounds	9
2.2 Measurements	9
3. RESULTS AND DISCUSSION	9
4. QUATERNIZATION	12
5. CONCLUSIONS	23
LITERATURE CITED	25

LIST OF FIGURES AND TABLES

	Figure	Page
	2D-NMR	8
Tables		
1	Dihydroimidazo-pyridinium Salts	10
2	Pyridinium Protons	13
3	Dihydroimidazo [1,2-a]-Pyridines and Pyridinium Salts	15
4	Pyridinium Protons	16
5	Pyridinium Carbons	20
6	Pyridinium Carbons of Imidazo-[1,2-a]-Pyridines	21
	Pyridinium Carbons of Imidazo-[1,2-a]-Pyridines Due to Quaternization	22

¹H AND ¹³C NUCLEAR MAGNETIC RESONANCE OF DIHYDROIMIDAZO-PYRIDINE AND IMIDAZO-[1,2-a]-PYRIDINE DERIVATIVES

1. INTRODUCTION

Because of the structural similarity between the purines and the imidazo-[1,2-a]pyridines, the latter have attracted considerable interest. Several attempts have been made to study this 10 π -electron aromatic system and correlate electron densities with their reactivity, ultraviolet (UV) absorption, and other parameters.¹⁻⁵ The ¹H NMR and ¹³C NMR chemical shifts have been correlated with the total π -electron densities of the ring positions.^{1, 3-7} However, difficulties were encountered in calculating electron densities due to the effect of σ -electron polarization.³ Cross-ring effects, para spin-spin coupling, peri-anisotropic effects, and shielding effects associated with atomic dipoles contributed to the chemical shifts of the protons of the azines.⁴ The NMR and UV studies led to the conclusion that the imidazo-pyridine molecule is a 10 π -electron aromatic system with considerable delocalization of electrons and the ability to sustain ring currents.

In the study of the aromatic properties and the interplay of ring currents, the ¹³C NMR has proved to be a versatile tool.⁷⁻⁹ A linear correlation was reported between ¹³C chemical shifts and the net electronic charges.^{4,9} The NMR spectra of the imidazo-pyridinium salts are considerably different from their free bases.¹ Quaternization affects the chemical shifts of carbons adjacent to the quaternized nitrogen atom. The electron charge buildup on the carbon atom moves its resonance upfield, and the shielding pattern is significantly altered by the quaternization.^{7,10} Due to a variety of effects, both N₁ and N₄ are capable of inducing positive and negative net charges.³

Imidazo-[1,2-a]-pyridines are no exception to the direct steric effects observed in aromatic heterocyclics.^{11,12} The peri-interactions between N₁-R and C₈-R on one hand and C₃-R and C₅-R on the other have been reported.^{1,13} The shielding effect for the para position of the substituent has been seen in the ¹H and ¹³C NMR.^{1,15} The substituent induced ortho-effect has been reported to affect ¹H chemical shifts.¹⁷ The presence of a substituent on the pyridine ring of the imidazo-pyridines has no effect on the reactivity of the imidazole moiety and vice versa.¹⁸

In connection with the synthesis of the dihydroimidazo-[1,2-a]-pyridines,¹⁹ ¹H and ¹³C NMR spectra were studied. The NMR spectra of the dihydroimidazo-pyridinium salts are strikingly different from those of the free bases and those of the completely aromatic imidazo-pyridinium derivatives. The pyridine protons appear between 6.7 and 9.0 ppm. Most noticeable differences are seen in the spectra of the pyridinium moiety. The substitution of the pyridine hydrogen with a methyl group causes an upfield displacement of the chemical shift of the ortho-protons. A similar observation was reported in the case of imidazo-[1,2-a]-pyridines.^{17,20} Clearly seen in the ¹³C NMR spectra of the dihydroimidazo-pyridinium derivatives is the displacement of the C₇ signal to higher frequency due to the positive charge on N₄. The chemical shifts of C₅ and C₇ are interchanged in all but one; in that, the C₇ signal is displaced to higher frequency, whereas the C₅ signal has shifted to lower frequency. This was further confirmed by 2D-NMR (see the figure). A similar argument may be advanced in the




Figure. 2D-NMR

case of the 1-benzyl-2-alkyl-2-hydroxy-2,3-dihydroimidazo-[1,2-a]-pyridinium salt.¹⁴ The one exception being 1b. This may be due to the peri-interactions.^{1,10,13}

2. EXPERIMENTAL PROCEDURES

2.1 Compounds.

The synthesis of the dihydroimidazo-[1,2-a]-pyridinium salts was accomplished in a straightforward manner by condensing the respective aminopyridines with 1,2-dibromoethane. The compounds obtained were purified and characterized by elemental and spectrometric analysis.¹⁹ For a review on the synthesis of imidazo-[1,2-a]-pyridines, see Blewitt.³ Attempted O-demethylation of pyrilamine with trimethylsilyl iodide to obtain tripelennamine, a metabolite of the antihistamine drug, gave a 79% yield of the N-benzyl dihydroimidazo-[1,2-a]-pyridinium salt.²¹

2.2 Measurements.


The ¹H NMR chemical shift data were obtained on a Varian E-390 NMR spectrometer for 0.2 M solutions of the appropriate compound (the salts in D₂O and the free bases in CDCl₃) at the probe temperature (34 °C). The chemical shift values were determined relative to the internal standard sodium 3-trimethylsilyl propionate (TSP) or tetramethylsilane (TMS) as appropriate. All signals were downfield from the reference, and the values reported to the nearest 0.05 ppm were obtained by direct measurement using a 10-ppm sweep width. The ¹³C data were obtained on a Varian XL-200 or VXR-400S spectrometer using the same parameters as above. The ¹H and ¹³C values are relative to the respective unsubstituted parent compound for each group of compounds. The sign of the upfield displacement of the respective signal in comparison with the parent compound is considered negative, and the downfield displacement is indicated by a positive sign.

3. RESULTS AND DISCUSSION

The nitrogen atoms of the imidazo-pyridine molecule can exert a variety of effects on electron distribution. In principle, they can induce both positive and negative charges on the carbon atom of the imidazo-pyridines and, therefore, present an interesting situation to study the substituents' effects arising from quaternization of the nitrogens, the cross-ring currents, the effect of the presence and absence of the double bond in the imidazole ring, and the influence of the pyridine moiety on the reactivity of the imidazole ring. This report discusses the effects of the substituents and quaternization on the ¹H and ¹³C NMR chemical shifts of the title compounds. The influence of the double bond of the imidazole ring on the ¹³C chemical shifts of the pyridine moiety is also discussed.

A satisfactory correlation between the calculated and the predicted chemical shifts for all protons in the imidazo-[1,2-a]-pyridines has been reported.²² Because the general characteristics of the NMR spectra of these types of compounds are straightforward, the assignment of the chemical shifts does not pose any special problem. Replacement of the pyridine hydrogen with a methyl group causes an increase in the shielding of the proton on the carbon ortho to the carbon carrying the substituent in the imidazo-[1,2-a]-pyridine series.^{1,17} Table 1 shows that a similar effect prevails in the dihydroimidazo-pyridinium salts. A para effect is also displayed by the substituent. In other words, both ortho and para effects due to substitution are observed. The presence of the electron withdrawing substituents (NO₂ group in 1g; N₆ and N₈ in 1m and 1n) causes deshielding. As a consequence, the ortho protons are

Table 1. Dihydroimidazo-pyridinium Salts




[¹H-NMR Chemical Shift Differences (ΔH) of Dihydroimidazo[1,2-a]pyridinium and -6-azapyridinium and -8-azapyridinium salts.]]

Compound	H5	ΔH5	H6	ΔH6	H7	ΔH7	H8	ΔH8	Ref
1a	8.00	-	6.92	-	7.91	-	7.03	-	20
1b	CH ₃	-	6.74	-0.18	7.77	-0.14	6.84	-0.19	20
1c	7.80	-0.20	CH ₃	-	6.96	-0.81	7.80	+0.77	20
1d	7.82	-0.18	6.79	-0.13	CH ₃	-	6.84	-0.19	20
1e	7.88	-0.12	6.88	-0.11	7.74	-0.17	CH ₃	-	20
1f	8.25	+0.25	Br	-	7.88	-0.03	6.98	-0.05	20
1g	9.25	+1.25	NO ₂	-	8.61	+0.70	7.13	+0.10	20
1h	8.60	+0.60	6.90	±	7.46	±	8.14	+1.11	14
1i	8.48	+0.48	7.05	±	8.00	±	8.22	+1.19	14
1j	8.50	+0.50	6.75	±	7.57	±	7.75-8.23	+ve	14
1k	8.61	+0.61	6.83	±	7.30	±	7.95-8.40	+ve	14
1l	7.97	-0.03	6.80	-0.12	7.95	+0.04	7.14	+0.09	21
1m	8.7	+0.70	N	-	8.21	+0.30	6.92	-0.11	20
1n	8.74	+0.74	7.03	+0.11	8.51	+0.60	N	-	20

[± means: ΔH could not be accurately determined due to the complexity of the signals of these protons.]

Table 1. Dihydroimidazo-pyridinium Salts (continued)



- 1a: $R_1, R_2, R_3, R_4, R_5, a, b, c, d = H$
- b: $R_1 = CH_3; R_2, R_3, R_4, R_5, a, b, c, d = H$
- c: $R_2 = CH_3; R_1, R_3, R_4, R_5, a, b, c, d = H$
- d: $R_3 = CH_3; R_1, R_2, R_4, R_5, a, b, c, d = H$
- e: $R_4 = CH_3; R_1, R_2, R_3, R_5, a, b, c, d = H$
- f: $R_2 = Br; R_1, R_3, R_4, R_5, a, b, c, d = H$
- g: $R_2 = NO_2; R_5 = CH_3; R_1, R_3, R_4, a, b, c, d = H$
- h: $R_5 = CH_3; a = CH_3; b = OH; c = CH_3; R_1, R_2, R_3, R_4, d = H$
- i: $R_5 = CH_3; a = C_6H_5; b = OH; R_1, R_2, R_3, R_4, c, d = H$
- j: $R_5 = CH_2C_6H_5; a = CH_3; b = OH; R_1, R_2, R_3, R_4, c, d = H$
- k: $R_5 = CH_2C_6H_5; a = C_6H_5; b = OH; R_1, R_2, R_3, R_4, c, d = H$
- l: $R_5 = -CH_2C_6H_4-OH(p); R_1, R_2, R_3, R_4, a, b, c, d = H$
- m: dihydroimidazo [1,2-a]-6-azapyridine
- n: dihydroimidazo [1,2-a]-8-azapyridine

shifted downfield. An increased shielding effect due to the electron-donating groups such as -OCH₃, -OC₂H₅, and -N(C₂H₅)₂ on ortho and para protons is also observed in the free bases (Table 2 compounds 3-12). This leads to increased electron density of C₆ and C₈, and this finding is in general agreement with their chemical reactivity vis-a-vis ⁺NO₂.¹⁸ The reversal of the ortho-effect of compounds 13 and 14 may be due to the peri-interaction.

Contrary to the claim that the substituents on the pyridine ring had very little effect on the properties of the imidazole moiety and vice versa,¹⁸ the substituents on the imidazole ring do cause significant changes in the ¹H and ¹³C chemical shifts of the pyridine moiety of the molecule (Tables 1 and 2). The size (bulk) of the alkyl substituent had no significant effect on the chemical shifts of the ortho protons (compounds 3, 7, and 9).

QUATERNIZATION


In principle, quaternization is possible at both sites, N₁ and N₄. The N₄ quaternized compounds such as 2,3-dihydroimidazo-[1,2-a]- and imidazo-[1,2-a]-pyridinium compounds can be synthesized from suitably substituted precursors. Protonation decreases the N-C bond, spin-pairing, and paramagnetic contribution and enhances cross-ring perturbations and anomalous upfield shifts of ¹³C signal of the α-carbons upon protonation of the adjacent nitrogen.²³ Tables 3 and 4 describe the effect of protonation of the bridgehead nitrogen, which produces resonance and inductive effects in the pyridine ring. In the case of dihydro derivatives, the H₅ is displaced upfield in all cases, whereas it exerts mixed effects in 18.

In general, the pyridinium protons of the imidazo-[1,2-a]-pyridinium salts are severely deshielded by the quaternization of the pyridine nitrogen (see Table 4). This effect is more pronounced than in the case of the dihydroanalogs. The presence of the double bond in the imidazole ring also contributes to the chemical shift displacement of H₅, H₆, H₉, and H₈. Quaternization of the free bases generally leads to N₁ quaternized products.^{2,24} As a consequence, the signal due to H₅ is displaced downfield (Table 4, structures 44-46). This paramagnetic effect is directly attributable to the resonance structures (Table 4, structures 48-49).²⁴ Table 5 shows the effect produced by the presence of a positively charged nitrogen (N₄) on the ¹³C chemical shifts of the dihydroderivatives. Except in the case of 1b, the signal due to C₅ is displaced upfield in all compounds examined. The C_{8a} appears to suffer a similar fate.

The quaternization of N₄ produces an upfield displacement of the ¹³C_{8a} signal (e.g., 37, Table 6), whereas the quaternization of N₁ causes a downfield displacement of ¹³C_{8a} (47). The Δ values of C₅, C₆, C₇, and C₈ reflect cross-ring substituent effects, C₅ experiencing the smallest perturbation. Comparison of the ¹³C NMR chemical shifts of the α-carbon carbons of 36 and 37 indicates that the substituent on N₁ does not make a significant contribution to their chemical shifts. The removal of the double bond between C₂ and C₃ of the N₁ quaternized compound (44) causes definite changes in the ¹³C chemical shifts (see 44 and 47); more particularly, the Δc values of C₅, C₇, and C₈ and the signal due to C_{8a} is displaced downfield. Nonetheless, the quaternization does produce significant variation in electronic effects.

Table 7 gives the chemical shift differences (Δc) due to the quaternization of the free bases. Although the quaternization occurs at N₁, the overall contribution of the resonance

Table 2. Pyridinium Protons



[(¹H-NMR Chemical Shift Differences (ΔH) of Pyridinium Protons of the Imidazo-[1,2-a]-pyridines.)]

Compound	H ₅	ΔH_5	H ₆	ΔH_6	H ₇	ΔH_7	H ₈	ΔH_8	Ref
2	8.16	-	6.78	-	7.15	-	7.70	-	1,26
3	CH ₃	-	6.78	0.0	7.23	+0.08	7.60	-0.10	1,25
4	7.90	-0.26	CH ₃	-	6.99	-0.16	7.60	-0.10	1,2,17
5	7.96	-0.20	6.45	-0.33	CH ₃	-	7.36	-0.34	1,2,20
6	8.04	-0.12	6.68	-0.10	6.98	-0.12	CH ₃	-	1,2
7	C ₂ H ₅	-	6.62	-0.16	7.19	+0.04	7.61	-0.09	13
8	OC ₂ H ₅	-	5.91	-0.87	7.10	-0.05	7.85	+0.15	18
9	CH(CH ₃) ₂	-	6.62	-0.16	7.19	+0.04	7.61	-0.09	13
10	N(C ₂ H ₅) ₂	-	6.35	-0.43	7.18	+0.03	7.44	-0.16	25
11	7.83	-0.33	6.38	-0.10	N(C ₂ H ₅) ₂	-	6.51	-1.19	25
12	7.70	-0.46	6.66	-0.12	6.28	-0.67	N(C ₂ H ₅) ₂	-	25
13	NHAc	-	7.09	+0.31	7.28	+0.13	7.47	-0.23	25
14	9.22	+1.06	NHAc	-	7.15	0.0	7.55	-0.15	25
15	8.24	+0.08	6.84	+0.06	7.97	+0.82	NHAc	-	25
16	Br	-	6.89	+0.11	6.86	-0.29	7.48	-0.22	26
17	8.32	+0.16	Br	-	7.20	+0.05	7.59	-0.11	25
18	8.19	+0.03	6.95	+0.17	7.25	+0.10	7.67	-0.03	25
19	7.89	-0.27	6.79	+0.01	7.12	-0.03	7.64	-0.06	25,26
20	CH ₃	-	6.48	-0.30	7.01	-0.14	7.43	-0.27	26

Table 2. Pyridinium Protons (continued)

Com- pound	H ₅	ΔH_5	H ₆	ΔH_6	H ₇	ΔH_7	H ₈	ΔH_8	Ref.
21	OC ₂ H ₅	-	-	-	8.30	+1.15	6.13	-1.57	18
22	CH ₃	-	7.35	+0.57	7.85	+0.70	7.85	+0.15	25
23	Br	-	6.89	+0.09	6.86	-0.29	7.48	-0.22	26
24	NH ₂	-	5.8	-0.98	6.93	-	6.93	-	26
25	9.0	+0.84	7.40	+0.62	8.40	+1.25	NHAc	-	25
26	Br	-	6.89	+0.11	6.86	-0.29	7.48	-0.22	26
27	7.90	-0.26	CH ₃	-	7.00	-0.15	7.51	-0.19	18
28	8.05	-0.09	6.70	-0.08	6.98	-0.17	CH ₃	-	18
29	8.21	+0.05	Cl	-	7.15	0.0	7.60	-0.10	18
30	NO ₂	-	-	-	6.48	-0.67	8.60	+0.90	18
31	9.45	+1.27	7.53	+0.75	7.83	+0.68	8.05	+0.35	25

Table 3. Dihydroimidazo-[1,2-a]-Pyridines and Pyridinium Salts

[(¹H-Chemical Shift Differences (ΔH) of Dihydroimidazo[1,2-a]-pyridines and-Pyridinium salts.)]

Compounds* ++	ΔH_5	ΔH_6	ΔH_7	ΔH_8
1a and 2	-0.16	+0.14	+0.76	-0.66
1b and 3	-	-0.04	+0.54	+0.76
1c and 4	-0.10	-	-0.03	+0.20
1d and 5	-0.10	+0.33	-	-0.31
1e and 6	-0.16	+0.20	+0.24	-


* Values are taken from Tables 1 and 2.

++ Downfield displacement +ve, upfield displacement -ve with respect to the free base.

Table 4. Pyridinium Protons


(¹ H-NMR Chemical Shift Differences (ΔH) of Pyridinium Protons of the Imidazo[1,2-a]pyridinium salts.)										
Com- pound	+ N	H ₅	ΔH ₅	H ₆	ΔH ₆	H ₇	ΔH ₇	H ₈	ΔH ₈	Ref
2	-	8.20	-	6.80	-	7.21	-	7.56	-	26
38	+ N ₄	8.62	+0.42	CH ₃	-	7.87	+0.66	7.87	+0.31	2
39	+ N ₄	8.58	+0.38	7.33	+0.53	CH ₃	-	7.68	+0.12	2
40	+ N ₄	8.73	+0.53	7.53	+0.73	7.40	+0.69	CH ₃	-	2
41	+ N ₄	8.56	+0.36	CH ₃	-	7.93	+0.72	7.93	+0.37	2
42	+ N ₄	8.62	+0.42	7.40	+0.60	CH ₃	-	7.84	+0.28	2
43	+ N ₄	8.67	+0.47	7.46	+0.66	7.83	+0.62	CH ₃	-	2
44	+ N ₁	9.70	+1.5	-	-	-	-	-	-	14
45	+ N ₁	9.14	+0.94	-	-	-	-	-	-	14
46	+ N ₁	8.89	+0.79	-	-	-	-	-	-	14

Table 4. Pyridinium Protons (continued)




- 2 $R_1, R_2, R_3, R_4, R_5, R_6 - H$
- 3 $R_1-CH_3; R_2, R_3, R_4, R_5, R_6 - H$
- 4 $R_2-CH_3; R_1, R_3, R_4, R_5, R_6 - H$
- 5 $R_3-CH_3; R_1, R_2, R_4, R_5, R_6 - H$
- 6 $R_4-CH_3; R_1, R_2, R_3, R_5, R_6 - H$
- 7 $R_1-C_2H_5; R_2, R_3, R_4, R_5, R_6 - H$
- 8 $R_1-OC_2H_5; R_2, R_3, R_4, R_5, R_6 - H$
- 9 $R_1-CH(CH_3)_2; R_2, R_3, R_4, R_5, R_6 - H$
- 10 $R_1-N(C_2H_5)_2; R_2, R_3, R_4, R_5, R_6 - H$
- 11 $R_3-N(C_2H_5)_2; R_1, R_2, R_4, R_5, R_6 - H$
- 12 $R_4-N(C_2H_5)_2; R_1, R_2, R_3, R_5, R_6 - H$
- 13 $R_1-NHAc; R_2, R_3, R_4, R_5, R_6 - H$
- 14 $R_2-NHAc; R_1, R_3, R_4, R_5, R_6 - H$
- 15 $R_4-NHAc; R_1, R_2, R_3, R_5, R_6 - H$
- 16 $R_1-Br; R_2, R_3, R_4, R_5, R_6 - H$
- 17 $R_2-Br; R_1, R_3, R_4, R_5, R_6 - H$
- 18 $R_6-Br; R_1, R_2, R_3, R_4, R_5 - H$
- 19 $R_6-CH_3; R_1, R_2, R_3, R_4, R_5 - H$
- 20 $R_1-CH_3; R_6-Br, R_2, R_3, R_4, R_5 - H$
- 21 $R_1-OCH_3; R_6-NO_2; R_2, R_3, R_4, R_5 - H$
- 22 $R_1-CH_3; R_6-NO_2; R_2, R_3, R_4, R_5 - H$
- 23 $R_1-Br; R_6-CH_3; R_2, R_3, R_4, R_5 - H$
- 24 $R_1-NH_2; R_6-CH_3; R_2, R_3, R_4, R_5 - H$
- 25 $R_4-NHAc; R_6-NO_2; R_2, R_3, R_4, R_5, - H$
- 26 $R_1-Br; R_5-CH_3; R_2, R_3, R_4, R_6 - H$

Table 4. Pyridinium Protons (continued)



(36-43)




(44-46)


- 27 $R_7 = \text{CH}_3$; $R_5 = \text{CO}_2\text{C}_2\text{H}_5$; $R_1, R_3, R_4, R_6 = \text{H}$
 28 $R_4 = \text{CH}_3$; $R_5 = \text{CO}_2\text{C}_2\text{H}_5$; $R_1, R_2, R_3, R_6 = \text{H}$
 29 $R_2 = \text{Cl}$; $R_5 = \text{CO}_2\text{C}_2\text{H}_5$; $R_1, R_3, R_4, R_6 = \text{H}$
 30 $R_1 = \text{NO}_2$; $R_5 = \text{CO}_2\text{C}_2\text{H}_5$; $R_2, R_3, R_4, R_6 = \text{H}$
 31 $R_6 = \text{NO}_2$; $R_1, R_2, R_3, R_4, R_5 = \text{H}$
 32 $R_5 = \text{CO}_2\text{C}_2\text{H}_5$; $R_1, R_2, R_3, R_4, R_6 = \text{H}$
 33 $R_2 = \text{NO}_2$; $R_5 = \text{CO}_2\text{C}_2\text{H}_5$; $R_1, R_3, R_4, R_6 = \text{H}$
 34 $R_4 = \text{NO}_2$; $R_5 = \text{CO}_2\text{C}_2\text{H}_5$; $R_1, R_2, R_3, R_6 = \text{H}$
 35 $R_2 = \text{Cl}$; $R_1, R_3, R_4, R_5, R_6 = \text{H}$
 36: $R_1, R_2, R_3, R_4, R_5, R_6, R_7 = \text{H}$
 37: $R_5 = \text{CH}_3$; $R_1, R_2, R_3, R_4, R_6, R_7 = \text{H}$
 38: $R_2 = \text{CH}_3$; $R_1, R_3, R_4, R_5, R_6, R_7 = \text{H}$
 39: $R_3 = \text{CH}_3$; $R_1, R_2, R_4, R_5, R_6, R_7 = \text{H}$
 40: $R_4 = \text{CH}_3$; $R_1, R_2, R_3, R_5, R_6, R_7 = \text{H}$
 41: $R_2, R_5 = \text{CH}_3$; $R_1, R_3, R_4, R_6, R_7 = \text{H}$
 42: $R_3, R_5 = \text{CH}_3$; $R_1, R_2, R_4, R_6, R_7 = \text{H}$
 43: $R_4, R_5 = \text{CH}_3$; $R_1, R_2, R_3, R_6, R_7 = \text{H}$

- 44: $R_5 = \text{CH}_2\text{C}_6\text{H}_5$; $R_6 = \text{CH}_3$; $R_1, R_2, R_3, R_4, R_7 = \text{H}$
 45: $R_5 = \text{CH}_2\text{C}_6\text{H}_5$; $R_6 = \text{C}_6\text{H}_5$; $R_1, R_2, R_3, R_4, R_7 = \text{H}$
 46: $R_5 = \text{CH}_2\text{C}_6\text{H}_5$; $R_6, R_7 = \text{CH}_3$; $R_1, R_2, R_3, R_4 = \text{H}$


Table 4. Pyridinium Protons (continued)



4.7



4.8



4.9

Table 5. Pyridinium Carbons

[¹³C-NMR: Chemical Shift Differences (Δ c) of the Pyridinium Carbons
of the Dihydroimidazo-[1,2-a]pyridinium Derivatives.]

Com- pound	C ₅	Δ C ₅	C ₆	Δ C ₆	C ₇	Δ C ₇	C ₈	Δ C ₈	C _{8a}	Δ C _{8a}	Ref
1a	139.20	-	116.50	-	147.30	-	112.20	-	158.90	-	20
1b	150.80	+11.60	116.30	-0.20	147.30	0.0	108.80	-3.40	159.20	+0.30	20
1c	135.70	-3.50	127.30	+10.80	137.00	-10.30	118.80	+6.60	149.30	-9.60	20
1d	138.70	-1.20	118.60	+2.10	161.00	+13.70	111.10	-1.10	158.60	-0.30	20
1e	136.20	-3.00	116.60	+0.10	145.60	-1.70	123.00	+10.80	158.90	0.0	20
1f	137.50	-1.70	114.90	-1.60	145.60	-1.70	108.20	-4.00	153.40	-5.50	20
11	144.60	+5.40	112.70	-3.80	138.00	-9.30	107.70	-4.50	157.20	-1.70	21

Table 6. Pyridinium Carbons of Imidazo-[1,2-a]-Pyridines

[¹³ C-NMR: Effect of the Substituents on Chemical Shift Difference (ΔC) of Pyridinium Carbons of Imidazo[1,2-a]-Pyridines.]											
Com- pound	C ₅	ΔC ₅	C ₆	ΔC ₆	C ₇	ΔC ₇	C ₈	ΔC ₈	C _{8a}	ΔC _{8a}	Ref
2*	125.80	-	112.20	-	124.30	-	117.70	-	145.40	-	10
3*	134.40	+8.60	112.20	0.0	124.30	0.0	115.00	-2.7	145.80	+0.4	13
4*	123.50	-2.30	121.90	+9.7	127.60	+3.3	117.00	-0.7	144.60	-0.8	13
5*	125.10	-0.7	115.00	+2.8	135.30	+11.0	116.40	-1.6	146.00	+0.6	13
6*	123.70	-2.1	112.00	+0.8	123.30	-1.2	127.50	+9.9	146.00	+0.6	13
7*	139.90	+14.1	109.30	-2.9	124.6	+0.3	115.20	-2.5	146.10	+0.7	13
8*	148.30	+22.50	88.50	-23.70	125.90	+1.6	108.70	-9.0	145.90	+0.50	18
9*	144.30	+18.5	107.20	-5.0	124.60	+0.3	115.30	-2.4	146.10	+0.70	13
36*	129.29	+3.49	117.12	+4.92	133.59	+9.29	112.06	-5.64	139.18	-6.22	10
37*	129.32	+3.52	117.08	+4.88	133.27	+8.97	118.42	+0.72	139.16	-6.25	10
47*	137.50	+11.70	114.90	+2.70	145.60	+21.30	108.20	-9.5	153.43	+8.03	14
32**	127.60	-	117.80	-	126.50	-	118.00	-	144.60	-	18
28**	125.20	-2.40	118.50	+0.70	124.70	-1.8	127.40	-9.4	145.20	+0.6	18
29**	127.50	-0.10	120.50	+2.7	125.40	-1.10	118.70	+0.7	142.90	-1.70	18
33**	120.20	-7.4	138.50	+20.7	128.80	+2.3	120.50	+2.5	144.50	-0.10	18
34**	139.90	+6.30	119.80	+2.00	125.70	-0.80	137.20	+19.2	137.50	-7.1	18
35**	125.20	-0.60	118.90	+6.7	124.90	+0.60	117.60	-0.40	143.00	-2.40	18

*ΔC values with respect to 2.

**ΔC values with respect to 32.

Table 7. Pyridinium Carbons of Imidazo-[1,2-a]-Pyridines Due to Quaternization

[¹³ C-NMR: Chemical Shift Difference (ΔC) of Pyridinium Carbons of Imidazo-[1,2-a]Pyridines Due to Quaternization of N ₁ .]*						
Compound	ΔC ₅	ΔC ₆	ΔC ₇	ΔC ₈	ΔC _{8a}	Ref
4	+4.2	+0.8	+4.2	-0.7	-0.8	18
6	-0.1	+0.5	+2.4	-0.7	-0.9	18
8	+0.4	+2.7	+2.7	-2.2	-2.2	18
28	+0.8	+0.5	+1.4	+0.7	-1.5	18
29	+5.4	+2.0	+1.4	-2.6	-1.9	18
32	+1.9	-2.8	+1.9	-1.9	-1.3	18
35	+2.1	+1.2	+0.6	-1.0	-1.2	18

* With respect to 2.

structure. By comparing the chemical shifts of the pyridine protons, we can detect and decide which of the two nitrogen atoms is quaternized.

Comparison of the ^{13}C chemical shifts of 2, 3, and 34 indicates that the introduction of a strong electron withdrawing group causes a significant downfield displacement of the ^{13}C NMR signal of the carbon carrying the substituent.

5. CONCLUSIONS

The origin, magnitude, and mode of transmission of the substituent induced electronic effects of the pyridines and their salts have been the subject of intensive investigations. These effects seem to persist even in the case of the bicyclic systems such as the imidazo pyridines. Based on the study of ^1H and ^{13}C NMR spectra of dihydromidazo and imidazo pyridines, we conclude that

- The presence of a large electron donating group (e.g., para $-\text{NMe}_2$, $-\text{OCH}_3$, $-\text{NH}_2$, $-\text{OH}$, $-\text{OR}$, $-\text{NR}_2$, $-\text{N}(\text{R})_2$) causes an upfield shift of the pyridine protons.
- In general, the electron donating groups cause an upfield displacement of the pyridine protons, whereas the electron withdrawing groups induce a downfield displacement. (This is reminiscent of the substituent effects observed in the case of the methylenedioxy pyridinium derivatives^{2,3,7} and substituted pyridinium compounds^{2,8,14}).
- The size (bulk) of the alkyl substituent does not affect the magnitude of the chemical shifts of the pyridine protons.
- The sign of the displacement of the C_5 and the C_{8a} can be used to detect the site of quaternization (see Table 7).
- The ^{13}C signal of the C_7 of dihydromidazo and imidazo pyridinium salts is shifted to higher frequency, and the C_5 signal is shifted to lower frequency, due to quaternization of the bridgehead nitrogen.
- The presence of a positive charge on the nitrogen causes an upfield shift of the amino proton; namely, the H_5 (see Table 3).
- As confirmed by ^{13}C NMR measurements in all but one compound, the chemical shift of the C_7 of the dihydromidazo and imidazo pyridinium salts is shifted to higher frequency, and the C_5 is shifted to lower frequency, due to quaternization of the bridgehead nitrogen.

Blank

LITERATURE CITED

1. Paudler, W.W., and Blewitt, H.L., "NMR Spectra and π -Electron Densities of Some Imidazo(1,2-a) Pyridines," Tetrahedron Vol. 31, pp 353-361 (1965).
2. Paudler, W.W., and Blewitt, H.L., "Ten π Electron Nitrogen Heterocyclic Compounds. V. The Site of Protonation and N-Methylation of Imidazo[1,2-a]pyridines and the Planarity of the Ring System," J. Org. Chem. Vol. 31, p 1295 (1966).
3. Blewitt, H.L., Special Topics in Heterocyclic Chemistry, Vol. 50, pp 117-178, A. Weissberger and E.C. Taylor, Eds., Wiley and Sons, New York, NY, 1977.
4. Black, P.J., Brown, R.D., and Heffernan, M.L., "Proton Chemical Shifts and Electron Densities in Aromatic and Heteroaromatic Molecules. I. Procedure and Chemical Shift Corrections; Applications to Azines," Aust. J. Chem. Vol. 20, pp 1305-1323 (1967).
5. Galasso, V., De Altì, G., and Bigotto, A., "Electronic Structure and Absorption Spectra of Indolizines," Theor. Chim. Acta Vol. 9, pp 222-229 (1968).
6. Kleinpeter, E., Borsdorf, R., Fischer, G., and Hofmann, H.J., "NMR Studies of Azaindolizines," J. Prakt. Chem. Vol. 314, pp 5-524 (1972).
7. Grant, D.M., Pugmire, R.J., Robins, M.J., and Robins, R.K., "Carbon-13 Magnetic Resonance. XX. 4-Azaindene (Pyrrocoline) and Related Bridgehead Nitrogen Heterocycles," J. Am. Chem. Soc. Vol. 93, pp 1887-1893 (1971).
8. Grant, D.M., and Pugmire, R.J., "Carbon-13 Magnetic Resonance. XIX. Benzimidazole, Purine, and Their Anionic and Cationic Species," J. Am. Chem. Soc. Vol. 93, pp 1880-1887 (1971).
9. Witanowski, M., Stefaniak, L., Sicinska, W., and Webb, G.A., "Carbon 13 Chemical Shifts of Some Azaindolizines Versus Electron Charge Distribution," J. Mol. Struct. Vol. 64, pp 15-27 (1980).
10. Pugmire, R. J., Smith, J. C., Grant, D. M., Stanovick, B., and Tisler, M., "Carbon-13 NMR Investigation of the Protonation and Quaternization of Azoloazines with a Bridgehead Nitrogen," J. Heterocycl. Chem. Vol. 13, p 1057 (1976).
11. Zoltewicz, Z.A., and Deady, L.W., Adv. Heterocycl. Chem. Vol. 22, p 11 (1971).
12. Gallo, R., Chanon, M., Lund, H., and Metzger, J., "Quaternization of Hindered Amines. Correlation Between Geometry and Steric Parameters," Tet. Letters, p 3857 (1972).
13. Maury, G., and Pigiere, C., "Transmission of a Peri Steric Effect Through a Heterocyclic Nucleus. I. Synthesis and Spectral Properties of Peri Disubstituted Dialkylimidazo(1,2-a) Pyridines," Tetrahedron. Vol. 37, pp 83-90 (1981).
14. Kuhia, D.E., and Watson, H.A., "Synthesis of Imidazo(1,2-a)Pyridinium Salts from 2-Alkylaminopyridines," J. Heterocycl. Chem. Vol. 15, pp 1149-1152 (1978).
15. Parker, R.G., and Roberts, J.D., "Nuclear Resonance Spectroscopy. ^{13}C Spectra of Indole and Methylindoles," J. Org. Chem. Vol. 35, pp 996-999 (1970).

16. Platzner, N., Basselier, J.J., and Demerseman, P., "Carbon-13 NMR Study of Benzofurans. Effect of Methyl Substituents. Additivity and Steric Effects," Bull. Soc. Chim. (France), pp 905-912 (1974).
17. Otsuru, M., Tori, K., and Watanabe, H., "Correlation of the Methyl Substitution Effect on the Chemical Shifts of Ortho-Protons with ρ -Bond Orders in Aromatic Molecules," Chem. Pharm. Bull. (Japan) Vol.15, pp 1015-1020 (1967).
18. Teulade, J.-C., Escalé, R., Grassy, G., Girard, J.-P., and Chapat, J.-P., "Reactivite de derives de l'imidazo(1,2-a) Pyridine vis-a-vis de la reaction de nitration Effets de substituant par RMN ^{13}C et CNDO," Bull. Soc. Chim. (France), p 529 (1979).
19. Munavalli, S., Hsu, F.-L., and Poziomek, E.J., "Facile Synthesis of 2,3-Dihydro-imidazopyridines and -Pyrimidines," Chem. Ind. (London), pp 243-244 (1987).
20. Hand, E.S., and Paudler, W.W., "Selenium Derivatives of Imidazo(1,2-a) Pyridines," J. Org. Chem. Vol. 40, pp 2916-2920 (1975).
21. Hsu, F.L., Munavalli, S., and Yeh, Y., "O-Demethylation of Pyrilamine," J. Pharm. Sci. Vol. 77, p 727 (1988).
22. Paudler, W.W., and Kuder, J.E., "Ten- π -Electron Nitrogen Heterocyclic Compounds. VIII. Semi-Empirical Calculations of Chemical-Shifts of Ring-Protons and of Methyl-Protons," J. Heterocyclic Chem. Vol. 3, p 33 (1966).
23. Pugmire, R.J., and Grant, D.M., "Carbon-13 Magnetic Resonance. XII. Five-Membered Nitrogen Heterocycles and Their Charged Species," J. Am. Chem. Soc. Vol. 90, pp 4232-4238 (1968).
24. Amarego, W.L.F., "Ionization and Ultraviolet Spectra of Indolizines," J. Chem. Soc., p 4226 (1964).
25. Hand, E.S., and Paudler, W.W., "Telemination of the Imidazo(1,2-a) Pyridine System," J. Am. Org. Chem. Vol. 43, p 2900 (1978).
26. Hand, E.S., and Paudler, W.W., "Reaction of 3-Substituted Imidazo(1,2-a) Pyridines with Bromine(1+) and the Alleged 5-Bromo-Substituted Product," J. Org. Chem. Vol. 45, pp 3738-3745 (1980).
27. Munavalli, S., Czafraniec, L. L., Beaudry, W., and Poziomek, E. J., "NMR Studies of Methylenebispyridinium Derivatives," Magn. Reson. Chem. Vol. 24, p 743 (1986).
28. Wehrli, F. W., Giger, W., and Simon, W., " ^{13}C -, ^{14}N -und ^1H -kernresonanz-spektroskopische Untersuchungen an *m-p*-substituierten N-Methylpyridiniumjodiden," Helv. Chim. Acta, Vol. 54, p 229 (1971).
29. Deady, L. W., and Willett, G. D., "N-Methyl Chemical Shifts in Some 2-, 3-, and 4-Substituted N-Methylpyridinium Salts," Org. Magn. Reson. Vol. 6, p 53 (1974).

END

FILMED

10-89

DTIC