

AD-A215 265

DTIC FILE COPY

REPORT DOCUMENTATION PAGE			Form Approved 0148 No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE	3. REPORT TYPE AND DATES COVERED	
			Final Report, 0 Aug 86 to 31 Jul 89	
4. TITLE AND SUBTITLE			5. FUNDING NUMBERS	
NONLINEAR WAVE PHENOMENA IN HYDRODYNAMICS			AFOSR-86-0192	
AUTHOR(S)			PE - 61102F	
Professor David J. Benney			PR - 2304 TA - A4	
6. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)			8. PERFORMING ORGANIZATION REPORT NUMBER	
Massachusetts Institute of Technology Department of Mathematics Cambridge, MA 02139			AFOSR-TR-89-1586	
7. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
AIR FORCE OFFICE OF SCIENTIFIC RESEARCH Mathematical and Information Sciences Building 410 Dolling AFB, DC 20332-6448				
11. SUPPLEMENTARY NOTES				
DTIC ELECTE DEC 08 1989 S B D				
12a. DISTRIBUTION/AVAILABILITY STATEMENT			12b. DISTRIBUTION CODE	
Distribution unlimited; approved for public release.				
13. ABSTRACT (Maximum 200 words)				
The principal accomplishment of this study is the development of quite different theory for nonlinear instabilities in shear flows. This theory is an asymptotic one and not ad hoc. Despite many attempts in the past the existing theories appear to be too weak mathematically and fail to capture the essentials of the nonlinear dynamical process in the breakdown of laminar flow. In particular the crucial elements of strong mean flow distortion and three dimensionality have not played an important role in past studies. The new theory, called the mean flow-first harmonic theory, remedies these two defects and shows why they are expected to be critical factors in a nonlinear theory. It is shown that there exists a truly nonlinear interaction between the evolving mean flow and one or more first harmonic waves. Both viscous and inviscid flow problems have been considered. In particular detailed calculations have been made for free surface flows, channel flow, and the Blasius boundary layer. Wake and mixing layer problems are currently under study. The second development concerns the critical layer singularity that arises in inviscid shear flows. In the past this difficulty for perfect fluids has been circumvented by using viscosity or nonlinearity. Each of these two limits lead to different results concerning the phase change and therefore the eigenvalues. A third and possibly more relevant limit is to consider a wave packet analysis in the neighborhood of the critical level. Such an analysis suggests that most amplified and damped disturbances are permitted in this limit. Further work is in progress.				
14. SUBJECT TERMS			15. NUMBER OF PAGES	
			1	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT	
Unclassified	Unclassified	Unclassified	SAR	

Final Report for AFOSR-86-0192B

Nonlinear Wave Phenomena In Hydrodynamics

The principal accomplishment of this study is the development of quite different theory for nonlinear instabilities in shear flows. This theory is an asymptotic one and not 'ad hoc'. Despite many attempts in the past the existing theories appear to be too weak mathematically and fail to capture the essentials of the nonlinear dynamical process in the breakdown of laminar flow. In particular the crucial elements of strong mean flow distortion and three dimensionality have not played an important role in past studies. The new theory, called the mean flow-first harmonic theory, remedies these two defects and shows why they are expected to be critical factors in a nonlinear theory. It is shown that there exists a truly nonlinear interaction between the evolving mean flow and one or more first harmonic waves. Both viscous and inviscid flow problems have been considered. In particular detailed calculations have been made for free surface flows, channel flow, and the Blasius boundary layer. Wake and mixing layer problems are currently under study.

The second development concerns the critical layer singularity that arises in inviscid shear flows. In the past this difficulty for perfect fluids has been circumvented by using viscosity or nonlinearity. Each of these two limits lead to different results concerning the phase change and therefore the eigenvalues. A third and possibly more relevant limit is to consider a wave packet analysis in the neighborhood of the critical level. Such an analysis suggests that most amplified and damped disturbances are permitted in this limit. Further work is in progress.



Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

89 12 05 085