

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE December 1989		3. REPORT TYPE AND DATES COVERED presentation/paper	
4. TITLE AND SUBTITLE ON THE VARIABILITY OF SPORADIC-E				5. FUNDING NUMBERS PE: 0601152N WU: DN306 212 PN: ZT43	
6. AUTHOR(S) A. K. Paul					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Ocean Systems Center San Diego, CA 92152-5000				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Chief of Naval Research Independent Research Programs (IR) OCNR-10P Arlington, VA 22217-5000				10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES					
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The analysis of digital ionograms taken in short time intervals show rapid changes of sporadic-E parameters. The lifetime (observation time) of an individual layer can be shorter than 1 minute. Frequently two or more Es-layers are observed in a given ionogram. Angle of arrival measurements show that the echoes are returned from two or more distinctly different areas in the sky implying a relatively small scale structure in horizontal directions. This conclusion is supported by the fact that almost 25% of the Es-layers found in our data set of more than 4,000 ionograms showed tilt angles of 20 degrees or more. The tilt angles were often highly variable. In many cases where continuity of the top frequency fEs was observed over relatively long sequences of ionograms, the variation of the tilt angles still indicated some instability of the layer. 					
Published in the <i>Proceedings URSI</i> , National Radio Science Meeting, January 1989.					
14. SUBJECT TERMS E-region variability E-region tilts sporadic E-layer tilts				15. NUMBER OF PAGES	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UNLIMITED		

AD-A216 660

A. Paul

**NATIONAL ACADEMIES OF SCIENCE AND ENGINEERING
NATIONAL RESEARCH COUNCIL**

**of the
UNITED STATES OF AMERICA**

**UNITED STATES NATIONAL COMMITTEE
International Union of Radio Science**



**National Radio Science Meeting
4-6 January 1989**

**Sponsored by USNC/URSI
in cooperation with
Institute of Electrical and Electronics Engineers**

**University of Colorado
Boulder, Colorado
U.S.A.**

G-6 Fr-PM

ON THE VARIABILITY OF SPORADIC-E

Adolf K. Paul
Naval Ocean Systems Center, Code 542
Ocean and Atmospheric Sciences Division
San Diego, California 92152-5000

G6-6
1520

The analysis of digital ionograms taken in short time intervals show rapid changes of sporadic-E parameters. The lifetime (observation time) of an individual layer can be shorter than 1 minute. Frequently two or more Es-layers are observed in a given ionogram. Angle of arrival measurements show that the echoes are returned from two or more distinctly different areas in the sky implying a relatively small scale structure in horizontal directions. This conclusion is supported by the fact that almost 25% of the Es-layers found in our data set of more than 4,000 ionograms showed tilt angles of 20 degrees or more. The tilt angles were often highly variable. In many cases where continuity of the top frequency fEs was observed over relatively long sequences of ionograms, the variation of the tilt angles still indicated some instability of the layer.

X	
By <u>NOSC</u>	
Distribution/	
Availability Codes	
Dist	A-1
21	

