

2

DTIC FILE COPY

Office of Naval Research
Contract N00014-87-K-0438

R&T Code 413j009

Technical Report No. 6

AD-A219 505

New Synthesis Route for 1,1,2,2-Tetracyanocyclopropanes

by

Ju-Yeon Lee and H.K. Hall, Jr.

J. Org. Chem.

submitted

University of Arizona
Department of Chemistry
Tucson, AZ 85721

March, 1990

DTIC
ELECTE
MAR 22 1990
S E D

Reproduction in whole, or in part, is permitted for any purpose of the United States Government.

*This document has been approved for public release and sale: its distribution is unlimited.

00 03 22 017

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION None			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY None			3. DISTRIBUTION / AVAILABILITY OF REPORT Unlimited		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE None					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report #6			5. MONITORING ORGANIZATION REPORT NUMBER(S) ONR N00014-87-K-0437		
6a. NAME OF PERFORMING ORGANIZATION University of Arizona		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Office of Naval Research		
6c. ADDRESS (City, State, and ZIP Code) Department of Chemistry University of Arizona Tucson, AZ 85721			7b. ADDRESS (City, State, and ZIP Code) 800 North Quincy Avenue Arlington, VA 22217		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c. ADDRESS (City, State, and ZIP Code) 800 N. Quincy Avenue Arlington, VA 22217			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
11. TITLE (Include Security Classification) New Synthesis Route for 1,1,2,2-Tetracyanocyclopropanes					
12. PERSONAL AUTHOR(S) Ju-Yeon Lee and H.K. Hall, Jr.					
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day)	
15. PAGE COUNT					
16. SUPPLEMENTARY NOTATION J. Org. Chem.					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Det.		
FIELD	GROUP	SUB-GROUP			
19. ABSTRACT (Continue on reverse if necessary and identify by block number) 1,1-Dimethoxy-2-bromoethylene 1 reacts with tetracyanoethylene (TCNE, 3) at -10°C to form methyl 2,2,3,3-tetracyanocyclopropanecarboxylate 6 in high yield. A similar reaction occurs with 1,1-diethoxy-2-bromoethylene 2 to yield ethyl 2,2,3,3-tetracyanocyclopropanecarboxylate 7. Further, <i>B</i> -bromoethyl 2,2,3,3-tetracyano-1-methylcyclopropanecarboxylate 10 and <i>B</i> -bromoethyl 2,2,3,3-tetracyano-1-ethylcyclopropanecarboxylate 11 were obtained by reacting TCNE with 2-bromomethylidene 8 and 2-bromoethylidene-1,3-dioxolane 9, respectively. A reaction mechanism via a zwitterion and intra-molecular displacement of bromide is proposed.					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT ☒ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT. ☐ DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION		
22a. NAME OF RESPONSIBLE INDIVIDUAL H.K. Hall, Jr.			22b. TELEPHONE (Include Area Code) 602-621-5325		22c. OFFICE SYMBOL

New Synthesis Route for 1,1,2,2-Tetracyanocyclopropanes

by

Ju-Yeon Lee and H. K. Hall, Jr.

C. S. Marvel Laboratories
Department of Chemistry
University of Arizona
Tucson, AZ 85721

Mr. Tolson _____
Mr. Boardman _____
Mr. Nichols _____
Mr. Belmont _____
Mr. Ladd _____
Mr. Clegg _____
Mr. Glavin _____
Mr. Harbo _____
Mr. Rosen _____
Mr. Tracy _____
Mr. Egan _____
Mr. Gurnea _____
Mr. Hendon _____
Mr. Pennington _____
Mr. Quinn Tamm _____
Tele. Rm. _____
Mr. Nease _____
Miss Gandy _____

Special Agent in Charge

A-1

A-1



Abstract

1,1-Dimethoxy-2-bromoethylene 1 reacts with tetracyanoethylene (TCNE, 3) at -10°C to form methyl 2,2,3,3-tetracyanocyclopropanecarboxylate 6 in high yield. A similar reaction occurs with 1,1-diethoxy-2-bromoethylene 2 to yield ethyl 2,2,3,3-tetracyanocyclopropanecarboxylate 7. Further, β -bromoethyl 2,2,3,3-tetracyano-1-methylcyclopropanecarboxylate 10 and β -bromoethyl 2,2,3,3-tetracyano-1-ethylcyclopropanecarboxylate 11 were obtained by reacting TCNE with 2-bromomethylidene 8 and 2-bromoethylidene-1,3-dioxolane 9, respectively. A reaction mechanism via a zwitterion and intra-molecular displacement of bromide is proposed.

Introduction

1,1,2,2-Tetracyanocyclopropane can be prepared rather easily by reaction of aqueous formaldehyde and malononitrile¹ or tetracyanoethylene with diazomethane².



A large number of substituted 1,1,2,2-tetracyanocyclopropanes are available by the Wideqvist reaction^{3,4}, in which a carbonyl compound is reacted with two equivalents of bromomalononitrile. A similar cyclopropanation procedure was reported by Hart^{5,6}.

In a previous report, we described the novel formation of a cyclopropane ring in the reaction of 1,1-diethoxy-2-bromoethylene with ethyl α -cyanoacrylate.⁷ A zwitterionic tetramethylene is formed by reaction of the electron-rich olefin with the electron-poor olefin. The expected cyclobutane cycloadduct is not formed. Instead, elimination of the bromide anion takes place with formation of a cyclopropane ring. Dealkylation of the dialkoxycarbocation by Br^- leads to an ester substituent. A similar reaction has been described by Scheeren: β,β -dicyanostyrene with 1,1-diethoxy-2-chloroethylene also yield a cyclopropane derivative as reaction product⁸.

In the present work, we react bromoketene acetals with tetracyanoethylene to lead to tetracyanocyclopropanecarboxylates.

Results and Discussion


1,1-Dialkoxy-2-bromoethylenes (1 or 2) react with tetracyanoethylene (TCNE 3) at -10°C to form methyl (6) or ethyl 1,1,2,2-tetracyanocyclopropane carboxylate (7), respectively. In analogy to the α -cyanoacrylate case, the following mechanism is proposed:



The initially formed 1,4-zwitterion 4 undergoes intramolecular elimination of bromide to form the dialkoxycation 5, which in turn undergoes dealkylation to form cyclopropane 6 or 7 and alkyl bromide. When the reaction is carried out in THF as solvent, a large quantity of poly-THF is formed together with the desired cyclopropane. This is additional evidence for the formation of 1,4-zwitterion in the course of reaction, which can initiate the cationic polymerization of THF⁹.

1,1-Diethoxy-2-bromopropene was much less reactive than 1,1-diethoxy-2-bromoethylene. Even at room temperature, no reaction occurred when 1,1-diethoxy-2-bromopropene was mixed with TCNE in a 1:1 molar ratio in THF.

Cyclic ketene acetals also undergo the reaction. When 2-(bromomethylidene)-1,3-dioxolane (8) or 2-(bromoethylidene)-1,3-dioxolane (9) is reacted with TCNE, a high yield of 2-bromoethyl 1,1,2,2-tetracyanocyclopropanecarboxylate 10 or 11 is obtained. The cyclic ketene acetal 8 has a tendency to undergo spontaneous cationic polymerization and the cyclopropane was always contaminated by the homopolymer of 8.



In these cases, the extra methyl group on the double bond of the ketene acetal did not reduce the reactivity significantly.

All 1,1,2,2-tetracyanocyclopropanecarboxylates (6, 7, 10, 11) were very sensitive to base or nucleophile. Saponification to carboxylic acid has failed so far. Even weak bases such as silver acetate, attacked the ring or the cyano groups. We were unable to obtain the vinyl ester derivatives in attempted dehydrobrominations of 10 and 11.

Experimental Section

Instrumentation. All melting points were obtained from a Thomas-Hoover capillary melting point apparatus. 1H -NMR and ^{13}C -NMR spectra were taken on a Bruker WM 250 nuclear magnetic resonance spectrometer at 250 MHz. Infrared spectra were recorded on a Perkin-Elmer 983 spectrometer. Elemental analyses were performed by Desert Analytics, Tucson, AZ.

Materials. Tetrahydrofuran was purified by refluxing with LiAlH_4 and distilled. Tetracyanoethylene (3) was purchased from Aldrich and was purified by two successive recrystallizations from chlorobenzene and sublimations (125°C , 0.5 mmHg).

1,1-Dimethoxy-2-bromoethylene (1) was obtained by dehydrobromination of 1,1-dibromo-2,2-dimethoxyethane according to a literature procedure¹⁰. Dibromoacetal was synthesized by bromination of 2-bromo-1,1-dimethoxyethane with bromine and calcium carbonate in dry diethyl ether.

1,1-Diethoxy-2-bromoethylene (2) was prepared following a similar procedure as for 1,1-dimethoxy-2-bromoethylene (1).

Methyl 1,1,2,2-tetracyanocyclopropane carboxylate (6). TCNE (2.56 g, 20 mmole) was dissolved in 30 ml of tetrahydrofuran. The solution under nitrogen was cooled to -10°C in dry ice-ethanol-water. Freshly distilled 1,1-dimethoxy-2-bromoethylene (4.18g, 25 mmole) was added slowly with stirring, and the reaction mixture was stirred for 1 hour. Solvent and methyl bromide were then evaporated by rotary evaporator. The obtained crude product was washed with n-hexane and purified by column chromatography (column size: 4.0 cm/40 cm, diameter/height, adsorbent: Silica Gel, 70-230 mesh, 60Å, eluent: ethylacetate/n-hexane, 40/60, V/V). Yield: 3.52 g (88%). $\text{Mp} = 175^\circ - 176^\circ\text{C}$ (dec.). IR (KBr): 3051 (CH, cyclopropane), 2261 (CN), 1752 (ester); $^1\text{H-NMR}$ (acetone- d_6): δ 3.93 (s, $-\text{COOCH}_3$) 4.55 (s, CHCOO-). Anal. Calcd for $\text{C}_5\text{H}_4\text{N}_4\text{O}_2$: C, 53.89; H, 2.00; N, 27.94. Found: C, 53.66; H, 1.95; N, 27.86.

Ethyl 1,1,2,2-tetracyanocyclopropane carboxylate (7) was synthesized using a similar procedure as for cyclopropane 6 from TCNE and 1,1-diethoxy-2-bromoethylene. Yield: 3.62 g (85%). $\text{Mp} = 158^\circ - 159^\circ\text{C}$ (dec.). IR (KBr): 3057 (CH, cyclopropane), 2263 (CN), 1753 (ester); $^1\text{H-NMR}$ (acetone- d_6): δ 1.33 (t, $-\text{CH}_3$), 4.38 (q, $-\text{CH}_2-$), 4.52 (s, CHCOO-); $^{13}\text{C-NMR}$ (acetone- d_6): δ 14.2 (s, $-\text{CH}_3$), 22.3 (s, $>\text{CH}<$), 38.5 (s, $>\text{CH-}$), 64.5 (s, $-\text{OCH}_2-$), 108.6 (s, cis-CN) 110.8 (s, trans-CN), 161.7 $-\text{COO-}$ (s).

-COO-). Anal. Calcd for $C_{10}H_6N_4O_2$: C, 56.08; H, 2.82; N, 26.16. Found: C, 56.03; H, 2.67; N, 26.14.

2-(Bromomethyldene)-1,3-dioxolane (8) was obtained by dehydrobromination of 2-(dibromomethyl)-1,3-dioxolane as described by McElvain¹¹. 2-(Dibromomethyl)-1,3-dioxolane was prepared by an alcohol exchange between 1,1-dibromo-2,2-diethoxyethane and glycol¹⁰.

2-(Bromoethyldene)-1,3-dioxolane (9) was synthesized using a procedure similar to that for 2-(bromomethylidene)-1,3-dioxolane (8) from 2-(β,β -dibromoethyl)-1,3-dioxolane which was obtained from propionaldehyde by two successive bromination and alcohol exchange reaction. The total yield of the reactions was 66%. Bp = 59° - 60° C/0.5 mmHg. This compound crystallized in the refrigerator. IR (neat): 2968 (CH), 1644 (C=C); 1 H-NMR ($CDCl_3$): δ 2.10 (s, $-CH_3$), 4.32 (m, $-CH_2-CH_2-$). Anal. Calcd. for $C_5H_7BrO_2$: C, 33.52; H, 3.91; Br, 44.69. Found: C, 33.39; H, 3.88; Br, 44.48.

Bromoethyl 1,1,2,2-tetracyanocyclopropanecarboxylate (10). TCNE (3.84 g, 30 mmole) was dissolved in 30 ml of tetrahydrofuran. The solution under nitrogen atmosphere was cooled in ice bath. 2-(Bromomethylidene)-1,3-dioxolane (6.6 g, 40 mmole) was added slowly with stirring under nitrogen, and the reaction mixture was stirred for 2 hr. Homopolymer of 2-(bromomethylidene)-1,3-dioxolane was separated by filtration and the resulting filtrate was concentrated by rotary evaporator. The obtained crude product was washed with n-hexane and purified by column chromatography. After washing with cold diethyl ether, the white crystals were dried under vacuum. Yield: 6.32 g (72%). Mp = 152° - 153° C. IR (KBr): 3051 (CH, cyclopropane), 2265 (CN), 1746 (ester); 1 H-NMR (acetone- d_6): δ 3.72 (t, $-CH_2Br$), 4.67 (t, $-COOCH_2-$); ^{13}C -NMR (acetone- d_6): δ 22.5 (s, $>C<$), 29.2 (s, $-CH_2Br$), 38.2 (s, $>CH-$), 67.9 (s, $-OCH_2-$), 108.5 (s, cis-CN), 110.7 (s, trans-CN), 161.9 (s, $-COO-$).

Anal. Calcd for $C_{10}H_3BrN_4O_2$: C, 40.96; H, 1.70; Br, 27.30; N, 19.11. Found: C, 41.05; H, 1.65; Br, 27.18; N, 19.16.

Bromoethyl 1,1,2,2-tetracyano-3-methylcyclopropanecarboxylate (11) was synthesized according to a procedure similar to that of bromoethyl 1,1,2,2-tetracyanocyclopropanecarboxylate (10) from TCNE (3.84 g, 30 mmole) and 2-(bromoethylidene)-1,3-dioxolane (7.17 g, 40 mmole). Yield: 8.3 g (90%). Mp=179°-180°C (dec). IR (KBr): 2964 (CH), 2257 (CN), 1745 (ester); 1H -NMR (DMSO- d_6): δ 1.77 (s, -CH₃), 3.71 (t, -CH₂Br), 4.54 (t, -COOCH₂-); ^{13}C -NMR (DMSO- d_6): δ 16.4 (s, -CH₃), 27.5 (s, >C(CN)₂), 29.2 (s, -CH₂Br), 43.8 (s, >C<), 68.2 (s, -OCH₂), 109.1 (s, cis-CN), 110.2 (s, trans-CN), 163.3 (s, -COO-). Anal. Calcd for $C_{11}H_7BrN_4O_2$: C, 42.99; H, 2.28; Br, 26.05; N, 18.24. Found: C, 43.12; H, 2.26; Br, 26.00; N, 18.14.

Acknowledgement

We are deeply indebted to the Office of Naval Research for partial financial support of this work and to Dr. Anne Padias for helpful discussions.

References

1. Scribner, R.M.; Sausen, G.N.; Prichard, W.W. J. Org. Chem., 1960, 25, 1440.
2. Bestus, J.; Castells, J. J. Proc. Chem. Soc. (London), 1962, 216.
3. Ramberg, L; Widequist, S. Arkiv Kemi., 1937, 12A(22).
4. Ramberg, L; Widequist, S. ibid, 1941, 14B(37).
5. Hart, H; Freeman, F. J. Org. Chem., 1963, 28, 1220.
6. Hart, H; Kim, Y.C. ibid, 1966, 31, 2784.
7. Hall, Jr., H.K.; Padias, A.B.; Deutschman, Jr., A.; Westerman, I.J. J. Org. Chem., 1979, 44, 2038.
8. Scheeren, W.J. Rec. Trav. Chim. Pays-Bas., 1986, 105, 75.
9. Padias, A.B.; Hall, Jr., H.K. Acc. Chem. Res., 1990, (in press).

10. Beyerstedt, F.; McElvain, S.M. J. Am. Chem. Soc., 1937, 59, 2266.

11. McElvain, S.M.; Curry, M.J. ibid, 1948, 70, 3781.