

②

REPORT DOCUMENTATION PAGE

AD-A220 729

1b. RESTRICTIVE MARKINGS

3. DISTRIBUTION/AVAILABILITY OF REPORT

4. PERFORMING ORGANIZATION REPORT NUMBER(S)

04

5. MONITORING ORGANIZATION REPORT NUMBER(S)

6a. NAME OF PERFORMING ORGANIZATION

Northwestern University

6b. OFFICE SYMBOL
(If applicable)

7a. NAME OF MONITORING ORGANIZATION

6c. ADDRESS (City, State and ZIP Code)

Department of Chemistry
Evanston, Illinois 60208-3113

7b. ADDRESS (City, State and ZIP Code)

8a. NAME OF FUNDING/SPONSORING
ORGANIZATION

Office of Naval Research

8b. OFFICE SYMBOL
(If applicable)

9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

8c. ADDRESS (City, State and ZIP Code)

Department of the Navy
Arlington, Virginia 22217

10. SOURCE OF FUNDING NOS.

PROGRAM
ELEMENT NO.PROJECT
NO.TASK
NO.WORK UNIT
NO.

11. TITLE (Include Security Classification)

Molecular Structure of $[(O)_2Re(bipyridine)]$..12. PERSONAL AUTHOR(S) Robert L. Blackburn, Lisa M. Jones, M. S. Ram, Michael Sabat and
Joseph T. Hupp

13a. TYPE OF REPORT

Technical Report

13b. TIME COVERED

FROM _____ TO _____

14. DATE OF REPORT (Yr., Mo., Day)

3/30/90

15. PAGE COUNT

7

16. SUPPLEMENTARY NOTATION

17. COSATI CODES

FIELD GROUP SUB. GR.

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

An x-ray crystal structure investigation has revealed that the stable octahedral complex $[(O)_2Re(2,2'bipyridine)(pyridine)_2] \cdot (ClO_4)$ exists in a *cis*-dioxo ligand configuration. The new complex represents only the second confirmed example of the *cis*-dioxo structure in hexacoordinate d^2 transition-metal chemistry. The *cis* assignment provides an explanation for the marked electrochemical and spectral differences between the bipyridyl complex and the previously known *trans*-dioxo rhenium tetrapyridine complex.

20. DISTRIBUTION/AVAILABILITY OF ABSTRACT

UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT. ☐ DTIC USERS ☐

21. ABSTRACT SECURITY CLASSIFICATION

22a. NAME OF RESPONSIBLE INDIVIDUAL

Joseph T. Hupp

22b. TELEPHONE NUMBER
(Include Area Code)

(708) 491-3504

22c. OFFICE SYMBOL

OFFICE OF NAVAL RESEARCH

TECHNICAL REPORT

FOR

Contract N0014-88K-0369

Technical Report No. 4

Molecular Structure of $[(O)_2Re(bipyridine)(pyridine)_2](ClO_4)$:

An Unusual Example of a d^2 Metal Complex with

a cis-Dioxo Ligand Configuration

by

Robert L. Blackburn, Lisa M. Jones, M. S. Ram,
Michal Sabat and Joseph T. Hupp

Department of Chemistry
Northwestern University
Evanston, Illinois 60208-3113



Reproduction in whole, or in part, is permitted for any purpose of
the United State Government.

Accepted for publication in Inorganic Chemistry.

STATEMENT "A" per Robert Nowak
ONR/Code 1113ES
TELECON

4/10/90

VG

Accession For	
NTIS	☒
DTIC	☒
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Special
A-1	

Dioxo rhenium(V) species, especially with pyridyl-type ligands, have attracted significant interest recently - both on account of their electrochemical properties¹ and their photophysical and photochemical reactivity characteristics.^{2,3} We have been examining these species from the point of view of multi-electron transfer kinetics (i.e. $\text{Re(V)} = \text{Re(III)}$ and $\text{Re(V)} = \text{Re(II)}$), especially at electrochemical interfaces.⁴ In the course of those studies we have had occasion to introduce relatively minor modifications to the ligand environment. One of these - the replacement of a pair of pyridine(py) ligands by 2,2'-bipyridine(bpy) in the prototypical complex, $\text{trans}-(\text{O})_2\text{Re(py)}_4^+$ - had profound chemical consequences. Formal potentials shifted by up to 600 mV, photophysical activity disappeared, and a rare display of three-electron redox chemistry emerged at high pH's. Furthermore, the kinetics of the multi-electron electrochemical reduction were greatly accelerated. Finally, there were marked changes in NMR, Raman UV-vis and infrared spectra.⁴

On the basis of extensive circumstantial evidence we were forced to conclude that the changes were brought about by trans to cis isomerization upon bpy substitution. (The conclusion was particularly attractive, however, because it could account for the rather remarkable kinetic effects.) It was noted, nevertheless, that a rigorous structural assignment was impossible based upon either magnetic or vibrational spectroscopy - the reason being that both *cis*- and *trans*- $(\text{O})_2\text{Re(bpy)(py)}_2^+$ would possess C_2V symmetry.^{4,5} Furthermore, there were theoretical reasons to doubt the *cis* geometry.⁶

We now report, based upon x-ray crystal structure investigations,⁷ that the product of bipyridyl substitution at $\text{trans}-(\text{O})_2\text{Re(py)}_4^+$ is indeed *cis*- $(\text{O})_2\text{Re(bpy)(py)}_2^+$. To our knowledge this is one of only two confirmed


examples of a hexacoordinate d^2 metal complex with *cis*-dioxo structure.^{8,9} (It should be noted, however, that two additional examples have been proposed by Meyer and co-workers^{10,11} based on polypyridyl complexation of oxidation state VI of either osmium or ruthenium. The osmium complex is moderately stable and has been isolated as a perchlorate salt;¹⁰ the Ru species displays only a transient solution existence.¹¹ Neither has been confirmed by crystallographic measurements. Nevertheless, in view of the present findings the earlier structural assignments almost certainly are correct.)

For the current work a powdered sample of *cis*- $[(O)_2Re(bpy)(py)_2](ClO_4)$ was prepared and purified as described by Ram, et al.⁴ Red crystals of x-ray quality (ca. 0.4 mm diameter) were obtained (~3 days) by dissolution of the powder in an acetone/ethyl acetate mixture, followed by cooling to $-30^\circ C$.¹²

An ORTEP drawing of the complex cation is shown in figure 1. The Re atom is present within a distorted octahedral environment. The *cis*-(O)₂Re moiety is characterized by Re=O bond lengths of 1.733(8) Å and 1.736(7) Å and an O=Re=O bond angle of 121.4(4). The Re=O bond lengths are only slightly shorter than those found in the d^2 trans complexes $[(O)_2Re(py)_4]Cl$ and $[(O)_2Re(ethylenediamine)_2]Cl$.¹³ They also compare very well to those found in a trigonal bipyramidal *cis*-(O)₂Re species.¹⁴ The O=Re=O angle, however, is much smaller in $(O)_2Re(bpy)(py)_2$ than in the five-coordinate structure ($138.7(6)^\circ$).¹⁴ On the other hand, it is appreciably larger than in most d^2 *cis*-dioxo complexes.^{15,16} The bipyridine N1-Re-N1' "bite" angle is $70.6(3)^\circ$, a quite typical value for bpy complexes. Likewise the inter-ring "twist" of 7.2° is typical of transition-metal bpy complexes.¹⁷ Finally, it is worth noting that the pyridyl ligands are tilted significantly towards the *cis*-(O)₂Re moiety. Thus the dihedral angles between the (O)₂Re plane and the N1

and N3 rings are 74.6° and 74.8° respectively. (The N1-Re-N3 angle itself is $169.0(3)^\circ$.) At present, we have no explanation for this peculiar geometry.

In view of Mingos' work,⁶ the cis configuration is perhaps not the most stable form thermodynamically for $(O)_2\text{Re}(\text{bpy})(\text{py})_2^+$.¹⁸ Evidence of chemical instability comes from solution studies⁴ and from the detection of $\text{pyridineH}^+\text{ClO}_4^-$ in the sample. Obviously, the appearance of the cis structure at the expense of the trans would then reflect a kinetic preference at some stage in the substitution process. Although we have not pursued the issues of substitution kinetics or mechanism, one simple sequence which might account for the isomerization would be the following:¹⁹



The crucial second step presumably would be driven by the need to relieve steric crowding at the α -sites on adjacent pyridine ligands.


Acknowledgement

We thank the Office of Naval Research for support of this work.

Supplementary Material: Tables of positional parameters, thermal parameters, interatomic distances and bond angles for the Re cation and a table of interatomic distances for the disordered perchlorate and acetone are available. Ordering information is given on any current masthead.

References

1. Pipes, D. W.; Meyer, T. J., Inorg. Chem., 1986, 25, 3256.
2. (a) Winkler, J. R.; Gray, H. B., J. Am. Chem. Soc., 1983, 105, 1371.
(b) Winkler, J. R.; Gray, H. B., J. Am. Chem. Soc., 1985, 24, 346. (c)
Thorp, H. H.; Van Houten, J.; Gray, H. B., Inorg. Chem., 1989, 28, 889.
(d) Thorp, H. H.; Kumar, C. V.; Turro, N. J.; Gray, H. B., J. Am. Chem. Soc., 1989, 111, 4364.
3. Newsham, M. D.; Giannelis, E. P.; Pinnavia, T. J.; Nocera, D. E., J. Am. Chem. Soc., 1988, 110, 3885.
4. Ram, M. S.; Johnson, C. S.; Blackburn, R. L.; Hupp, J. T., Inorg. Chem., 1990, 29, 238.
5. NMR studies did, however, enable us to rule out an even lower symmetry *cis* structure:



6. Mingos, D. M. P., J. Organomet. Chem., 1979, 179, C29.
7. **Crystal data:** formula = $[\text{ReO}_2(\text{py})_2(\text{bpy})]\text{ClO}_4 \cdot 0.33(\text{pyH})\text{ClO}_4 \cdot 0.33(\text{CH}_3)_2\text{CO}$,
M = 711.25, rhombohedral, space group $\bar{R}3$ (no. 148), $a = 24.627(5) \text{ \AA}$,
 $c = 19.748(5) \text{ \AA}$, $V = 10,372(7) \text{ \AA}^3$, $Z = 18$. The structure was solved by
Patterson and Fourier techniques and refined to $R(F)$ and $R_w(F)$ of 0.052
and 0.070, respectively. 2260 absorption corrected reflections with
 $I > 3\sigma(I)$ were measured on an Enraf Nonius CAD4 diffractometer up to
 $2\theta_{\text{max}} = 55^\circ$ (Mo $K\alpha$ radiation, $\lambda = 0.71069 \text{ \AA}$).

8. The sole previous example is potassium tris(acetato)dioxoosmate(VI). Here the *trans* acetato ligands are monodentate while the remaining acetate is bidentate. It has been argued that the *cis*-dioxo configuration is obtained because the molecule effectively conforms to pseudo trigonal bipyramidal (*tbp*) symmetry, with the bidentate acetate effectively occupying only one coordination site. (Genuinely pentacoordinate *cis*-dioxo species are well known.¹⁵) Pseudo *tbp* symmetry was also invoked to explain the rather large O=Os=O bond angle (125°) (Behling, T.; Capparelli, M. V.; Skapski, A. C.; Wilkinson, G., Polyhedron, 1982, 1, 840). Our results for the more nearly octahedral rhenium complex would seem to be at variance with these "geometrical" explanations.
9. For a review of this area, see: Holm, R. H., Chem. Rev., 1987, 1401.
10. Dobson, J. C.; Takeuchi, K. J.; Pipes, D. W.; Geselowitz, D. A.; Meyer, T. J., Inorg. Chem., 1986, 25, 2357.
11. Dobson, J. C.; Meyer, T. J., Inorg. Chem., 1988, 27, 3283.
12. Identical IR spectra were obtained in the metal-oxo stretching region for crystalline and powdered samples.
13. Lock, C. J. L.; Turner, G., Acta Cryst., 1978, B34, 923.
14. Ciani, G. F.; D'Alfonso, G.; Romiti, P. F.; Sironi, A.; Freni, M., Inorg. Chim. Acta, 1983, 72, 29.
15. Nugent, W. A.; Mayer, T. M., "Metal-Ligand Multiple Bonds", John Wiley and Sons, New York, 1988.
16. Bond angles for the latter almost always fall in the 102-112° range.¹⁵
17. See, for example: Ghosh, M.; Siddhartha, R., Acta Cryst., 1983, C39, 1367.

18. The studies by Meyer and co-workers^{10,11} of $(O)_2Ru(bpy)_2^{2+}$ and $(O)_2Os(bpy)_2^{2+}$ clearly support this contention. A referee has pointed out, however, that in $(O)_2Re(bpy)(py)_2^+$ the π -donating oxo ligands might be sufficiently stabilized by the π -accepting bipyridyl ligand in a trans configuration (relative to the oxo groups) to render the cis dioxo species the more stable isomer. Obviously, a definitive resolution will require the synthesis of one or both trans isomers - a goal which has thus far proven elusive.
19. A reviewer has argued (based on the work of Kashani and Murmann, Int. J. Chem. Kinet., 1985, 17, 1005) that a six coordinate intermediate, $Re(py)_3(H_2O)(O)_2^+$, is more likely. (It is worth noting that neither intermediate has been isolated.) While we have no strong feelings on this point, we are inclined to favor the five-coordinate pathway. One piece of evidence in support of isomerization via a pentacoordinate species is a recently discovered route to $cis-(O)_2Re(bpy)(py)_2^+$ through a known²⁰ five-coordinate precursor, $(O)_2Re(py)_2I$ (M. S. Ram, unpublished).
20. Freni, M.; Giusto, D.; Romiti, P.; Minghetti, G., Gazz. Chim. Ital., 1969, 99, 286.

Figure caption: ORTEP drawing of *cis*-(O)₂Re(bpy)(py)₂⁺.

