

20030206263

Best Available Copy

AD-A220 929

2

5E1

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION (U) DTIC			1b. RESTRICTIVE MARKINGS N/A	
2a. SECURITY CLASSIFICATION AUTHORITY N/A			3. DISTRIBUTION/AVAILABILITY OF REPORT Distribution Unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE N/A			5. MONITORING ORGANIZATION REPORT NUMBER(S) N/A	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) N/A			7a. NAME OF MONITORING ORGANIZATION Office of Naval Research	
6. NAME OF PERFORMING ORGANIZATION The Agouron Institute		6b. OFFICE SYMBOL (If applicable) N/A	7b. ADDRESS (City, State, and ZIP Code) 800 North Quincy Street Arlington, Virginia 22217-5000	
6c. ADDRESS (City, State, and ZIP Code) 505 Coast Boulevard South La Jolla, California 92037		8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		
8b. OFFICE SYMBOL (If applicable) ONR		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00017-87-K-0322		
8c. ADDRESS (City, State, and ZIP Code) 800 North Quincy Street Arlington, Virginia 22217-5000		10. SOURCE OF FUNDING NUMBERS		
		PROGRAM ELEMENT NO 61153N	PROJECT NO RR04106	TASK NO 441h002
11. TITLE (Include Security Classification) (U) Genetic and Molecular Basis of Marine Fouling				
12. PERSONAL AUTHOR(S) Silverman, Michael R.				
13a. TYPE OF REPORT Final	13b. TIME COVERED FROM 4/01/87 TO 3/31/90	14. DATE OF REPORT (Year, Month, Day) 4/17/90	15. PAGE COUNT 3	
16. SUPPLEMENTARY NOTATION				
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP		
06	03		Surface recognition; gene regulation; bacterial differentiation.	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)				
The goal of our research is to use molecular and genetic techniques to investigate the process of bacterial attachment to and colonization of surfaces in the marine environment. We are focussing on exploring the mechanism which controls surface-induced swarmer cell differentiation of <i>Vibrio parahaemolyticus</i>. Gene fusions which couple transcription of swarmer cell genes, <i>laf</i>, to luminescence reporter genes, <i>lux</i>, have been used to analyze how environmental signals regulate differentiation, and a novel mechanism of surface recognition involving a tactile sensor and an osmotic sensor has been discovered. Work is continuing to develop a refined understanding of the sensors and other elements in the information transduction circuit which controls expression of swarmer cell genes.				
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT ☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT ☐ DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION (U)	
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. M. Marron			22b. TELEPHONE (Include Area Code) (202) 696-4760	22c. OFFICE SYMBOL ONR

DD Form 1473, JUN 86

Previous editions are obsolete

SECURITY CLASSIFICATION OF THIS PAGE

S/N 0102-LF-014-6603

00 04 99 148

DATE: April 17, 1990

FINAL REPORT ON CONTRACT N00014-87-K-0322, R&T CODE 441h002

PRINCIPAL INVESTIGATOR: Michael R. Silverman
CONTRACTOR: The Agouron Institute, La Jolla, California 92037
CONTRACT TITLE: Genetic and Molecular Basis & Marine Fouling
PERIOD OF PERFORMANCE: 4/1/87 - 3/1/90

RESEARCH OBJECTIVE: Our goal is to understand the molecular mechanisms which control microbial colonization of surfaces in the marine environment. We have been focussing on the surface-induced differentiation of *V. parahaemolyticus* because exploration of this adaptive response could reveal a mechanism for recognition of surface contact.

PROGRESS (YEAR 1, 2 AND 3): *V. parahaemolyticus* has two distinct cell types, the swimmer cell and the swarmer cell, which are adapted for survival in different habitats. The swimmer cell, produced when the bacterium is grown in a liquid medium, is a short rod with a single sheathed polar flagellum. The swarmer cell, produced when *V. parahaemolyticus* is grown in contact with surfaces, is greatly elongated and synthesizes, in addition to the polar flagellum, numerous lateral flagella which are responsible for translocation over surfaces. Differentiation to the swarmer cell appears to be an appropriate adaptation for life on surfaces because cells with lateral flagella adhere more firmly to surfaces and swarming expands the area of colonization on the surfaces. But, how does the bacterium recognize contact with a surface and activate the genetic program (*laf* genes) encoding the swarmer cell phenotype?

The transcription of genes encoding the swarmer cell phenotype has been analyzed with *laf::lux* gene fusion strains constructed in *V. parahaemolyticus* with transposon mini-Mulux. This approach simplified examination of gene regulation since light emission, encoded by the *lux* genes, was measured instead of complex morphological and behavioral events. Gene fusion strains were used to show that swarmer cell genes were induced by a variety of conditions, including a high viscosity environment or antibody cross-linking, which had in common the constraint of movement of the polar flagellum. To test the hypothesis that the polar flagellum is functioning as a tactile sensor which controls swarmer cell formation, we constructed a variety of mutations in genes encoding components of the polar flagellum (*fla*). The consequence of such mutations was the constitutive, surface-independent, expression of *laf* genes. So, the performance of the polar flagellum (*Fla*) is coupled to the transcription of the *laf* genes such that when *Fla* function is perturbed, either physically or genetically, swarmer cell genes are induced. Because the polar flagellum appears to be capable of sensing external forces influencing its motion, we suggest it is operating as a dynamometer.

Another environment input has been found to influence swarmer cell formation. In addition to stimulation of the tactile sensor (the polar flagellum), a second signal, iron limitation, is required for swarmer cell differentiation. Differentiation requires a large investment of cellular resources, and by basing the "decision" to differentiate on multiple inputs an appropriate response to a specific environmental condition could best be accomplished. We have also found that the polar (*Fla*) and lateral (*Laf*) flagellar systems

show behavioral coupling. The two appendages are assembled from different motor-propeller components, but chemotactic control of the behavior of swimmer and swarmer cells is controlled by one common information processing apparatus.

We have identified signals which induce swarmer cell differentiation and have discovered that the polar flagellum functions as a tactile sensor controlling differentiation. Research is now focussing on understanding how this sensor works at the molecular level. The polar flagellum can be expected to be very complex with many components involved in the assembly of the motor-propeller structure, in the energy transduction machinery driving propeller rotation, in the chemosensory system directing flagellar movement in response to environmental stimuli and also in the tactile sensor function. We are using mutants to attempt to separate tactile sensor function from the behavioral response function of the polar flagellum and to determine what component or specific flagellar activity is directly involved in controlling expression of the swarmer cell phenotype.

We have used mutant analysis to explore the mechanism of chemotactic control of the two flagellar systems of *V. parahaemolyticus*. Studying chemotaxis with *V. parahaemolyticus* is complicated because it is capable of producing two propulsive organelles, Fla (polar flagellum) and Laf (lateral flagella) either or both of which could be responsible for movement toward a particular chemical attractant or away from a particular chemical repellent. We could eliminate this confusion by constructing swim-only, Fla⁺ Laf⁻, and swarm-only, Fla⁻ Laf⁺, strains for analysis of chemotaxis in capillary assays. With these specially constructed strains, we have shown that swim-only and swarm-only bacteria respond to the same spectrum of chemical substances. However, swim-only bacteria chemotact in a low viscosity medium while swarm-only bacteria chemotact best in a high viscosity environment. Furthermore, we have cloned and mutated a chemotaxis gene locus in *E. coli* and then recombined these defects into the chromosomes of various swim-only and swarm-only strains of *V. parahaemolyticus*. The results of introducing these *che* gene defects is the disruption of chemotactic control of both the swimming and swarming organelles. So, the two propulsive systems are controlled by a single chemical sensing mechanism. Put another way, *V. parahaemolyticus* has two structurally separate propulsive devices which are directed by one "brain".

We are particularly interested in extending the analysis of chemotaxis mutants. These can be constructed by localized mutagenesis of the cloned *che* genes and subsequent transfer of mutations to *V. parahaemolyticus* by a gene replacement procedure. It is known from analysis of paralyzed (Mot⁻) mutants that flagellar rotation is required for tactile sensing, and with Che⁻ mutants it should be possible to determine if chemosensory function is also necessary. And, Che⁻ mutants are usually locked into either a clockwise or counter-clockwise rotational mode so the influence of the direction of propeller rotation on sensing can be examined. Another approach is to search for genes whose products directly regulate the expression of swarmer cell genes. Swarmer cell genes, *laf*, appear to be regulated by negative rather than by positive control of transcription since mutants with defects in the tactile sensor genes are constitutive for *laf* expression rather than being uninducible which would be the consequence of a defect in a positive effector of transcription. We are searching for genes encoding a repressor of *laf* transcription by programming expression of cloned genes positioned on an expression vector *in trans* in *V. parahaemolyticus*.

PUBLICATIONS FROM THIS PROJECT:

1. McCarter, L. and M. Silverman. 1987. Phosphate regulation of gene expression in *V. parahaemolyticus*. *J. Bacteriol.* 169:341-3449.
2. Belas, R., D. Bartlett and M. Silverman. 1988. Cloning and gene replacement mutagenesis of a *Pseudomonas atlantica* agarase gene. *Appl. and Env. Microbiol.* 54:30-37.
3. McCarter, L., M. Hilmen and M. Silverman. 1988. Flagellar dynamometer controls swarmer cell differentiation of *V. parahaemolyticus*. *Cell* 54:345-351.
4. Simon, M., R. Belas, L. McCarter, D. Bartlett and M. Silverman. 1988. Microbial role in biodeterioration: Advanced Techniques Applicable to the Indian Ocean, Oxford and IBH Publishing, New Delhi, India, Eds. American Institute of Biological Sciences, p. 265.
5. McCarter, L. and M. Silverman. 1989. Iron regulation of swarmer cell differentiation of *V. parahaemolyticus*. *J. Bacteriol.* 171:731-736.
6. Sar, N., L. McCarter, M. Simon and M. Silverman. 1990. Chemotactic control of the two flagellar systems of *V. parahaemolyticus*. *J. Bacteriol.* 172:334-341.
7. McCarter, L. and M. Silverman. 1990. Surfaced-induced differentiation of *Vibrio parahaemolyticus*. *Mol. Microbiol.* (in press).
8. Crawford, I., C. Han and M. Silverman. 1990. Sequence and features of the tryptophan operon of *Vibrio parahaemolyticus*. *Sequence* (in press).
9. Silverman, M., R. Showalter and L. McCarter. 1990. Genetics of *Vibrio*. *Meth. Enzymol.* (in press).

INVENTIONS:

None

PERSONNEL SUPPORT:

This contract supported the salary of Research Scientist Dr. Linda McCarter.

Account No.	
THIS CONTRACT	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Date	
Approved	
Dis	
A-1	