

AD-A221 733

ALL COPY

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER No. 35	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Quantum Statistical Properties of an FEL Amplifier	5. TYPE OF REPORT & PERIOD COVERED Technical Report 10/01/84 - 09/30/85	
6. PERFORMING ORG. REPORT NUMBER		
AUTHOR(s) G. Dattoli, J. Gallardo, A. Renieri and M. Richetta		8. CONTRACT OR GRANT NUMBER(s) N00014-80-C-0308
PERFORMING ORGANIZATION NAME AND ADDRESS University of California Quantum Institute Santa Barbara, CA 93106		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61153N; RR011-07-06; NR603-001
CONTROLLING OFFICE NAME AND ADDRESS ONR 1030 E. Green St. Pasadena, CA 91106		12. REPORT DATE 1985
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES
15. SECURITY CLASS. (of this report) Unclassified/Unlimited		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES IEEE Journal of Quantum Electronics Special Issue on FELs, 1985		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Statistical, Properties, Amplifier.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) We discuss the problem of photon quantum statistics of a single particle Free Electron Laser (FEL) amplifier in the small-signal cold beam regime to first order in the electron quantum recoil. The initial radiation wave is an arbitrary coherent state. We show that Glauber coherence is not preserved by the FEL interaction if the initial coherent state is not the vacuum even if we neglect the electron quantum recoil (absence of gain). We evaluate the first two moments of the final photon distribution and find sub (super)-poissonian photon statistics for negative (positive) resonance parameter.		

DTIC
ELECTE
MAY 21 1990

ADVISORY GROUP
DEFINITION
FEB 1985
ON ELECTRON DEVICES

Quantum Statistical Properties of an FEL Amplifier

G.Dattoli, J.Gallardo, A.Renieri, M.Richetta

Submitted to IEEE

QIFEL035/84

Nov. 2, 1984

Quantum Statistical Properties of an FEL Amplifier

G. Dattoli* , J. Gallardo

Quantum Institute
University of California
Santa Barbara, CA 93106

and

A. Renieri, M. Richetta+

Comitato Nazionale Energie Nucleare
Centro di Frascati
C.P. 65, 00044 Frascati, Rome, Italy

* Permanent address ENEA, Frascati, Italy

+ ENEA student

Accession For	
NTIS	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution /	
Availability Codes	
Dist	Special
A-1	

ABSTRACT.

We discuss the problem of photon quantum statistics of a single particle Free Electron Laser (FEL) amplifier in the small-signal cold beam regime to first order in the electron quantum recoil. The initial radiation wave is an arbitrary coherent state. We show that Glauber coherence is not preserved by the FEL interaction if the initial coherent state is not the vacuum even if we neglect the electron quantum recoil (absence of gain). We evaluate the first two moments of the final photon distribution and find sub (super)-Poissonian photon statistics for negative (positive) resonance parameter.

The problem of Glauber coherence(1) in a FEL has been investigated by different groups in several papers.(2-7)

The first analysis was carried out using a group theoretical approach (2,3) in which the symmetry properties of the quantized non-relativistic FEL Hamiltonian was exploited to analyze both the single mode and multi-mode regimes. The main conclusion of these papers was that coherence was maintained. Therefore, provided the electron motion could be treated classically, the FEL interaction "rotates" a Glauber state into another one. This result is reminiscent of the Bloch-Nordsieck Theorem(8). The validity of the assumption of classical electron motion were discussed in Ref.(9).

A quantized dynamical approach to the FEL problem was started in Ref.(4); see also Ref.(7). They considered quantized electrons and electromagnetic fields in the framework of the interaction picture, showing that a FEL amplifier or oscillator is an interesting device that exhibits genuine quantum effects like sub or super-Poissonian photon statistics, and that the radiation field could be in a so-called squeezed state; also they conjectured that Glauber coherence is only preserved by a gain-less process.

A different dynamical method was proposed in Refs.(5,6) in which the Schrodinger equation of a single electron interacting with the combined radiation and wiggler field is solved to first order in the quantum electron recoil. In this way, all

the quantum effects previously mentioned, were recovered and furthermore we proved in a rigorous manner the conjecture of preservation of coherence to zeroth-order in the electron recoil. All these results were obtained starting from the vacuum state, therefore, we could say that departure from Poisson photon statistics and the presence of squeezed states are closely connected with the radiation field vacuum fluctuations.

In this letter we address the same problem with an arbitrary Glauber coherent state as an initial state. The single mode non-relativistic Hamiltonian describing the FEL process

$$H = \frac{P^2}{2m} + \hbar\omega \left(a_L^\dagger a_L + \frac{1}{2} \right) + \hbar\omega \left(a_U^\dagger a_U + \frac{1}{2} \right) + \hbar\Omega \left(a_L^\dagger a_U e^{-2ikz} + a_U^\dagger a_L e^{2ikz} \right) \quad (1)$$

The symbols used throughout the paper are summarized in Table 1.; we refer the reader to Refs. (5,10) for a discussion of their physical meaning.

Assuming an initial state consistent of an electron of momentum P_0 and a radiation field in a coherent state with average number of photon $|\alpha_0|^2$ and using the conservation laws of total number of photons and total momentum, we can write the state describing the evolution of the coupled electron and radiation field as

$$|\psi(t)\rangle = e^{i\frac{P_0 t}{2m\hbar} + i\omega t(n_0^\circ + 1)} \sum_{n=0}^{\infty} \frac{e^{-\frac{1}{2}|\alpha_0|^2}}{\sqrt{n!}} \alpha_0^n \sum_{\ell=n}^{\infty} C_\ell(t) |P_0 - 2\ell\hbar k, n+\ell\rangle \quad (2)$$

where l is the number of exchanged photons during the interaction and $\alpha_0 = |\alpha_0| e^{i\omega t}$. We recall that in the FEL process for each exchanged photon, the electron momentum changes by $2\hbar k$.

The coefficient $C_l(t)$ which is the amplitude probability for the process satisfy the following differential difference-equation (Raman-Nath equation)(5)

$$i C'_l(\tau) = (-\omega_0 + \epsilon l) C_l(\tau) + \Omega_R [\sqrt{m+l+1} C_{l+1} + \sqrt{m+l} C_{l-1}]$$

$$C_l(0) = \delta_{l,0} \quad (3)$$

the prime denotes derivative respect to $\tau = \frac{c}{L} t$ ($0 \leq \tau \leq 1$) where L is the length of the undulator and $\frac{L}{c}$ is the interaction time.

An exact analytical solution of Eq.(3) is not known; however, a perturbative solution in terms of the electron recoil parameter ϵ can be constructed. The zeroth-order solution has been found in Ref.(10) and reads,

$$C_l(\tau) \Big|_{\epsilon=0} = \exp \left\{ \frac{i\omega_0}{2} \int_0^\tau d\tau' |\alpha(\tau')|^2 \right\} \phi_n^l(\alpha(\tau)) \quad (4,a)$$

where the orthonormal functions ϕ_n^l are,

$$\phi_n^l(\alpha(\tau)) = \sqrt{\frac{n!}{(n+l)!}} e^{-\frac{1}{2}|\alpha|^2} (\alpha(\tau))^l L_n^l(|\alpha|^2) \quad (4,b)$$

and

$$\alpha(\tau) = (-i) e^{i\omega_0 \frac{\tau}{2}} \Omega_R \left(\frac{\sin \omega_0 \frac{\tau}{2}}{\omega_0/2} \right)$$

L_n^l are the Laguerre associated polynomials. Replacing the

expression of the coefficient $C_\ell(\tau)$ into Eq. 2 we obtain the zeroth-order state evolution of the FEL system. In Ref. 5 and 6 it was shown that $|\psi(\tau)\rangle_{\epsilon=0}$ is a coherent state in the Glauber sense under the assumption that the initial state is the vacuum.

We introduce creation and annihilation operators of the form

$$A = a_L e^{2ikz}, \quad A^\dagger = e^{-2ikz} a_L^\dagger$$

with $[A, A^\dagger] = 1$.

The action of them on the state $|\psi\rangle$ is

$$A |\psi\rangle = (\alpha(\tau) + \alpha_0 e^{2ikz}) |\psi\rangle \quad (5)$$

where we have used the relation $L_m^\ell(x) = L_{m-1}^\ell(x) + L_m^{\ell-1}(x)$.

From Eq.(5) we cannot conclude that $|\psi\rangle$ is a coherent state; it is clear that this comes from the presence of an average number of photons $|\alpha_0|^2$ in the initial state.

The average number of photons in the state $|\psi\rangle$ is

$$\langle \psi | A^\dagger A | \psi \rangle = \langle m + \ell \rangle_{\epsilon=0} = |\alpha_0|^2 + |\alpha|^2 \quad (6)$$

Next, we wish to compute the gain of this device to first order in the electron recoil and investigate the photon statistics of the final state of the radiation field.

The solution of our basic differential-difference equation (R.N.) has been derived in Ref.(10) and reads,

$$|\psi(\tau)\rangle = \exp \left\{ i \frac{W_0}{2} \int_0^\tau d\tau' |\alpha(\tau')|^2 + i \tau \omega (m_0^0 + 1) \frac{L}{c} + \frac{i P_0 L \tau}{2 m \hbar c} \right\} \times \quad (7)$$

$$\times \sum_{m=0}^{\infty} \frac{e^{-\frac{1}{2}|\alpha_0|^2}}{\sqrt{m!}} \alpha_0^m \sum_{\ell=-n}^{\infty} \sqrt{\frac{n!}{(n+\ell)!}} e^{-\frac{1}{2}|\alpha(\tau)|^2} (\alpha(\tau))^\ell [A_{\ell,n} + i D_{\ell,n}] |P_0 - 2\ell k \hbar, n+\ell\rangle$$

The coefficients $A_{\ell,n}$ and $D_{\ell,n}$ are lengthy expressions; in the small signal regime, i.e. first order in Ω_R they are given by,

$$A_{\ell,n} \simeq L_n^\ell(\cdot) + \frac{\epsilon}{|\alpha|} \frac{\partial |\alpha|^2}{\partial \omega_0} \left\{ (2\ell+1) |\alpha|^2 L_n^{\ell+1}(\cdot) - (2\ell-1)(n+\ell) L_n^{\ell-1}(\cdot) \right\} - \frac{\epsilon}{|\alpha|} \frac{\partial |\alpha|^2}{\partial \omega_0} \left\{ |\alpha|^4 L_n^{\ell+2}(\cdot) - (n+\ell)(n+\ell-1) L_n^{\ell-2}(\cdot) \right\} + \dots$$

and

$$D_{\ell,n} \simeq +\frac{1}{2} \tau \epsilon \left\{ (2\ell+1) |\alpha|^2 L_n^{\ell+1}(\cdot) + (2\ell-1)(n+\ell) L_n^{\ell-1}(\cdot) \right\} + \frac{\tau}{2} \epsilon \left\{ |\alpha|^4 L_n^{\ell+2}(\cdot) + (n+\ell)(n+\ell-1) L_n^{\ell-2}(\cdot) \right\} + \dots$$

the argument of the Laguerre polynomials $L_n^\ell(\cdot)$ is $|\alpha(\tau)|^2$. From the above expression for the state vector $|\psi\rangle$ we can evaluate the average value of photons in the final state,

$$\langle n+\ell \rangle_{\epsilon \neq 0} = \sum_{m=0}^{\infty} \frac{e^{-|\alpha_0|^2}}{m!} |\alpha_0|^{2m} \sum_{\ell=-n}^{\infty} \frac{m!}{(n+\ell)!} e^{-|\alpha|^2} |\alpha|^{2\ell} (n+\ell) [A_{\ell,n}^2 + D_{\ell,n}^2] \quad (8)$$

Using recurrence relations for the Laguerre polynomials and after simple but tedious manipulations we can write the coefficients $A_{\ell,n}$ and $D_{\ell,n}$ to first order in the recoil ϵ

$$D_{\ell,n}^2 \simeq 0$$

$$A_{l,n}^2 \simeq \left(L_n^{l(\cdot)} \right)^2 + \frac{2\epsilon}{|\alpha|} \frac{\partial |\alpha|^2}{\partial \omega_0} \left(L_n^{l(\cdot)} \right)^2 \left[l^2 + |\alpha|^2 - |\alpha|^4 \right] \\ - 2 L_n^{l(\cdot)} L_n^{l-1(\cdot)} \left[l^2 + l(n - |\alpha|^2) - n |\alpha|^2 \right] \Bigg\}.$$

The photon distribution function for the radiation field is given by

$$W(n, l, \epsilon) = \frac{e^{-|\alpha_0|^2}}{n!} |\alpha_0|^{2n} P_n(l, \epsilon)$$

where

$$P_n(l, \epsilon) = \frac{n!}{(n+l)!} e^{-|\alpha(\tau)|^2} |\alpha(\tau)|^{2l} A_{l,n}^2$$

which satisfy

$$\sum_{n=0}^{\infty} \sum_{l=n}^{\infty} W(n, l, \epsilon) = 1;$$

to show this we make use of the generating function(11)

$$\sum_{l=n}^{\infty} \lambda^l P_n(l, \epsilon) = e^{(\lambda-1)|\alpha|^2} L_n(-|\alpha|^2 \frac{(\lambda-1)^2}{\lambda})$$

and the explicit definition of the Laguerre polynomials.

Along the same lines the average value of photons in the final state for an arbitrary initial coherent state, reads

$$\langle n+l \rangle_{\epsilon \neq 0} = |\alpha_0|^2 + |\alpha|^2 - \epsilon \frac{\partial |\alpha|^2}{\partial \omega_0} \left(1 + 2|\alpha_0|^2 \right) \quad (9)$$

The $|\alpha|^2$ term in the right-hand-side of the above equation corresponds to spontaneous emission; the second term (gain) consists of the sum of two contributions: stimulated quantum vacuum field fluctuations and a classical stimulated one, which

is proportional to the average number of photons in the input laser wave.

Similarly, the second moment of the distribution $W(n, l, \epsilon)$ is given by,

$$\begin{aligned} \langle (n+l)^2 \rangle_{\epsilon \neq 0} = \langle (n+l)^2 \rangle_{\epsilon=0} - \\ - 2\epsilon \frac{\partial |\alpha|^2}{\partial W_0} \left\{ 2|\alpha_0|^4 + |\alpha_0|^2 (3 + 4|\alpha|^2) + 2|\alpha|^2 + \frac{1}{2} \right\} \end{aligned} \quad (10)$$

A radiation state in which the fluctuation of the number of photons $\langle \Delta N^2 \rangle$ is larger (smaller) than the average value of photons $\langle N \rangle$, corresponds to a photon distribution super (sub)-Poissonian (12). The sub-Poissonian behavior of the photon distribution is an effect which reveals the quantum properties of the radiation field and cannot be explained if we use a classical description of it. Hence, we evaluate the normalized second factorial moment

$$\langle \Delta(n+l)^2 \rangle_{\epsilon \neq 0} - \langle n+l \rangle_{\epsilon \neq 0}^2 = -2\epsilon \frac{\partial |\alpha|^2}{\partial W_0} \left\{ |\alpha_0|^2 (2|\alpha|^2 + 1) + |\alpha|^2 \right\} \quad (11)$$

This expression reduces to the known result of Refs. (4,5) for an initial vacuum state, i.e.

$$\langle \Delta l^2 \rangle_{\epsilon \neq 0} - \langle l \rangle_{\epsilon \neq 0}^2 = -2\epsilon |\alpha|^2 \frac{\partial |\alpha|^2}{\partial W_0}$$

In Fig. 1 we plot Eq.(11) as a function of the resonance parameter W_0 for several values of the initial mean number of photons $|\alpha_0|^2$. For an initial vacuum state and for an arbitrary initial coherent state the final stimulated radiation field is super-Poissonian for positive resonance parameter $W_0 > 0$ (above resonance), sub-Poissonian for $W_0 < 0$ and Poissonian at resonance $W_0 = 0$. This effect could be

detected in an amplification FEL experiment operating in the visible part of the spectrum.

Concerning the problem of squeezed states, since we know the expression of the amplitude probability in terms of $A_{0,n}$ and $B_{0,n}$, we can easily evaluate the quantities $\langle \Delta A_1^2 \rangle$ and $\langle \Delta A_2^2 \rangle$ where

$$A = \frac{A + A^*}{2}$$

$$A = \frac{A - A^*}{2i}$$

see also Refs.(6,9). In this way we could show that, in presence of gain, we have squeezed states for the FEL amplification. Anyway since A and A^* are not pure field operators but electron + field ones the interpretation of squeezing is doubtful and we do not dwell on this point.

In summary, we have shown that:

- a) Starting from an initial coherent state we cannot conclude that coherence is preserved at zero gain.
- b) The FEL amplified radiation field exhibits both squeezing and sub-Poissonian photon statistics⁺. These quantum effects could be susceptible to experimental verification.

⁺ The terms photon antibunching, bunching, have been used to describe sub-Poissonian, super-Poissonian, photon distributions.

ACKNOWLEDGEMENT.

We wish to acknowledge enlightening discussions with W.B.Colson.

(JG) was supported by The Office of Naval Research Contract
N00014-80-C-0308.

REFERENCES.

1. R. J. Glauber, "Quantum Optics and Electronics", Lectures at Les Houches, 1964. Editors C. de Witt, A. Blaudin and C. Conen-Tannoudji.
2. G.Dattoli, A.Renieri, F.Romanelli and R.Bonifacio, Opt.Comm. 34,240(1980)
3. G.Dattoli, A.Renieri and F.Romanelli, Opt.Comm.35,245(1980)
4. W. Becker, M. O. Scully and M. S. Zubairy, Phys. Rev. Lett. 48, 475 (1982); W. Becker, M. S. Zubairy, Phys. Rev.A25, 2200 (1982); W. Becker and J. K. McIver, Phys. Rev.A27, 1030 (1983); W. Becker, J. K. McIver, M. S. Zubairy and M. O. Scully, "Free Electron Generators of Coherent Radiation", C. A. Brau, S. F. Jacobs, M. O. Scully, Editors, Proceedings SPIE 453, 306 (1983).
5. G.Dattoli and A.Renieri, J.de Physique No.2,44,125(1983)
6. G. Dattoli and M. Richetta, Opt.Comm.50,165(1984)
7. W. B. Colson, Phys. Lett. 59A, 187 (1976), "Free Electron Laser Theory", PhD dissertation, Stanford University, Stanford, CA (1977); P. Bosco, W. B. Colson and R. A. Freedman, IEEE J. of Quantum Electron QE-19, 272 (1983).; J. M. J. Madey, J. Appl. Phys. 42, 1906 (1971); J. M. J. Madey and D. A. G. Deacon, "Cooperative Effects in Matter and Radiation", Editors, C. M. Bowden, D. W. Howgate and H. R. Robl (1977) Plenum Pub. Co.
8. F.Bloch and A.Nordsieck,Phys.Rev.52,54(1937);Bjorken and Drell, "Relativistic Quantum Fields",McGraw-Hill, New York (1965).
9. G. Dattoli and A. Renieri, "Laser Handbook", Vol. 4,

Editors: M.S.Stich and M.S.Bass (to be published); G.Dattoli,
A.Renieri and M.Richetta, "Quantum Free-Electron Laser
Statistics: A critical review", Brookhaven Conference on Free
Electron Generation of Extreme Ultraviolet Coherent Radiation,
September 1983, to be published by Optical Society of America.

10. F.Ciocci, G.Dattoli and M.Richetta, J.Phys.A17,1333(1984)
11. W.Becker,Opt.Comm.33,69(1980).
12. M.C.Teich, B.E.Saleh and D.Stoler, Opt.Comm.46,244(1983);
L.Mandel, "Sub-Poissonian Photon Statistics and Squeezed Quantum
States", invited paper XIII International Quantum Electronics
Conference,Anaheim,Ca,June 1984.

TABLE 1.

c	speed of light
m	electron mass
$\hbar$ ($=\hbar/2\pi$)	Planck constant
P, Z	electron longitudinal linear momentum and coordinate
$a_{L,U}, a_{L,U}^\dagger$	annihilation and creation operators of laser (L) and undulator (U) photons
$\vec{k}_L = -\vec{k}_U$	laser and undulator wavevectors in the moving frame
$\omega = \vec{k} c$	laser frequency
$n_{L,U}$	number of photons, laser (L) and undulator (U)
L/c	Interaction time
$r_0 = e^2/mc^2$	classical electron radius
V	interaction volume
$\Omega = \frac{2\pi c^2}{\omega} r_0/V$	coupling constant
$2\omega P_0 L/mc^2 = W_0$	resonance parameter
$2\hbar k^2 L/mc = \epsilon$	electron recoil parameter
$\Omega_R =$	$L/c \Omega \sqrt{m_0^2}$

FIGURE CAPTIONS.

Fig.1. Deviation from a Poisson distribution. $\{ \langle \Delta(n+l)^2 \rangle - \langle n+l \rangle \} / \epsilon$
vs. resonance parameter W_0 for two different values of the
initial number of photons a) $|\alpha_0|^2 = 0$; b) $|\alpha_0|^2 = 10^{14}$. The coupling
constant has been set $\Omega_R = 0.25$ (small signal regime)



