

REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS
BEFORE COMPLETING FORM

1. REPORT NUMBER
No. 34

2. GOVT ACCESSION NO.

3. RECIPIENT'S CATALOG NUMBER

TITLE (and Subtitle)

The Spherical Raman-Nath Equation
with Time-Dependent Coefficients

5. TYPE OF REPORT & PERIOD COVERED
Technical Report
10/01/84 - 09/30/85

6. PERFORMING ORG. REPORT NUMBER

AUTHOR(s)

G. Dattoli and J. Gallardo

8. CONTRACT OR GRANT NUMBER(s)

N00014-80-C-0308

PERFORMING ORGANIZATION NAME AND ADDRESS

University of California
Quantum Institute
Santa Barbara, CA 93106

10. PROGRAM ELEMENT, PROJECT, TASK
AREA & WORK UNIT NUMBERS

61153N; RR011-07-06;
NR603-001

CONTROLLING OFFICE NAME AND ADDRESS

ONR
1030 E. Green St.
Pasadena, CA 91106

12. REPORT DATE

1985

13. NUMBER OF PAGES

14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)

15. SECURITY CLASS. (of this report)
Unclassified/Unlimited

15a. DECLASSIFICATION/DOWNGRADING
SCHEDULE

16. DISTRIBUTION STATEMENT (of this Report)

Approved for public release; distribution.

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)

ADVISORY GROUP
RECEIVED
FEB 10 1986
ON ELECTRON DEVICES

18. SUPPLEMENTARY NOTES

Lettere al Nuovo Cimento, 42, 163, 1985.

19. KEY WORDS (Continue on reverse side if necessary and identify by block number)

Raman-Nath, Time-Dependent, Coefficients

DTIC
ELECTE
MAY 21 1986
S D

20. ABSTRACT (Continue on reverse side if necessary and identify by block number)

In this letter we write the exact solution of the spherical Raman-Nath equation with time dependent coefficients in terms of a hypergeometric function. Finally we indicate how already known results can be derived from this more general expression.

AD-A221 741

ADVISORY COPY

THE SPHERICAL RAMAN-NATH EQUATION WITH
TIME-DEPENDENT COEFFICIENTS

G. Dattoli and J. Gallardo

Lettere al Nuovo Cimento
1985

QIFEL034/84



Accession For	
NTIS	RAAI <input checked="" type="checkbox"/>
OTIC	FAB <input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution /	
Availability Codes	
Dist	▲ _____
A-1	

THE SPHERICAL RAMAN-NATH EQUATION WITH TIME-DEPENDENT
COEFFICIENTS

G. Dattoli,+ J. Gallardo,* and A. Torre#

+ E.N.E.A. Centro Ricerche Energia, Frascati C.P. 65-00044,
Frascati, Italy.

* Quantum Institute, University of California at Santa
Barbara, California 93106.

INFN Sezione di Napoli, Italy.

ABSTRACT.

In this letter we write the exact solution of the Spherical Raman-Nath equation with time dependent coefficients in terms of a hypergeometric function. Finally we indicate how already known results can be derived from this more general expression.

The Raman-Nath (R.N.) type equations are a set of differential recursive equations (1), which have become familiar to the physics community in recent times. They are indeed relevant to a large number of physical problems; to quote a few of them we recall light diffraction by ultrasounds (2), atomic and molecular dynamics (3), and Free Electron Lasers (4). (For a more extended list, the reader is referred to Ref. 1.)

A general and systematic analysis of these equations has been put forward in Ref. (1,5). In these papers an operational technique, helpful to look for both exact and non trivial perturbed solutions, has been developed.

Together with the R.N. equations discussed in Refs. (1,5), a more general class of recursive differential equations has been considered(6). These equations, which appear in the analysis of stimulated Compton-Scattering (4) or may be exploited to write the time evolution of the so called coherent Block states (7-9), has been named spherical or SU2-R.N. equations .

In this note we will find a solution for the most general form of SU2-R.N. equation, solvable in terms of known functions, and we will indicate how all the previously found solutions (1,5,6) are only particular cases of this more general expression.

The equation we consider is the following :

$$i \frac{d}{d\tau} C_{\ell}(\tau) = \omega(\tau) \left(\frac{n_+ - n_-}{2} + \ell \right) C_{\ell}(\tau) + \Omega(\tau) \sqrt{(n_- - \ell)(n_+ + \ell + 1)} C_{\ell+1}(\tau) \\ + \Omega^*(\tau) \sqrt{(n_+ + \ell)(n_- - \ell + 1)} C_{\ell-1}(\tau) \\ C_{\ell}(0) = \delta_{\ell,0} \quad (1)$$

where n_{\pm} are fixed integers, $\omega(t)$ and $\Omega(t)$ are time-dependent integrable functions, and ℓ is a discrete variable ($-n_- \leq \ell \leq n_+$).

We impose the initial condition $C_{\ell}(0) = \delta_{\ell,0}$. This choice is not restrictive and indeed for general initial condition of the form $C_{\ell}(0) = f_{\ell}$ the solution of Eq.(1) can be obtained following the procedure below with only very few changes (5).

The technique, adopted to find the solution of the above equation, is a generalization of that presented in Ref. (5). We rewrite Eq. (1) in a more compact form by defining

$$C_{\ell}(\tau) = (-i)^{\ell} \exp \left[-i \left(\frac{n_+ - n_-}{2} + \ell \right) \int_0^{\tau} d\tau' \omega(\tau') \right] M_{\ell}(\tau) \quad (2)$$

Introducing angular momentum-type operators, we easily find

$$\frac{d}{d\tau} M_\ell(\tau) = -\Omega(\tau) \exp\left[-i \int_0^\tau d\tau' \omega(\tau')\right] \hat{J}_+ M_\ell(\tau) + \Omega^*(\tau) \exp\left[i \int_0^\tau d\tau' \omega(\tau')\right] \hat{J}_- M_\ell(\tau) \quad (3)$$

The J operators "act" on M_ℓ according to the rules

$$\hat{J}_3 M_\ell(\tau) = \left(\frac{n_+ - n_-}{2} + \ell\right) M_\ell(\tau) \quad (4)$$

$$\hat{J}_\pm M_\ell(\tau) = \sqrt{(m_\mp \mp \ell)(m_\pm \pm \ell + 1)} M_{\ell \pm 1}(\tau)$$

They obey the well-known rules of commutation

$$[\hat{J}_+, \hat{J}_-] = 2 \hat{J}_3, \quad [\hat{J}_3, \hat{J}_\pm] = \pm \hat{J}_\pm \quad (5)$$

The use of the above operators clarifies the term "SU2- R.N. equation" to indicate those equations which can be cast in the general form of Eq. (1).

The key-point in our derivation will be the use of the so called Wie-Norman Lie-algebraic analysis (10) of linear differential equations. We can, therefore, bypass all the problems relevant to the time ordering and coupling exponentials to write

$$M_\ell(\tau) = e^{2h(\tau) \hat{J}_3} e^{-g(\tau) \hat{J}_+} e^{f(\tau) \hat{J}_-} i \delta_{\ell,0} \quad (6)$$

where the functions $h(t)$, $g(t)$, and $f(t)$ obey the following system of differential equations

$$\begin{aligned} \frac{d}{d\tau} h(\tau) &= g(\tau) \frac{d}{d\tau} f(\tau) \\ \frac{d}{d\tau} g(\tau) &= \Omega(\tau) \exp\left[-2h(\tau) - i \int_0^\tau d\tau' \omega(\tau')\right] - g(\tau) \frac{d}{d\tau} h(\tau) \\ \frac{d}{d\tau} f(\tau) &= \Omega^*(\tau) \exp\left[2h(\tau) + i \int_0^\tau d\tau' \omega(\tau')\right] \end{aligned} \quad (7)$$

with initial conditions, $h(0)=g(0)=f(0)=0$. This system can be solved if the solution of the following Riccati equation is known

$$\begin{aligned} \frac{d}{d\tau} u(\tau) - u^2(\tau) + p(\tau)u(\tau) + q(\tau) &= 0 \\ u(0) &= 0 \end{aligned} \quad (8)$$

where we have set

$$\begin{aligned} u(\tau) &= \frac{d}{d\tau} h(\tau) \\ p(\tau) &= -i\omega(\tau) - \left[\frac{1}{\Omega(\tau)} \frac{d}{d\tau} \Omega(\tau) \right]^* \\ q(\tau) &= -|\Omega(\tau)|^2 \end{aligned} \quad (9)$$

However, even if we do not know the explicit solution of the Eq.(8), we can write down the expression of $C_\rho(\tau)$ in terms of the functions $f(\tau)$, $g(\tau)$, $h(\tau)$. This can be accomplished in a rather direct way. We use a series expansion

of the exponentials and the operational rules Eq.(4), so that, after a lengthy algebra we find

$$C_l(\tau) = \sqrt{\binom{n_-}{l} \binom{n_+ + l}{l}} \exp\left[\frac{n_- - n_+}{2} \mathcal{C}(\tau)\right] \\ \times \left(\mathcal{F}(\tau)\right)^l {}_2F_1(-n_+, n_+; l+1; \mathcal{F}(\tau) G(\tau)) \quad (10)$$

where to simplify the notation we have defined

$$\mathcal{C}(\tau) = i \int_0^\tau d\tau' \omega(\tau') - 2 h(\tau) \\ \mathcal{F}(\tau) = -i f(\tau) \exp\left[-i \int_0^\tau d\tau' \omega(\tau')\right] \\ \mathcal{G}(\tau) = i g(\tau) \exp\left[i \int_0^\tau d\tau' \omega(\tau')\right] \quad (11)$$

also ${}_2F_1(a, b; c; z)$ is the hypergeometric function (11).

The result in Eq.(10) is a very general one and it allows us to calculate the probability amplitude for the stimulated Compton scattering (4). Furthermore, it may be exploited to recover all previously known solutions (1,5,6).

If we take $\Omega(\tau)$ and $\omega(\tau)$ as time independent functions and $n_+ = 0$, it is trivial to recover the solution written in Ref. (6) in terms of the binomial generalization of the Poisson-

Charlier function(12).

The connection with the cases discussed in Refs. (1,5) requires the use of the asymptotic properties of ${}_2F_1(a,b;c;z)$; taking the large n - limit the hypergeometric function reduces to a Laguerre function (11) so that Eq.(10) yields the solution of the Harmonic R.N. equation . Finally taking both η_+ and η_- very large , ${}_2F_1(a,b;c;z)$ becomes a Bessel function (11) and thus we recover the solution of the "shifted" R.N. equation .

In summary, we have found an exact solution of the SU2-R.N. equation. In a forthcoming paper we will apply this solution to the analysis of the time evolution of the coherent Block-states(9). In that paper we will discuss, in more details, the problem of the asymptotic limits touched upon in the last part of this letter.

Acknowledgement

The authors express their gratitude to A. Renieri, S. Solimeno and M. Richetta for enlightening discussions.

One of the authors (JG) has been supported by the Office of Naval Research, ONR contract N00014-80-C-0308.

REFERENCES.

1. G. Dattoli, M. Richetta and I. Pinto, E.N.E.A. Report 83.39/p Frascati, submitted for publication.
2. M.V. Berry, "The Diffraction of Light by Ultrasound", Academic Press, London-New York (1966).
3. J. H. Eberly, B. W. Shore, Z. Bialyuincka-Birula and I. Bialyincka-Birula, Phys. Rev A16, 2048 (1977).
4. G. Dattoli and A. Renieri, "Experimental and Theoretical aspects of the Free Electron Laser" to appear in the IV th volume of Laser Handbook, M.L.Stitch and M.S.Bass, eds., North-Holland.
5. P.Bosco and G.Dattoli, J.Phys.A16,4409(1983); F.Ciocci, G.Dattoli and M.Richetta, J.Phys.A17, 1393(1984).
6. P.Bosco, G.Dattoli and M.Richetta, J.Phys.A17,L395(1984); P.Bosco, J.Gallardo and G.Dattoli, J.Phys.A17,2739(1984).
7. F.T. Arecchi, E. Courtens, G. Gilmore and H. Thomas Phys.Rev.A6,2211(1972); J.M. Radcliffe, J.Phys.A4,313(1970).
8. R.Bonifacio, D.M.Kim and M.O.Scully, Phys.Rev.187,441(1969).
9. G.Dattoli, J.Gallardo and A.Torre, in preparation
10. J.Wei and E.Norman, J.of Math.Phys.4A,575(1963).
11. A. Erdely et al., "Higher Trascendental Functions", vol.1, Chap.VI, McGraw-Hill.

12. G.Szëgo, "Orthogonal Polynomials", Amer.Math.Soc.Colloquium
Pub. vol 23, New York (1939).