

AD-A223 114

FTD-ID(RS)T-0792-88

FOREIGN TECHNOLOGY DIVISION



SPECIAL FEATURES OF THE STRONG INTERACTION OF A PLASMA WITH A MAGNETIC
FIELD AS IT EXPANDS RADIALLY IN A DISK CHANNEL

by

A. Yu. Kervis, V.S. Sokolov, et al.



DTIC
ELECTE
JUN 14 1990
S E D

Approved for public release;
Distribution unlimited.



MICROFORM CLASSIFICATION NOTICE/TARGET (For Unclassified Material)

(TO BE COMPLETED BY RECORDS CUSTODIAN OR CAMERA OPERATOR/DOCUMENT PREPARER IN GOVERNMENT OR CONTRACT MICROFORM SERVICE CENTER)


**CLASSIFICATION NOTICE
THE HIGHEST CONTAINED
ON THIS MICROFORM IS:**

UNCLASSIFIED

IF CHECKED THE FOLLOWING ACCESS WARNING NOTICES APPLY

☐	FOR OFFICIAL USE ONLY (AFR 12-30)
☐	PRIVACY ACT OF 1974 (AFR 12-35)
☐	SOURCE SELECTION SENSITIVE (AFR 70-15)
☐	OTHER (Specify) (For Unclassified Material)

DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

HUMAN TRANSLATION

FTD-ID(RS)T-0792-88

25 October 1988

MICROFICHE NR: FTD-88-C-000888

**SPECIAL FEATURES OF THE STRONG INTERACTION OF
A PLASMA WITH A MAGNETIC FIELD AS IT EXPANDS
RADIALLY IN A DISK CHANNEL**

By: A. Yu. Kerkis, V.S. Sokolov, et al.

English pages: 6

**Source: Novosibirsk Aerofizicheskiye Issledovaniya,
pp. 71-75**

Country of origin: USSR

Translated by: Charles T. Ostertag, Jr.

Requester: FTD/TQTD

Approved for public release; Distribution unlimited.

THIS TRANSLATION IS A RENDITION OF THE ORIGINAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DIVISION.

PREPARED BY:

TRANSLATION DIVISION
FOREIGN TECHNOLOGY DIVISION
WPAFB, OHIO.

U. S. BOARD ON GEOGRAPHIC NAMES transliteration SYSTEM

Block	Italic	Transliteration	Block	Italic	Transliteration
А а	<i>А а</i>	A, a	Р р	<i>Р р</i>	R, r
Б б	<i>Б б</i>	B, b	С с	<i>С с</i>	S, s
В в	<i>В в</i>	V, v	Т т	<i>Т т</i>	T, t
Г г	<i>Г г</i>	G, g	У у	<i>У у</i>	U, u
Д д	<i>Д д</i>	D, d	Ф ф	<i>Ф ф</i>	F, f
Е е	<i>Е е</i>	Ye, ye; E, e ^u	Х х	<i>Х х</i>	Kh, kh
Ж ж	<i>Ж ж</i>	Zh, zh	Ц ц	<i>Ц ц</i>	Ts, ts
З з	<i>З з</i>	Z, z	Ч ч	<i>Ч ч</i>	Ch, ch
И и	<i>И и</i>	I, i	Ш ш	<i>Ш ш</i>	Sh, sh
Й й	<i>Й й</i>	Y, y	Щ щ	<i>Щ щ</i>	Shch, shch
К к	<i>К к</i>	K, k	Ъ ъ	<i>Ъ ъ</i>	"
Л л	<i>Л л</i>	L, l	Ы ы	<i>Ы ы</i>	Y, y
М м	<i>М м</i>	M, m	Ь ь	<i>Ь ь</i>	'
Н н	<i>Н н</i>	N, n	Э э	<i>Э э</i>	E, e
О о	<i>О о</i>	O, o	Ю ю	<i>Ю ю</i>	Yu, yu
П п	<i>П п</i>	P, p	Я я	<i>Я я</i>	Ya, ya

^uye initially, after vowels, and after ъ, ь; e elsewhere.
When written as ѣ in Russian, transliterate as y^u or ѣ.

RUSSIAN AND ENGLISH TRIGONOMETRIC FUNCTIONS

Russian	English	Russian	English	Russian	English
sin	sin	sh	sinh	arc sh	sinh ⁻¹
cos	cos	ch	cosh	arc ch	cosh ⁻¹
tg	tan	th	tanh	arc th	tanh ⁻¹
ctg	cot	cth	coth	arc cth	coth ⁻¹
sec	sec	sch	sech	arc sch	sech ⁻¹
cosec	csc	cach	csch	arc cach	csch ⁻¹

Russian English

rot curl
lg log

GRAPHICS DISCLAIMER

All figures, graphics, tables, equations, etc.
merged into this translation were extracted
from the best quality copy available.



Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Avail and/or	Special
A-1 23	

**SPECIAL FEATURES OF THE STRONG INTERACTION OF A
PLASMA WITH A MAGNETIC FIELD AS IT EXPANDS
RADIALLY IN A DISK CHANNEL**

Kerkis, A.Yu., Sokolov, V.S.,
Trynkina, N.A., Pomichev, V.P.,

The investigation concerns the process of expansion of a plasma in a disk channel when a transverse magnetic field is present. The layout of the experimental unit is shown in Figure 2.19. The plasma source is a coaxial electric discharger, on which a capacitor bank of the IMM-5-150 type is discharged. The initial air pressure comprises 0.7-1.0 mm Hg. The plasma which forms as a result of the electric discharge moves along a straight tube, which is a continuation of the discharger, at a velocity of up to 20 km/s. This tube, which has a length of 30 cm and diameter of 5 cm, terminates in a disk channel with a width of 2 cm. The process develops in such a manner that the plasma, following impact on the end of the straight tube, is expanded in the disk channel. The leading edge of the moving plasma, already at a radius of 60 mm, is a shock wave, traveling over a quiescent gas at a velocity of up to 10 km/s (without a magnetic field). The results of the measurement of pressure along the length of the disk channel at different moments of time when the magnetic field is absent are given in Figure 2.20. Also given are the estimates and the indirect measurements of distribution of electrical conductivity over the radius at different moments of time. These measurements in combination with

data on pressure showed that the change in the parameters of the gas beyond the cylindrical shock wave take place primarily as a result of its adiabatic expansion.

In this case the presence of a transverse magnetic field, when its strength exceeds a certain critical magnitude, leads to a qualitative change in the nature of expansion of the gas. Direct measurements of the current density in the plasma, changes in the strength of the magnetic and electric fields, pressure (Figure 2.21), and also high-speed photography with the help of an SFRL (in the mode of photosweep and time magnifier), lead to the detection of a clearly expressed current layer (Figure 2.22), appearing in the gas beyond the direct shock wave and leading, as a result of its intensive stagnation in the magnetic field, to the formation of a reflected shock wave. In this case the measured current density reaches a magnitude of 2000 A/cm^2 , which at a field strength of $\sim 4000 \text{ Oe}$ under the conditions of the experiment leads practically to the complete standstill of the gas in this zone. Estimates show that the magnitude of electrical conductivity inside the current layer reaches values of $120\text{-}150 \text{ mho/cm}$ (without a magnetic field it is equal to $5\text{-}10 \text{ mho/cm}$).




Figure 2.19.

Key: (1) Pressure sensors; (2) Current sensors; (3) Magnetic probes;
(4) SFR.




Figure 2.20.




Figure 2.21.

Key: (1) μs .




Figure 2.22. Frame-by-frame photography of the process. Exposure time for one frame $1.33 \mu\text{s}$. Sequence of frames: from top to bottom, from left to right.

During the course of the experiment attention was turned to the stability of the current layer. It was revealed that it had exceptional stability as a geometric formation. In a number of cases on the external boundary of the layer a picture developed which was characteristic for Rayleigh-Taylor instability, which, however, disappeared quite rapidly, leading to the complete restoration of the structure of the current layer. The process of expansion of the conducting gas in a transverse magnetic field, which is being investigated at the present time, and which is accompanied by the formation of a clearly expressed current layer, coincides in its nature with the analogous process which was described in [1], and thereby confirms experimentally the conclusions of this work, obtained on the basis of a numerical solution of equations of magnetic hydrodynamics.

BIBLIOGRAPHY

1. A.M. Tikhonov, A.A. Samarskiy, L.A. Zaklyaz'minskiy,
B.M. Volosevich, L.M. Degtyarev, S.P. Kurdyumov,
Yu.P. Popov, V.S. Sokolov, A.P. Favorskiy, "Nonlinear effect of the
formation of a self-sustaining, high-temperature conducting layer of
gas in nonsteady-state processes of magnetic hydrodynamics," DAN
[Reports of the Academy of Sciences], No 4, v 173, 1967.

DISTRIBUTION LIST

DISTRIBUTION DIRECT TO RECIPIENT

<u>ORGANIZATION</u>	<u>MICROFICHE</u>
A205 DMAHTC	1
A210 DMAAC	1
CS09 BALLISTIC RES LAB	1
CS10 R&T LABS/AVEADCOM	1
CS13 AFRADCOM	1
CS35 AVRADCOM/TSARCOM	1
CS39 TRASANA	1
CS91 FSTC	4
C619 MIA REDSTONE	1
D008 MILSC	1
E053 HQ USAF/INET	1
E404 AEDC/DOF	1
E408 AFWL	1
E410 AD/IND	1
E429 SD/IND	1
P005 DOE/ISA/DDI	1
P050 CIA/OCR/ADD/SD	2
AFTT/LDE	1
FTD	
CCV	1
MIA/PHS	1
LLYL/CODE L-389	1
NASA/NST-44	1
NSA/T513/TDL	2
ASD/FTD/TOLA	1
FSL/NIX-3	1

MICROFORM CERTIFICATE OF AUTHENTICITY

START

THIS IS TO CERTIFY THAT THE MICROGRAPHIC IMAGES APPEARING ON THIS MICROFORM ARE ACCURATE REPRODUCTIONS OF THE RECORDS CREATED OR MAINTAINED BY: THE FOREIGN TECHNOLOGY DIVISION, WRIGHT-PATTERSON AFB, OHIO

PREVIOUS DESIGNATIONS

AND WERE MICROFILMED IN THE COURSE OF BUSINESS PURSUANT TO ESTABLISHED ROUTINE POLICY FOR SYSTEMS UTILIZATION AND/OR FOR THE MAINTENANCE AND PRESERVATION OF SUCH RECORDS THROUGH THE STORAGE OF SUCH MICROFORMS IN PROTECTED LOCATIONS.

IT IS FURTHER CERTIFIED THAT THE PHOTOGRAPHIC PROCESSES USED FOR MICROFILMING OF THE ABOVE RECORDS WERE ACCOMPLISHED IN A MANNER AND ON MICROFILM WHICH MEETS THE RECOMMENDED REQUIREMENTS OF THE NATIONAL BUREAU OF STANDARDS FOR MICROGRAPHIC REPRODUCTIONS.

DATE MICROFILMED <i>21 Nov 88</i>	CAMERA OPERATOR <i>KH</i>
SERIAL NUMBER <i>(3486) (2609) (1062)</i>	RECORDS MANAGEMENT DIVISION
FILE LOCATION FTD/DARM, WPAFB, OHIO	FTD/DA 24x
AUTHORITY AFR 12-40 AND MICROFORM SYSTEM NUMBER AFSC 1B-72	REDUCTION RATIO