

(U)

# DTIC FILE COPY

2

SECURITY

## REPORT PAGE

Form Approved  
OMB No 0704-0188

1a REPORT NUMBER

# AD-A223 360

1b. RESTRICTIVE MARKINGS

NA

2a SECURITY CLASSIFICATION

3 DISTRIBUTION/AVAILABILITY OF REPORT

Distribution unlimited

2b DECLASSIFICATION/DOWNGRADING SCHEDULE

NA

4 PERFORMING ORGANIZATION REPORT NUMBER(S)

University of Minnesota  
College of Biological Sciences

5 MONITORING ORGANIZATION REPORT NUMBER(S)

NA

6a NAME OF PERFORMING ORGANIZATION

University of Minnesota

6b. OFFICE SYMBOL  
(If applicable)

NA

7a. NAME OF MONITORING ORGANIZATION

Office of Naval Research

6c. ADDRESS (City, State, and ZIP Code)  
Department of Biochemistry  
1479 Gortner Ave.  
St. Paul, MN 55108

7b. ADDRESS (City, State, and ZIP Code)

800 N. Quincy St.  
Arlington, VA 22217-5000

8a NAME OF FUNDING/SPONSORING ORGANIZATION

Office of Naval Research

8b. OFFICE SYMBOL  
(If applicable)

ONR

9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

N00014-88-J-1220

8c. ADDRESS (City, State, and ZIP Code)

800 N. Quincy St.  
Arlington, VA 22217-5000

10 SOURCE OF FUNDING NUMBERS

PROGRAM ELEMENT NO	PROJECT NO.	TASK NO	WORK UNIT ACCESSION NO
61153N	RR04108		

11 TITLE (Include Security Classification)

(U) Effects of Electric Fields on Membrane-Bound Na,K-ATPase

12. PERSONAL AUTHOR(S)

Tsong, Tian Y.

13a. TYPE OF REPORT

Progress

13b TIME COVERED

FROM 7/1/89 TO 6/30/90

14. DATE OF REPORT (Year, Month, Day)

6/30/90

15 PAGE COUNT

8

16 SUPPLEMENTARY NOTATION

17 COSATI CODES

FIELD	GROUP	SUB-GROUP
08		

18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

Electric Fields, Cell Membranes, Enzymes,  
Biological Effects, *JES*

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

We continued to work on effects of oscillating electric fields on membrane functions, in particular the electric activation of Na,K-ATPase, and to develop theory of electroconformational coupling. We believe transmembrane electric fields are involved in the regulation of the internal activity of a cell and also in the cell-to-cell communications. An in depth study of Na,K-ATPase will provide useful information concerning the molecular design of a cell to sense and to transmit signals.

### DISTRIBUTION STATEMENT A

Approved for public release;  
Distribution Unlimited

20 DISTRIBUTION/AVAILABILITY OF ABSTRACT

UNCLASSIFIED/UNLIMITED  SAME AS RPT  DTIC USERS

21 ABSTRACT SECURITY CLASSIFICATION

(U)

22a NAME OF RESPONSIBLE INDIVIDUAL

Dr. Igor Vodnyanov

22b TELEPHONE (Include Area Code)

(202) 696-4056

22c OFFICE SYMBOL

ONR

## Progress Reports on ONR Grant N00014-88-J-1220



PRINCIPAL INVESTIGATOR: Tian Y. Tsong

CONTRACTOR: University of Minnesota

CONTRACT TITLE: Effect of Electric Fields on Membrane Bound Na,K-ATPase

COVERED PERIOD: 1/7/89 - 30/6/90

## PROGRESS:

1). In the a.c. activation of Na,K-ATPase, we began to obtain data for kinetic analysis and for comparing results in which ATP is the primary energy source. This comparison will examine whether the a.c. induced ion pumping activity is by a similar mechanism as the in vivo mechanism of ion pumpings. An a.c. field can also induce transfection of E. coli by plasmid DNA. This activity showed an optimal frequency at 1 Hz.

2). In the development of the electroconformational coupling model, we have obtained conditions in which the efficiency of the transduction of electric energy can approach that of the theoretical maximum, 100%. Other oscillating thermodynamic potentials have also been found to display similar properties. In particular, a pressure sensitive ion transporter has been made to absorb acoustic energy to pump an ion against its concentration gradient. The conversion of acoustic or mechanical signals into electric signals is an important ability of cells. This model will be called, piezo-electric coupling model and will be investigated in details.

3). Transmembrane potential induced by an oscillating electric field: The  $\Delta\psi_{\text{membr}}$  generated by an a.c. field depends strongly on the frequency of the field and can be calculated using the Schwan Equation.

$$\Delta\psi_{\text{membr}} = 1.5 a E_{\text{appl}} \cos \theta / [1 + (\omega \tau)^2]^{1/2} \quad (1)$$

$$\text{where } \tau = a C_{\text{membr}} (\rho_{\text{int}} + \rho_{\text{ext}}/2) \quad (2)$$

In the equations,  $\omega = 2\pi f$ , and  $f$  is the frequency of the electric field.  $\tau$ ,  $a$ ,  $\theta$ ,  $C_{\text{membr}}$ ,  $\rho_{\text{int}}$ , and  $\rho_{\text{ext}}$  are, respectively, membrane relaxation time, the radius of the cell (spherical), the angle between the field line and the normal to a position of interest, membrane capacitance, the resistivity of the cytoplasmic fluid, and of the external medium. We have measured the critical breakdown potential,  $\Delta\psi_{\text{crit}}$ , of the plasma membrane of murine myeloma cell line (Tib9) using a.c. field, by monitoring the entry of a fluorescence probe, propidium iodide, into the cells. The set up for these measurements is shown in Figure 1. This dye is weakly fluorescent in solution but becomes strongly fluorescent when it binds to DNA. Experiments were done under a

Accession For	
NTIS	CRA&I <input checked="" type="checkbox"/>
DTIC	TAB <input type="checkbox"/>
Unannounced <input type="checkbox"/>	
Justification _____	
By _____	
Distribution / _____	
Availability Codes	
Dist	Avail and/or Special
A-1	

microscope by direct visual examination of single cells or by examining photographic prints. When an a.c. field reached the intensity,  $E_{crit}$ , that generated a maximal membrane potential,  $\Delta\psi_{max}$ , equal to or greater than the  $\Delta\psi_{crit}$ , the membrane was perforated at the two loci facing the electrodes. The dye diffused into the cell, giving rise to two bright, narrow, moon-like bands, which expanded to the whole cell in 1-3 minutes (Figure 2).  $\Delta\psi_{crit}$ 's were measured in three media of different resistivities,  $\rho_{ext}$ , (52600, 7050 and 2380  $\Omega$  cm), over the range 0.1 kHz to 300 kHz, with the field duration of 200 ms. Regression analysis based on equation (3) showed that in a medium of given resistivity, the  $\Delta\psi_{crit}$  was constant over the frequency range studied. When the capacitance of the membrane,  $C_{membr}$ , was taken to be  $0.9 \mu F cm^{-2}$ , the resistivity of the cytoplasmic medium was determined to be 910-1100  $\Omega$  cm.

$$\Delta\psi_{crit} = 1.5 a E_{crit} / [1 + (\omega \tau)^2]^{1/2} \quad (3)$$

The  $\Delta\psi_{crit}$  were 0.33 V, 0.48 V and 0.53 V, respectively, for the three media in decreasing resistivities. The good fit of these data to the curves calculated using the Schwan Equation indicates that the equation may be used to describe the transmembrane potential of a living cell generated by an oscillating electric field (Figure 3). Some electric parameters are summarized in Table 1.

## Publications

1. Tsong, T.Y. (1989). Electroconformational coupling: A fundamental process of biomolecular electronics for signal transductions. In "Molecular Electronics: Biosensors and Biocomputers", Hong, F., Ed., Plenum Publ. Corp. Pp. 83-95.
2. Tsong, T.Y., Astumian, R.D. (1989). Charge-field interactions in cell membranes and electroconformational coupling: Transduction of electric energy by membrane ATPase. In "Charge-Field Effects in Biosystems II", Allen, M.J., Ed., Plenum Publ. Corp. Pp. 167-177.
3. Astumian, R.D., Robertson, B., Tsong, T.Y. (1989). Charge-field interactions in cell membranes and electroconformational coupling: Theory for the interactions between dynamic electric fields and membrane enzymes. In "Charge-Field Effects in Biosystems II", Allen, M.J., Ed., Plenum Publ. Corp. Pp. 179-190.
4. Robertson, B., Astumian, R.D., Tsong, T.Y. (1989). Nonlinear effects of periodic electric fields on membrane proteins. In "Charge-Field Effects in Biosystems II", Allen, M.J., Ed., Plenum Publ. Corp. Pp. 191-209.
5. Liu, D.-S., Astumian, R.D., Tsong, T.Y. (1990). Activation of the Na<sup>+</sup> and the Rb<sup>+</sup> modes of Na,K-ATPase by an oscillating electric field. *J. Biol. Chem.* 265:7260-7267.
6. Tsong, T.Y. (1990). Electric modulation of membrane proteins: Enforced conformational oscillation and cellular energy and signal transductions. *Annu. Rev. Biophys. Biophys. Chem.* 19:83-106.
7. DeBustros, A., Lee, R.Y., Compton, D., Tsong, T.Y., Baylin, S.B., Nelkin, B.D. (1990). Different cis-acting DNA elements regulate expression of the human calcitonin gene in cultured lines of lung carcinoma and medullary thyroid carcinomas. *Mole. Cell Biol.* 10:1773-1778.
8. Xie, T.-D., Sun, L., Tsong, T.Y. (1990). Study of mechanisms of electric field induced DNA transfection I. DNA entry by surface binding and diffusion through membrane pores. *Biophys. J.* In press.
9. Xie, T.-D., Tsong, T.Y. (1990). Study of mechanisms of electric field induced DNA transfection II. Transfection by low amplitude, low frequency alternating electric fields. *Biophys. J.* In press.
10. Markin, V.S., Tsong, T.Y., Astumian, R.D., Robertson, B. (1990). Energy transduction between a concentration gradient and an alternating electric field. *J. Chem. Phys.* In press.
11. Marszalek, P., Liu, T.-S., Tsong, T.Y. (1990). Schwan Equation and transmembrane potential induced by alternating electric field. *Biophys. J.* In press.
12. Tsong, T.Y., Tomita, M. (1990). Selective B-lymphocyte-myeloma cell fusion. *Methods in Enzymol.* In press.
13. Tsong, T.Y. (1990). On electroporation and some related cell membrane phenomena. *Bioelectrochem. Bioenerg.* In press.

And several abstracts in scientific meetings, several manuscripts submitted.

**Table 1 Some electric parameters of myeloma cells**

Parameter	Experiment 1	Experiment 2	Experiment 3
$a$ ( $\mu\text{m}$ )	6.5	6.5	6.5
$\rho_{\text{ext}}$ ( $\Omega\text{ cm}$ )	52600	7050	2380
$\rho_{\text{int}}$ ( $\Omega\text{ cm}$ )	n.a.	910	1100
$\alpha$	0.17	0.030	0.015
$\tau'$ ( $\mu\text{s}$ )	13 <sup>a</sup>	2.5 <sup>b</sup>	1.3 <sup>c</sup>
$\Delta\psi_{\text{crit}}$ (V)	0.33	0.48	0.53
$\Delta\psi_{\text{rms}}$ (V)	0.23	0.34	0.37

Notes: n.a. means not applicable.


a: Value obtained by calculation.

b: Values obtained by optimization.

$$\Delta\psi_{\text{rms}} = \Delta\psi_{\text{crit}} / \sqrt{2}.$$


$$\alpha = a G_{\text{membr}} \rho_{\text{ext}} / 2$$

$$\tau = \tau' (1 + \alpha)$$



**Fig. 1 Block diagrams of equipment and cell chamber for electroporation.**

The upper figure is the block diagram of the equipment used for electroporation: PG denotes the pulse generator (Wavetek, Model 801); AC, the functional generator (Wavetek, Model 148A); R, a  $47\ \Omega$  non-inductive resistor; and S, a switch. The lower figure illustrates the cell chamber. An a.c. field of appropriate amplitude and frequency was generated by the functional generator and applied to the cell suspension. The duration of the a.c. field, 200 ms, was controlled by a square wave triggering pulse. The distance between the two platinum electrodes was 0.1 mm. The dashed lines in the glass slide indicate the epoxy resin ring which formed a vessel to hold  $150\ \mu\text{l}$  of cell suspension. The conductivity of the cell suspension was monitored during the experiment by measuring the voltage drop on the non-inductive  $47\ \Omega$  resistor.



**Fig. 2 Observation of electroporation by the fluorescence changes of propidium iodide**

The chemical formula of propidium iodide is given in the upper figure. The middle figure shows a typical myeloma cell under the microscope (bright field). The relative positions of the platinum electrodes are indicated. The lower figure gives some photographs taken at different time after a cell was electroporated by an a.c. field,  $E_{crit}$ . Within 1-3 s, two narrow, moon-like bright bands appeared at the two loci facing the electrodes (the left most photo). The next 3 photos, from left to right, were taken at approximately 20 s, 1 min and 3 min, respectively. See text for details.




Fig. 3 Critical a.c. field strength,  $E_{crit}$  for electroporation as a function of the a.c. frequency

Data obtained in three media of different resistivities,  $52600 \Omega \text{ cm}$  (○),  $7050 \Omega \text{ cm}$  (●), and  $2380 \Omega \text{ cm}$  (□), are shown. The curves drawn through these data points were obtained by calculation or the optimization according to Eq. (6). See text for details.