

2

AD \_\_\_\_\_

IMMUNOLOGICAL APPROACH TO THE IDENTIFICATION AND DEVELOPMENT  
OF VACCINES TO VARIOUS TOXINS

ANNUAL REPORT

Tran C. Chanh, Ph.D.  
Deborah Armstrong, Ph.D.  
Ronald C. Kennedy, Ph.D.

June 18, 1990

Supported by

U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND  
Fort Detrick, Frederick, Maryland 21702-5012

Contract No. DAMD17-87-C-7005

SOUTHWEST FOUNDATION FOR BIOMEDICAL RESEARCH  
San Antonio, Texas 78228

DTIC  
ELECTE  
AUG 01 1990  
S E D

DOD DISTRIBUTION STATEMENT

Approved for public release; distribution unlimited. The findings in this report are not to be construed as an official Department of Army position unless so designated by other authorized documents.

AD-A225 047

90 08 01 0210

**REPORT DOCUMENTATION PAGE**

Form Approved  
OMB No 0704-0188

1a REPORT SECURITY CLASSIFICATION Unclassified		1b RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3 DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited	
2b DECLASSIFICATION / DOWNGRADING SCHEDULE		4 PERFORMING ORGANIZATION REPORT NUMBER(S)	
4 PERFORMING ORGANIZATION REPORT NUMBER(S)		5 MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION Southwest Foundation for Biomedical Research	6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code) P.O. Box 28147 San Antonio, Texas 78228-0147		7b. ADDRESS (City, State, and ZIP Code)	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION U.S. Army Medical Research & Development Command	8b. OFFICE SYMBOL (if applicable)	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER DAMD17-87-C-7005	
8c. ADDRESS (City, State, and ZIP Code) Fort Detrick, Frederick, MD 21702-5012		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO 61102A	PROJECT NO. 3M1- 61102Bs12
		TASK NO. AA	WORK UNIT ACCESSION NO. 126
11. TITLE (Include Security Classification) Immunological Approach to the Identification and Development of Vaccine to Various Toxins			
12. PERSONAL AUTHOR(S) Tran C. Chanh, Deborah Armstrong, and Ronald C. Kennedy			
13a. TYPE OF REPORT Annual	13b. TIME COVERED FROM 12-8-88 TO 2-6-90	14. DATE OF REPORT (Year, Month, Day) 1990 June 18	15. PAGE COUNT
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	Anti-idiotypic vaccine, in-vitro and in-vivo protection against toxicity, monoclonal anti-idiotypic antibody. RAI
06	01		
06	13		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Monoclonal antibodies specific for saxitoxin and tetrodotoxin with higher binding affinities than those generated under this contract are still being sought for. Attempts to generate internal image monoclonal anti-idiotypic antibodies specific for the highest binding anti-saxitoxin and anti-tetrodotoxin antibodies are continuing. A monoclonal anti-idiotypic antibody, designated DE8, specific for a protective anti-mycotoxin T-2 monoclonal antibody (HD11) was generated. Administration of DE8 into syngeneic BALB/c mice induced a protective antibody response against the <u>in-vitro</u> and <u>in-vivo</u> toxicity of T-2.			
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION	
22a. NAME OF RESPONSIBLE INDIVIDUAL Virginia M. Miller		22b TELEPHONE (Include Area Code) (301) 663-7325	22c OFFICE SYMBOL SGRD-RMI-S

FOREWORD

In the conducting the research described in the report, the investigators adhered to the "Guide for the Care and Use of Laboratory Animals," as prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Animal Resources - National Research Council (DHEW Publication No. (NIH) 78-23, Revised 1978).

Citations of commercial organizations and trade names in this report to do not constitute an official Department of the Army endorsement or approval of the products or services of these organizations.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
	Avail and/or
Dist	Special

A-1



TABLE OF CONTENTS

Report Documentation Page.....1

Foreword.....2

Table of Contents.....3

I. Introduction.....4

II. Body of the Report

    A. Studies with Saxitoxin.....5

    B. Studies with Tetrodotoxin.....5

    C. Studies with T-2 Mycotoxin.....6

III. Summary.....7

    Table 1.....8

    Publications.....9

## I. INTRODUCTION

The sodium channel blockers, saxitoxin (STX) and tetrodotoxin (TDT), and the protein synthesis inhibitor mycotoxin T-2 are potent, nonproteinaceous small molecular weight biological toxins. Because of their small size, they do not generally induce an antibody response with a high average binding affinity, even when conjugated to large protein carriers, which could protect against the toxicity induced by these biological toxins. Moreover, the current methodology for conjugation of toxins to protein carriers usually result in the formation of unstable complexes with the potential release of the native toxins in their toxic form. Therefore, it is impractical if not impossible to develop effective vaccines against these biological toxins using conventional technology such as synthetic peptides or recombinant technology. The very toxic nature of these biological compounds also preclude their use as immunogens in the induction of effective protection. Because of these difficulties, we have attempted to develop anti-idiotypic (anti-Id) based vaccines for the induction of protective immunity against STX, TDT and T-2. Our approach involves the generation of high affinity monoclonal antibodies (mAbs) specific for the respective toxins which are protective against toxicity. These anti-toxin antibodies (Ab1) are then used to generate syngeneic monoclonal anti-Id antibodies which serologically mimic the nominal antigen (or toxins). The generated anti-Id antibodies are then assessed for their ability to mimic the antigen and to induce in mice a protective immunity against toxicity.

Two specific anti-STX mAbs were generated that possess affinity constants ( $K_a$  values) of approximately  $10^6 \text{ M}^{-1}$ . They are designated S1A5 and S3E.2, and are of the  $\text{IgM}_k$  and  $\text{IgG}_{1k}$  isotypes, respectively. Two stable and specific clones were also generated against TDT among several thousand hybrids tested. The mAbs, termed TD13a1 and TD2C5, are  $\text{IgG}_{1k}$  mAbs with  $K_a$ 's values approaching  $10^7 \text{ M}^{-1}$ . In spite of their relatively low  $K_a$ 's values, these anti-STX and anti-TDT mAbs were found to protect against the toxicity induced by STX and TDT, respectively.

In the T-2 mycotoxin system, an  $\text{IgG}_{1k}$  anti-T-2 mAb (HD11) was isolated that had a relative affinity binding constant of approximately  $4.8 \times 10^8 \text{ M}^{-1}$ . This mAb was shown to protect against the in-vitro toxicity of T-2 mycotoxin. HD11 mAb was used to generate a monoclonal anti-Id antibody, designated DE8, which when injected into mice induced a significant protective immunity against the in-vitro and in-vivo toxicity induced by T-2.

## II. RESULTS

### A. Studies with Saxitoxin.

The results of studies involving the sodium channel blocker STX with respect to generation of specific mAbs and their ability to inhibit STX toxicity in-vitro were discussed in previous annual reports and published (1). In summary, STX was conjugated to keyhole limpet hemocyanin (KLH) and used to immunize BALB/c mice. Anti-STX antibodies were detected in serum by an enzyme-linked immunosorbent assay (ELISA) within a week or two after the first immunization (1). Spleens from immunized mice were fused with NS-1 myeloma cells and approximately 7000 resultant hybrids were screened by ELISA for reactivity to STX. Two stable hybrids were isolated, subcloned, and characterized. These hybrids, termed S1A5 and S3E.2, secreted specific anti-STX antibodies that did not recognize the closely related toxin TDT, as determined by competition ELISA. The S1A5 mAb was of the IgM<sub>k</sub> class and S3E.2 of the IgG<sub>1k</sub> subclass with affinity constants of approximately  $10^6 \text{ M}^{-1}$ . The protective ability of these antibodies was tested by a competitive displacement assay for [<sup>3</sup>H]STX binding on rat brain membranes. Purified S3E.2 strongly displaced [<sup>3</sup>H]STX binding, whereas S1A5 weakly inhibited [<sup>3</sup>H]STX binding to membranes. One nanomole of S2E.2 or S1A5 was able to bind 0.03 nmol or 0.005 nmol, respectively, of STX. The S2E.2 mAb offered partial protection against STX-induced reduction of peripheral nerve action potential in rat tibial nerve when administered in situ at concentrations 10- to 30-fold greater than STX. The S1A5 mAb, despite its ability to inhibit STX binding in vitro, was completely ineffectual in situ (1). These antibodies, particularly S3E.2, thus represent potentially useful reagents for neurobiologic research, detection of toxin contamination, and diagnosis of poisoning, and may provide protection against the toxicity of STX in vivo.

Because of the relatively low affinity constants of the generated anti-STX mAbs, two approaches were taken in an attempt to increase the affinity of binding (see second annual report). Briefly, the first approach involves the in-vitro secondary immunization of in-vivo STX-primed mouse spleen cells prior to fusion. Mice were also immunized with STX-KLH coupled via the linker N-succinimidyl 3-2 (pyridyldithio)propionate (SPDP). However, neither approach resulted in higher affinity mAbs than the ones described above. The S3E.2 anti-STX mAb has been used to immunize mice for the generation of internal image monoclonal anti-Id antibodies.

### B. Studies with Tetrodotoxin.

Monoclonal anti-TDT antibodies were generated as described above for anti-STX mAbs (2). Spleens from immunized mice were fused with NS-1 mouse myeloma cells and ~ 9,329 resultant hybrids were screened by ELISA for reactivity to TDT. Two stable hybrids, termed TD13a1 and TD2C5, secreted specific anti-TDT antibodies that recognized TDT but not the related sodium channel blocker, saxitoxin (STX), as determined by competition ELISA. Both antibodies were of the IgG<sub>1k</sub> subclass with  $K_a$ 's approaching  $10^7 \text{ M}^{-1}$ . The inhibitory ability of these antibodies was tested by a competitive displacement assay for [<sup>3</sup>H]STX on rat brain membranes. Both antibodies strongly inhibited TDT binding to membranes. A nanomole of TD2C5 was able to

- 1.8 nmol of TDT, whereas a comparable amount of TD13a1 bound half as much. Furthermore, TD2C5 was able to protect against TDT-induced reduction of peripheral nerve action potentials in rat tibial nerve when administered *in situ*. Because of the protective efficacy of TD2C5 anti-TDT mAb *in-vitro* and *in-situ*, TD2C5 mAb was conjugated to KLH and used to immunize BALB/C mice and rabbits for the production of anti-Id antibodies (see second annual report). A number of fusions was performed with spleen cells from mice immunized with TD2C5-KLH. To date, no monoclonal anti-Id antibody with internal image has been isolated. Efforts are continuing to generate anti-Id with vaccine potential.

### C. Studies with T-2 Mycotoxin.

Using T-2 conjugated to ovalbumin (OVA), generously provided by Dr. J. Hewetson (USAMRIID, Fort Detrick, Frederick, MD), to immunize BALB/C mice, a monoclonal anti-T-2 antibody (HD11) was generated that had a relatively high binding affinity constant. HD11 anti-T2 mAb was also found to be protective against the T-2-induced cytotoxicity on the human epidermoid carcinoma cell lines Hep-2 and KB (3). Rabbit anti-Id antibodies specific for HD11 were generated and characterized (see second annual report and 3). HD11 conjugated to KLH was also used to immunize syngeneic BALB/C mice for the generation of monoclonal anti-Id antibodies. One mAb specific for HD11 anti-T2 was isolated. The characteristics of this anti-Id mAb, designated DE8, have been described in details elsewhere (4 and second annual report). Briefly, DE8 anti-Id mAb is specific for HD11 anti-T-2. DE8 recognizes an Id determinant on HD11 associated with the T-2 binding site since the interactions between DE8/HD11 are inhibited by free T-2 mycotoxin. Addition of DE8 mAb to HD11 completely abrogated the protective effect of HD11 against T-2 cytotoxicity, suggesting that the binding of DE8 to HD11 prevented the latter from binding to and inactivating T-2 (4). Administration of DE8-KLH or unconjugated DE8 into BALB/C mice resulted in the induction of a specific and significant circulating anti-T-2 antibody response characterized by HD11 Id-positive antibodies. Moreover, the DE8-induced anti-T-2 antibody response was able to abolish T-2 cytotoxicity as determined by [<sup>3</sup>H]-leucine uptake of the Hep-2 and KB cells (4). Taken together, these results suggest that DE8 anti-Id mAb possesses the "internal image" of the nominal antigen (T-2) and that it can serologically mimic T-2 in the induction of an anti-T-2 immune response.

Encouraged by these results of the *in-vitro* protection, we assessed the ability of DE8-KLH to induce an anti-T-2 antibody response that would protect mice against a lethal challenge with T-2. Groups of five BALB/c mice each were immunized with four injections of DE8-KLH or control anti-Id mAb. One week after the last immunization, the mice were challenged intradermally with various doses of T-2 mycotoxin. The results are presented in Table 1. In our hands, the LD<sub>50</sub> of T-2 in mice challenged intradermally is approximately 5.0 mg/kg body weight. Mice actively immunized with DE8-KLH were protected against T-2 succumbed to T-2 toxicity, all the remaining DE8-KLH-treated mice, even the groups that received 20 mg/kg (4 times the LD<sub>50</sub>), survived. The LD<sub>50</sub> of the control antibody group was comparable to that of the untreated control group. Experiments using higher doses of T-2 will be performed to determine the LD<sub>50</sub> of DE8-KLH-immunized mice. This represents to our knowledge the first demonstration of anti-Id-induced protective immunity against a small molecular weight biological toxin.

### III. Summary

Our demonstration of the successful use of anti-Id based vaccine in inducing a protective immunity against T-2 mycotoxin provides further evidence for the ability of anti-Id antibodies to serologically mimic nominal antigens of small molecular weight and of a nonproteinaceous nature. It is hoped that our efforts in the STX and TDT systems will also yield positive results.

Table 1

Administration of DES Anti-Idiotypic Protects Mice Against T-2 Toxicity In-Vivo

Mice immunized with	T-2 Mycotoxin (ug/ml)	Survival ratio <sup>a</sup>
Control Anti-Id	5.0	3/5
	10.0	4/5
	15.0	5/5
	20.0	5/5
DES Anti-Id	5.0	0/4
	10.0	1/5
	15.0	0/5
	20.0	0/5

<sup>a</sup>Number of dead mice/total number of mice tested. T-2 mycotoxin was administered intradermally.

## Publications

### Abstracts

Huot, R.I., Castro, J.A., and Chanh, T.C. Monoclonal antibodies specifically inhibit binding of sodium channel blockers to rat brain membranes, Fed. Am. Soc. Exp. Biol. 2(4):A678, 1988.

Chanh, T.C., Reed, R.C., Frenzel, G., and Huot, R.I. Anti-idiotypic antibodies against monoclonal antibody specific for the trichothecene mycotoxin T-2. Fed., Am. Soc. Exp. Biol. 2(4):A677, 1988.

Huot, R.I., J.A., Castro, D.L. Armstrong and T.C. Chanh. Polyclonal rabbit anti-idiotypic antibodies against a protective monoclonal antibody specific for the sodium channel blocker Tetrodotoxin. Sixth An. Texas Immunol. Conf. Galveston, TX, 1988.

Schick, M., Hewetson, J., Nabers, P., and Chanh, T.C. Monoclonal anti-idiotypic antibody as vaccine against mycotoxin (T-2)-induced cytotoxicity. Fed. Am. Soc. Exp. Biol. 3:A1135, 1989.

### Manuscripts

1. Huot, R.I., Armstrong, D.L., and Chanh, T.C. In vitro and in situ inhibition of the sodium channel blocker saxitoxin by monoclonal antibodies. J. Toxicol. Environ. Health. 27:281-393, 1989.

2. Huot, R.I., Armstrong, D.L., Hewetson, J.H., and Chanh, T.C. Monoclonal antibodies inhibit the sodium channel blocker tetrodotoxin in vitro and in situ, J. Clin. Invest. 83:1821-1826, 1989.

3. Chanh, T.C., Huot, R.I., Schick, M.R., and Hewetson, J.H. Anti-idiotypic antibodies against a monoclonal antibody specific for the trichothecene mycotoxin T-2. Toxicol. Appl. Pharmacol., 100:201-207, 1989.

4. Chanh, T.C., G. Rappocciolo, and J.F. Hewetson. Monoclonal Anti-Idiotypic induces a protective antibody response against the trichothecene mycotoxin T-2. In press J. Immunol., June 1990.

5. Chanh, T.C., and J.F. Hewetson. Fine specificity of a monoclonal anti-T-2 mycotoxin antibody. Submitted.