

AD-A228 755

REPORT DOCUMENTATION PAGE

DTIC FILE COPY

1a SECURITY CLASSIFICATION AUTHORITY SELECTED NOV 19 1990 S D C		1b RESTRICTIVE MARKINGS	
2b DECLASSIFICATION/DOWNGRADING SCHEDULE		3 DISTRIBUTION/AVAILABILITY OF REPORT This document has been approved for public release and sale, distribution of this document is unlimited.	
4 PERFORMING ORGANIZATION REPORT NUMBER Technical Report # 4		5 MONITORING ORGANIZATION REPORT NUMBER(S)	
6a NAME OF PERFORMING ORGANIZATION Naval Research Laboratory	6b OFFICE SYMBOL (If applicable) Code 6119	7a NAME OF MONITORING ORGANIZATION Office of Naval Research	
6c ADDRESS (City, State, and ZIP Code) Washington, DC 20375-5000		7b ADDRESS (City, State, and ZIP Code) 800 N. Quincy Street Arlington, VA 22217-5000	
8a NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research	8b OFFICE SYMBOL (If applicable) Code 1112AI	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-89-WX-24146	
8c ADDRESS (City, State, and ZIP Code) 800 North Quincy Street Arlington, VA 22217-5000		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO. 0601153N	PROJECT NO. R011-03-OK TASK NO. 412n006-01 WORK UNIT ACCESSION NO.
11 TITLE (Include Security Classification) Some One-Dimensional Molecular Dynamics Simulations of Detonation			
12 PERSONAL AUTHOR(S) M.L. Elert,* D.W. Brenner, C.T. White			
13a TYPE OF REPORT Technical (Interim)	13b TIME COVERED FROM 88/12/1 TO 89/9/30	14. DATE OF REPORT (Year, Month, Day) 89/8/28	15 PAGE COUNT
16 SUPPLEMENTARY NOTATION *U.S. Naval Academy, Annapolis, MD 21402-5026			
17 COSATI CODES		18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
		Solid-state Detonations, Chemical Reactions, Simulations	
		Molecular Dynamics, (7111)	
19 ABSTRACT (Continue on reverse if necessary and identify by block number) One dimensional molecular dynamics simulations of the onset of detonation have been performed using three-body potentials which accurately reproduce the effects of endothermic bond breaking and exothermic bond recombinations. A stable detonation wave of reasonable velocity can be produced from the impact of a plate upon a stationary array of diatomic molecules. The onset of detonation has been studied using a variety of potential forms including LEPS and Tersoff potentials. Properties of the detonation front, including from velocity, reaction zone width, and product distribution, have been studied as functions of barrier height, exothermicity, and other relevant potential function parameters. A model is proposed to explain the self-regulating detonation front velocities observed in these simulations in terms of atomic-scale kinetics.			
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT ☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT ☐ DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a NAME OF RESPONSIBLE INDIVIDUAL Dr. Carter T. White		22b TELEPHONE (Include Area Code) (202) 767-3270	22c OFFICE SYMBOL Code 6119

OFFICE OF NAVAL RESEARCH
CONTRACT # N00014-89-WX-24146
R&T CODE 412n006-01
TECHNICAL REPORT # 4

SOME ONE-DIMENSIONAL MOLECULAR DYNAMICS SIMULATIONS OF DETONATION*

Mark L. Elert^a, Donald W. Brenner^b, and C.T. White^b

^aU.S. Naval Academy
Annapolis, Maryland 21402

^bCode 6119
Naval Research Laboratory
Washington, D.C. 20375

Submitted for publication in the
Proceeding of the 1989 APS Topical Conference on
Shock Compression of Condensed Matter



Accession For	
NTIS CRABI	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Availability for Special
A-1	

*Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

ONE-DIMENSIONAL MOLECULAR DYNAMICS SIMULATIONS OF DETONATION

M. L. ELERT

Chemistry Department, U. S. Naval Academy, Annapolis, MD 21402

D. W. BRENNER and C. T. WHITE

Code 6110, Naval Research Laboratory, Washington, D.C. 20375

One-dimensional molecular dynamics simulations of the onset of detonation have been performed using three-body potentials which accurately reproduce the effects of endothermic bond breaking and exothermic bond recombinations. A stable detonation wave of reasonable velocity can be produced from the impact of a plate upon a stationary array of diatomic molecules. The onset of detonation has been studied using a variety of potential forms including LEPS and Tersoff potentials. Properties of the detonation front, including front velocity, reaction zone width, and product distribution, have been studied as functions of barrier height, exothermicity, and other relevant potential function parameters. A model is proposed to explain the self-regulating detonation front velocities observed in these simulations in terms of atomic-scale kinetics.

1. INTRODUCTION

The study of the onset of detonation at the molecular level is a difficult experimental task because of the short time scales involved and the destructive nature of the detonation process. Computer simulations are an obvious alternative, but the continuum methods which are so valuable for predictions at the macroscopic level are unable to give information regarding the primary chemical processes occurring in the reacting solid. Molecular dynamics simulations, on the other hand, are feasible under precisely the conditions of interest: short times and relatively small numbers of molecules. Indeed, molecular dynamics simulations of *nonreactive* shock waves have been performed for more than two decades. For the study of detonation, however, the simulations must explicitly include the chemical reactions which provide the energy to sustain the process. This requires the use of many-body potential functions which correctly reproduce the salient features of detonation reactions, but yet are simple enough to make the simulation computationally feasible.

We report two types of many-body potential

functions which are capable of accurately describing bond-breaking and bond-forming behavior in simple model systems of detonating solids. We demonstrate that one-dimensional molecular dynamics simulations using these potentials produce self-sustaining detonation waves with stable velocities independent of initiation conditions. Finally, we offer a molecular interpretation for the stability of the detonation velocities.

2. THE LEPS POTENTIAL

One form of potential function which can give qualitatively correct bond-breaking and bond-forming behavior for simple systems is the LEPS (London, Eyring, Polanyi, Sato) potential, developed in the 1930's to describe the H_2 potential energy surface. In the form devised by Tully¹, a three-body LEPS potential function can be described in terms of 2x2 matrices as follows: For each pair of atoms ij in the three-body system, a diagonal 2x2 matrix is constructed in which $H_{11} = V_{\text{bond}}(r_{ij})$ and $H_{22} = V_{\text{nonbond}}(r_{ij})$, where r_{ij} is the scalar distance between atoms i and j , V_{bond} is a typical diatomic bonding potential such

as a Morse function, and $V_{\text{Lennard-Jones}}$ is a weak van der Waals attraction between atoms i and j . The total potential energy of the three-body system, assuming nearest-neighbor interactions only, is then given as the lower eigenvalue of the 2×2 hamiltonian

$$H = H_{ij} + R^T H R$$

where R is the rotation matrix corresponding to a 60° two-dimensional rotation.

The LEPS potential is particularly suitable for one-dimensional simulations because in such systems each atom has exactly two nearest neighbors and therefore forms the central atom of a three-body system. The total potential energy of a one-dimensional crystal can be expressed as a sum of overlapping LEPS three-body potentials in which each atom (except those on the ends) appears as a left-hand, central, and right-hand member of three different trimers. The potential energy and atomic forces can be expressed analytically, so the method is computationally efficient.

We have recently reported² preliminary results of some one-dimensional molecular dynamics simulations using LEPS potentials designed to represent the features of detonating nitric oxide system, $2\text{NO} \rightarrow \text{N}_2 + \text{O}_2$. The initial NO crystal is arranged in the form ... NO ON NO ON ... so that formation of the appropriate products is possible. The NNO and NOO LEPS potentials are parameterized to match experimental exothermicities and calculated barrier heights for the three-body reactions $\text{N} + \text{NO} \rightarrow \text{N}_2 + \text{O}$ and $\text{N} + \text{O}_2 \rightarrow \text{NO} + \text{O}$. The detonation is initiated by impact of a small nitric oxide plate upon one end of the larger crystal. The simulation can readily be followed over the course of 20 picoseconds, during which time the detonation front propagates through several thousand atoms.

The most striking feature of these simulations is that the detonation velocity is independent of the plate impact velocity, as long as the impact

is strong enough to cause bond ruptures which initiate the detonation reaction. If the plate impact velocity is very high, the detonation is initially overdriven, but quickly returns to its steady-state value. If the impact velocity is too low to initiate detonation, it causes a gradually decaying nonreactive shock wave to propagate through the system. The lack of dependence of the detonation velocity on initiation conditions is in agreement with experimental observations of condensed-phase detonation.

Figure 1 shows the atomic density relative to the front for a plate impact speed of 6 km/sec, averaged over the course of a 20 picosecond simulation. Most of the chemical reactions take place in the high-density region within 400 Ångstroms of the front; behind this region the density falls off rapidly as the products and unreacted fragments separate. Further reaction in this gas-phase-like region is less likely because of the larger average interatomic spacing and small relative velocities which prevail in this region.




FIGURE 1
Density (g/mole/Å) versus position relative to the front for a plate impact speed of 6 km/sec.

3. THE TERSOFF POTENTIAL

One difficulty with the LEPS potential is the lack of a straightforward scheme for extending the method to higher dimensions with arbitrary numbers of near neighbors. We have therefore investigated the use of another form of empirical reactive potential based on the work of Tersoff. The Tersoff method uses a modified pair potential in which the bonding part of the interaction between a pair of atoms i and j is weakened by an amount proportional to the number of other near neighbors which compete for bonding to atom i , i.e.,

$$E = \sum_i E_i$$

where

$$E_i = (1/2) \sum_{j \neq i} (V_R(r_{ij}) - B_{ij} \cdot V_A(r_{ij}))$$

Here V_R represents a repulsive potential, V_A is an attractive potential, and B_{ij} represents a many-body coupling between bond ij the local coordination of atom i . These three functions can be implemented in a variety of ways^{4,5}, giving the Tersoff potential a great deal of flexibility. In particular, the coupling term B_{ij} may depend on the angle of a competing bond ik to the bond ij , thereby incorporating directed valence effects of importance for covalent compounds in two and three dimensions.

We have constructed a one-dimensional Tersoff potential to represent the generic reaction $2AB \rightarrow A_2 + B_2$, where the AB molecules have been given a bond strength of 4 eV and the A_2 and B_2 molecules have each been assigned a bond energy of 5 eV so that the overall reaction is exothermic. An initial AB molecular crystal was constructed in a manner identical to the nitric oxide simulations described above, ... AB BA AB ..., and the stability of the reactant solid was ensured by incorporating barrier heights of 0.5 eV for the two possible three-body reactions $A + AB \rightarrow A_2 + B$ and $B + BA \rightarrow B_2 + A$ (which are energetically equivalent in this case) and weak long-range van der Waals attractions between AB

molecules. Details of the potential function will be presented elsewhere.

As in the nitric oxide LEPS simulations described above, molecular dynamics simulations of the detonation of this system were carried out, with the detonation initiated by impact of a mass of variable velocity on the reactant solid. Although the density, exothermicity, and the potential surfaces in this case were quite different from those employed in the nitric oxide simulations, the results were qualitatively similar. For impact velocities ranging from 4 km/sec to 9 km/sec, the detonation velocity rapidly stabilized to a constant value of 12.6 km/sec. Exothermic chemical reactions in a high-density region behind the front provided the energy to sustain the detonation. Impact velocities less than 4 km/sec or greater than 9 km/sec did not produce detonation; in these cases the disturbance propagated through the chain as a gradually dissipating nonreactive shock wave.

4. VELOCITY REGULATION MECHANISM

The self-limiting detonation velocity observed in the simulations can be understood by investigating the behavior of an isolated three-body reaction process, $A + AB \rightarrow A_2 + B$. Classical trajectory calculations on the Tersoff potential surface described above lead to the results shown in Figure 2, where the velocity of product atom B is plotted as a function of the velocity given to an incident reactant atom A. The straight line marks the condition that the velocity of the outgoing atom equals the velocity of the incoming atom. The lower value of the curve is determined by the smallest velocity of atom A, which can overcome the potential barrier to reaction. The upper value of the curve represents the point beyond which the impact of atom A leads only to endothermic fragmentation ($A + A + B$) or to no reaction ($A + AB$).

If detonation in one dimension is viewed as being driven by a series of three-body reactions




FIGURE 2

Outgoing velocity of atom B following the reaction $A + AB \rightarrow A_2 + B$, as a function of the velocity of the incoming atom A relative to a stationary AB molecule. The straight line indicates equal incoming and outgoing velocities.

in which the products of one reaction collide with a molecule to initiate a new reaction, then the detonation regime will be limited to the range of impact velocities which cause initiation of the exothermic reactions $A + AB \rightarrow A_2 + B$ and $B + BA \rightarrow B_2 + A$. This analysis should also be valid in higher dimensions, as cross sections for bond dissociation generally increase for increasing atomic collision energies.

For incoming velocities near the low end of the reaction range, the velocity of the product atom is greater than that of the reactant atom. This is due to the net exothermicity of the reaction, and has the effect of accelerating the atomic velocities. For higher incident velocities, however, the outgoing atom has a lower velocity than the incoming atom. This occurs because the trajectory interacts with the repulsive wall of the potential surface, increasing the amount of energy deposited in the internal vibrational mode of the product molecule A_2 . If a series of isolated reactions in the orientation present in the chain described above are initiated with an atom anywhere within the range of reacting velocities, the series will "self regulate" to the point at which the two curves cross in Figure 2. Of course, in the solid

chain the detonation velocity will be significantly higher than the above analysis predicts, because the detonation wave is a collective disturbance which can propagate faster than individual atomic velocities. This effect can be accounted for quantitatively by including in the analysis the effect of the density of the chain, or more accurately the fraction of the unit cell length excluded by repulsive atomic core potentials.

5. CONCLUSION

Molecular dynamics simulations of detonation can be carried out in one dimension with many-body potential functions which incorporate realistic reaction dynamics. These simulations can provide valuable insight into the chemical mechanisms underlying the initiation and propagation of detonation. The Tersoff potential is particularly well suited to the extension of these investigations to two and three dimensions.

ACKNOWLEDGEMENTS

We are grateful to the Office of Naval Research for support of this work through contract number N0001489AF00001. We also thank the NRL Research Advisory Committee for a grant of computer time on the NRL Cray-XMP. M.L.E. acknowledges support from the NRL-USNA Cooperative Program for Scientific Interchange.

REFERENCES

1. J. Tully, J. Chem. Phys. 73 (1980) 6333.
2. M.L. Elert, D.M. Deaven, D.W. Brenner, and C.T. White, Phys. Rev. B 39 (1989) 1453.
3. J. Tersoff, Phys. Rev. B 37 (1988) 6991.
4. D.W. Brenner, M.L. Elert, and C.T. White, Incorporation of Reactive Dynamics in Simulations of Chemically-Sustained Shock Waves, this volume.
5. J. Tersoff, Phys. Rev. Lett. 61 (1988) 2879.

REPORTS DISTRIBUTION FOR ONR PHYSICS DIVISION
UNCLASSIFIED CONTRACTS

Director Defense Advanced Research Projects Agency Attn: Technical Library, TIO 1400 Wilson Blvd. Arlington Virginia 22209-2309	1 copy
Office of Naval Research Physics Division Office (Code 1112) 800 North Quincy Street Arlington, Virginia 22217-5000	2 copies
Office of Naval Research Director, Technology (Code 20) 800 North Quincy Street Arlington, Virginia 22217-5000	1 copy
Naval Research Laboratory Department of the Navy (Code 2625) Attn: Technical Library Washington, D.C. 20375-5000	1 copy
Office of the Director of Defense Research and Engineering Information Office Library Branch The Pentagon, Rm. 3E 1006 Washington, D.C. 20301	1 copy
U.S. Army Research Office Box 12211 Research Triangle Park North Carolina 27709-2211	2 copies
Defense Technical Information Center Cameron Station Alexandria, Virginia 22314	4 X copies
Director National Bureau of Standards Research Information Center Attn: Technical Library (Admin E-01) Gaithersburg, MD 20899	1 copy
Commander U.S. Army Belvoir Research, Development and Engineering Ctr. Attn: Technical Library (STRBE-BT) Fort Belvoir, Virginia 22060-5606	1 copy

interim/revised June 1988

ODDR&E Advisory Group on Electron Devices 201 Varick Street, 11th Floor New York, New York 10014-4877	1 copy
Air Force Office of Scientific Research Department of the Air Force Bolling AFB, DC 22209	1 copy
Air Force Weapons Laboratory Technical Library Kirtland Air Force Base Albuquerque, NM 87117	1 copy
Lawrence Livermore Laboratory Attn: Dr. W. F. Krupke University of California P.O. 808 Livermore, CA 94550	1 copy
Harry Diamond Laboratories Technical Library 2800 Powder Mill Road Adelphi, MD 20783	1 copy
Naval Weapons Center Technical Library (Code 753) China Lake, CA 93555	1 copy
Naval Underwater Systems Center Technical Center New London, CT 06320	1 copy
Commandant of the Marine Corps Scientific Advisor (Code RD-1) Washington, DC 20380	1 copy
Naval Ordnance Station Technical Library Indian Head, MD 20640	1 copy
Naval Postgraduate School Technical Library (Code 0212) Monterey, CA 93940	1 copy
Naval Missile Center Technical Library (Code 5632.2) Point Mugo, CA 93010	1 copy
Naval Ordnance Station Technical Library Louisville, KY 40214	1 copy

interim/revised June 1988

Commanding Office
Naval Ocean Research & Development Activity
Technical Library
NSTL Station, MS 39529 1 copy

Naval Oceans Systems Center
Technical Library
San Diego, CA 92152 1 copy

Naval Surface Weapons Center
Technical Library
Silver Spring, MD 20910 1 copy

Naval Ship Research and Development Center
Central Library (Codes L42 and L43)
Bethesda, MD 20084 1 copy

Naval Avionics Facility
Technical Library
Indianapolis, IN 46218 1 copy

a