

DTIC FILE COPY

2

Particle Simulations in Magnetospheric Plasmas
N00014-90-J-2000

AD-A228 893

Ken-Ichi Nishikawa
Department of Physics and Astronomy
The University of Iowa
Iowa City, IA 52242

11 October 1990
Final Technical Report for Period 1 October 1989 - 30 September 1990

Distribution Statement

Prepared for
Defense Technical Information Center
Building 5, Cameron Station
Alexandra, VA 22314

Naval Research Lab
4555 Overlook Avenue, S.W.
Washington, DC 20375-5000

DTIC
ELECTE
OCT 24 1990
S D
Lo D

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

90 10 22 056

TABLE OF CONTENTS

	Page
I. RESULTS FROM PRIOR SUPPORT ...	1
II. PUBLICATION FROM PRIOR SUPPORT ...	5
III. REFERENCES ...	6

I. RESULTS FROM PRIOR SUPPORT

1. Simulation Study of a New Mechanism for Excitation of Kinetic Waves in a Magnetoplasma

Nishikawa *et al.* [1988] have investigated the new ion-cyclotron-like waves by a localized transverse electric field by means of simulation with the assistance of the nonlocal kinetic theory. The linear theory shows that the growth rates of the kinetic Kelvin-Helmholtz (K-H) modes are strongly reduced with increasing $u = k_{\parallel}/k_y$, and they become unstable only when $b = k_y^2 \rho_i^2 < 1$ and $k_y L \approx 1$, where L is the scale length associated with the transverse electric field. On the other hand, the new modes have larger frequencies and become unstable at larger $b > 1$ and $k_y L \gg 1$ [Ganguli *et al.*, 1988].

Simulation results show that ion-cyclotron-like waves are excited in regions where $\mathbf{E} \times \mathbf{B}$ drift is localized. Linear growth rates of several modes are obtained from the wave analysis of the simulation. This linear analysis shows that the (0, 4) mode corresponds to large b , and large $k_y L$ has the maximum growth rate. Clearly, these are not K-H modes. Further, the simulation results show that density gradients help to enhance the growth rates. However, like the K-H mode, the real frequencies of this instability are approximately proportional to $k_y V_E^0$, where V_E^0 is the peak value of the $\mathbf{E} \times \mathbf{B}$ drift.

Nonlinear phenomena such as diffusion and coalescence of vortices are investigated. In the linear stage smaller vortices are generated and larger vortices with the lower frequencies are dominant in the nonlinear stage. In the nonlinear stage ions diffuse strongly due to large-scale vortices.

Recently, we have investigated the electrostatic waves driven by the combined effects of a localized transverse electric field and parallel electron drifts by means of simulation with the assistance of the nonlocal kinetic theory [Nishikawa *et al.*, 1989b,c,d].

We have performed a number of simulations for this instability. Simulation results show that electrostatic waves are excited in the regions where the $\mathbf{E} \times \mathbf{B}$ drift is localized in the simultaneous presence of parallel electron drifts and transverse electric fields. Simulations with only parallel electron drifts or transverse electric fields show that no instability grows out of the thermal noise. The simulation with both the parallel electron drift and the transverse electric field shows the growing waves out of the thermal noise. The Doppler shift due to the $\mathbf{E} \times \mathbf{B}$ drift can lower the phase velocity of waves along the magnetic field. Then this makes wave-particle resonance possible for smaller V_d^0 , which leads to this instability.

The nonlinear phenomena such as diffusion and coalescence of vortices are investigated [Nishikawa *et al.*, 1990a]. In the linear stage, smaller vortices are generated and larger vortices with the lower frequencies are dominant in the nonlinear stage. In the nonlinear stage the ions diffuse strongly due to the large scale vortices.

The spectrum for nonlinear stage was analyzed and the result is in good agreement with the satellite data. The details of this study will be reported in *J. Geophys. Res.* in near future.

STATEMENT "A" per ¹Dr. Peter Palmadesso
NRL/Code 4700.3
TELECON 10/22/90 VG

per call

Date	
A-1	

2. Beam Instability in the Foreshock

As an application of the simulation method used in the proposed research (Broadband electrostatic noise), the beam instability in the foreshock has been investigated. Electrons backstreaming into the Earth's foreshock generate waves near the plasma frequency by the beam instability. Two versions of the beam instability exist: the 'reactive' version in which narrowband waves grow by bunching the electrons in space, and the 'kinetic' version in which broadband growth occurs by a maser mechanism [Cairns, 1987a, b, and references therein]. Recently, Cairns [1987b] has suggested that (1) the backstreaming electrons have steep-sided 'cut-off' distributions which are initially unstable to the reactive instability, (2) the back-reaction to the wave growth causes the instability to pass into its kinetic phase, and (3) the kinetic instability saturates by quasilinear relaxation.

Cairns and Nishikawa [1989] have performed two-dimensional simulations of the reactive instability for Maxwellian beams and cutoff distributions. The results of the simulations are consistent with suggestions (1) and (2) above. In addition, we have demonstrated that the reactive instability is a bunching instability, and the reactive instability saturates and passes over into the kinetic phase by particle trapping. We found that the kinetic growth occurring after saturation of the reactive instability is presumably due to the spatially localized gradients in $y - v_{\parallel}$ phase space. Both simulation results and numerical solutions of the dispersion equation indicate that the center frequency of the intense narrowband waves near the foreshock boundary may be between $0.9\omega_{pe}$ and $0.98\omega_{pe}$, rather than being above ω_{pe} as previously believed. Nonlinear evolution of the electron plasma waves are investigated [Nishikawa *et al.*, 1990c]. They found the decay process in which a large-amplitude electron plasma wave driven by an electron beam decays into a backward propagating daughter wave and an ion acoustic wave.

Recent one-dimensional simulations of the electron beam-plasma instability [Dum, 1990] with parameters appropriate for the electron foreshock quantitatively tested the applicability of quasi-linear theory and other turbulence theories. A key requirement was that, in contrast to earlier simulations, the model at all times allowed the excitation of a large number of wave modes. Even in the case of an initially cold beam, quasi-linear theory is then applicable, except during a short period of strong beam modulation at the end of the reactive phase. Mode coupling is seen late in the kinetic phase but cannot prevent plateau formation.

It is important to investigate possible modifications of this evolution in a multidimensional system, especially for a low temperature beam, which initially allows the excitation of a large range of perpendicular wave numbers. We thus compare one- and two-dimensional simulations with large system size. The effects of increased magnetic field strength (e.g. along auroral field lines) is also investigated. This study will be reported in *J. Geophys. Res.*

3. Whistler Mode Driven by the Spacelab 2 Electron Beam

During the Spacelab 2 mission while an electron beam was being ejected from the shuttle, the Plasma Diagnostics Package (PDP) detected a clear funnel-shaped emission that is believed to be caused by whistler-mode emission from the electron beam [Gurnett *et al.*, 1986]. In order to understand the mechanism of this emission, the simulations have been performed using a three-dimensional magnetostatic code for low- β plasmas in which the beam electrons are initially located in the column [Nishikawa *et al.*, 1989a]. In order to simulate the continuous electron ejection from the shuttle, the simulations were also performed with the recycling of the beam electrons. The beam electrons excite whistler waves and lower hybrid waves. The brief fluid theory based on the magnetostatic code was checked with the simulation results. The propagating whistler mode was identified with the theory. The simulation results show that the quasi perpendicularly (the angle between the magnetic field and the wavenumber is larger than 50°) propagating whistler waves have larger amplitude whose real frequencies are smaller than the local electron cyclotron frequency. This fact is consistent with the fact that the funnel-shaped emission is observed below the electron cyclotron frequency away from the beam electron. The beam electrons initially in the column diffuse radially as well as slow down due to the $\mathbf{E} \times \mathbf{B}$ caused by the excited waves. The acceleration of the beam electrons also takes place due to the excited whistler waves.

In order to compare with the PDP data, the local magnetic fields $B_{x,y}$ and the perturbed electric fields $E_{y,z}$ are diagnosed at the several points in the simulation system. The results show that the waves are radially excited by the beam electrons localized in the center of the system. The wave spectra of the electric fields $E_{y,z}$ diagnosed at $x = 31\lambda_e$, $y = 16\lambda_e$, and $z = 17\lambda_e$ show the several kinds of waves generated by the beam electrons. The analysis shows that the lower hybrid waves and whistler waves are excited. The frequency range of these spectra extends from ω_{pi} to beyond ω_{pe} which is in qualitative agreement with the PDP data. As the PDP data show, the intense electrostatic narrowband emission around the electron plasma frequency has been observed dominantly in the spectra of the electric field (E_z) with and without the recycling of the beam electrons. This wave is identified as the parallel (quasi parallel) whistler wave. This simulation result is in good agreement with the PDP data [Gurnett *et al.*, 1986]. The spectra of the magnetic fields diagnosed at the same position show that whistler modes are excited. The cB/E was calculated for the whistler mode. In the case without the recycling of the beam electrons, the cB/E is approximately 0.35 for the whistler mode (around $0.368\omega_{pe}$), which is smaller than that calculated from the PDP data [Gurnett *et al.*, 1986].

Recently we have developed a 3-D electromagnetic and relativistic particle code without solving Poisson's equation. "Radiating" boundaries are introduced besides periodic boundary conditions. These boundary conditions allow radiation to escape freely out of the simulation domain without reflection. Periodic boundaries are applied along z direction. Radiating boundaries are used for the boundaries perpendicular to the magnetic field. The simulation domain is a mesh of dimension $100 \times 100 \times 200$ with a few million particles. Whistler emission as shown by wave data obtained from the Spacelab 2 mission will be the main topics of this simulation study.

4. Simulation of Electron Cyclotron Harmonic Waves by AMPTE Lithium Release

We have studied the generating electrostatic turbulence in the lithium cloud due to solar wind particles which are flowing through it [Nishikawa *et al.*, 1990b]. Several simulations have been performed in order to understand the processes generating electron cyclotron harmonic waves.

The simulations show that electron cyclotron harmonic waves are excited by mainly the solar wind electrons. The spectra of perturbations of electrostatic potential exhibit electron cyclotron harmonic waves.

The overall frequency range is obtained by observing the electric fields at some points in the simulation region. Power spectra of the electric fields exhibit electron cyclotron frequency and higher harmonics which is similar to the spectrum obtained by the satellite.

Possible mechanisms of generating electron cyclotron harmonic waves have been discussed in the previous work [Roeder *et al.*, 1987]. In this simulation study, the possible energy sources of the instability are included automatically. The relative velocities among different species become the energy source of instabilities. The relative velocity between solar wind electrons and lithium ions would be one possible energy source for the excitation of ECH waves. The fact that the instability can be excited without new-born cold electrons support this mechanism. Theoretically, the mode couplings among harmonics occur mainly due to the electron perpendicular drift, which lead to the instability [Tataronis and Crawford, 1970].

According to Figure 1 by Gurnett *et al.* [1986b], the higher frequency ranges of waves look very similar at both cases with $\bar{n}_{li} \approx \bar{n}_{pr}$ and $\bar{n}_{li} \ll \bar{n}_{pr}$. In this study, we found that ECH waves are excited also in the case of $\bar{n}_{li} = \bar{n}_{pr}$. The satellite data show that ECH waves can be observed only in some range of lithium ion density ($0.03 - 0.18 \text{ cm}^{-3}$) associated with lithium drift energy [Roeder *et al.*, 1987]. The observations show that in the case of $\bar{n}_{li} \approx \bar{n}_{pr}$ ECH waves are not found. The simulation results show that in the same case of $\bar{n}_{li} = \bar{n}_{pr}$ ECH waves are excited. This discrepancy may be explained by the setting of simulation geometry due to the two-dimensional code. If the simulation plane is set nearly parallel to the magnetic fields ($B_{oy}/B_o \leq 1$), the angle between the magnetic fields and the drift velocity of hot electrons and protons becomes very small. Then, in the case of $\bar{n}_{li} = \bar{n}_{pr}$, the parallel drift of solar wind protons and electrons may excite ion acoustic waves more dominantly than ECH waves, which is not included in this study [Gurnett *et al.*, 1986b; Ma *et al.*, 1987].

Further study is required on the observed plasma velocity distributions and comparisons of simulation wave results with the wave power spectra. More simulations are required for further progress in the interpretation of electrostatic noise.

II. PUBLICATIONS FROM PRIOR SUPPORT

(as of December 18, 1989)

Cairns, I. H. and K.-I. Nishikawa,

Simulations relevant to the beam instability in the foreshock plasma, *J. Geophys. Res.*, **94**, 79, 1989.

Nishikawa, K.-I., L. A. Frank, and C. Y. Huang,

Three-dimensional simulation of whistler mode excited by the Spacelab 2 electron beam, *J. Geophys. Res.*, **94**, 6855, 1989a.

Nishikawa, K.-I., G. Ganguli, Y. C. Lee, and P. Palmadesso,

Simulation of electrostatic ion instabilities in the presence of parallel currents and transverse electric fields, *Physics of Space Plasma (1988), SPI Conference Proceedings and Reprints Series, Number 8*, T. Chang, G. B. Crew, and J. R. Jasperse, eds., Scientific Publishers, Inc., Cambridge, MA, 1989b, p. 405; also in *Proceeding of an International School and Workshop on Plasma Astrophysics, ESA SP-285 (Vol. 1)* p. 385, ESA Paris, France, 1989c.

Nishikawa, K.-I., G. Ganguli, Y. C. Lee, and P. J. Palmadesso,

Simulation of electrostatic turbulence due to sheared flow parallel and transverse to the magnetic field, *J. Geophys. Res.*, **95**, 1029, 1990a.

Nishikawa, K.-I., J. L. Roeder, and H. C. Koons,

Simulation of electron cyclotron harmonic waves by AMPTE lithium release, to be re-submitted to *J. Geophys. Res.*, 1990b.

Nishikawa, K.-I. and I. H. Cairns,

Simulation of the nonlinear evolution of electron plasma waves, submitted to *J. Geophys. Res.*, 1990c.

C. T. Dum and K.-I. Nishikawa,

Simulation studies of the electron beam-plasma instability, to be submitted to *J. Geophys. Res.*, 1990d.

REFERENCES

- Cairns, I. H., The electron distribution function upstream from the Earth's bow shock, *J. Geophys. Res.*, **92**, 2315, 1987a.
- Cairns, I. H., A theory for Langmuir waves in the electron foreshock, *J. Geophys. Res.*, **92**, 2329, 1987b.
- Cairns, I. H. and K.-I. Nishikawa, Simulations relevant to the beam instability in the foreshock. *J. Geophys. Res.*, **93**, October, 1988. (in press) external magnetic field, *Phys. Fluids*, **28**, 761, 1985a.
- Dum, C. T., Simulation studies of plasma waves in the electron foreshock: The generation of Langmuir waves by a gentle bump-on-tail electron distribution, *J. Geophys. Res.*, **95**, 8095, 1990.
- Ganguli, G. and P. Palmadesso, Electrostatic ion instabilities in the presence of parallel currents and transverse electric fields, *Geophys. Res. Lett.*, **15**, 103, 1988.
- Gurnett, D. A., W. S. Kurth, J. J. Steinberg, P. M. Banks, R. I. Bush, and W. T. Raitt, Whistler-mode radiation from the Spacelab 2 electron beam, *Geophys. Res. Lett.*, **13**, 225, 1986a.
- Gurnett, D. A., T. Z. Ma, R. R. Anderson, O. H. Bauer, G. Haerendel, B. Hausler, G. Paschmann, R. A. Treumann, H. C. Koon, R. Holzworth, and H. Luhr, Analysis and interpretation of the shocklike electrostatic noise observed during the AMPTE solar wind lithium releases, *J. Geophys. Res.*, **91**, 1301, 1986b.
- Ma, T. Z., D. A. Gurnett, and N. Omid, An analysis of shocklike electrostatic noise observed during AMPTE solar wind ion release, *J. Geophys. Res.*, **92**, 2555, 1987.
- Nishikawa, K.-I., L. A. Frank, and C. Y. Huang, Three-dimensional simulation of whistler mode excited by the Spacelab 2 electron beam, *J. Geophys. Res.*, **94**, 6855, 1989a.
- Nishikawa, K.-I., G. Ganguli, Y. C. Lee, and P. Palmadesso, Simulation of ion-cyclotron-like modes in a magnetoplasma with transverse inhomogeneous electric field, *Phys. Fluids*, **31**, 1568, 1988.
- Nishikawa, K.-I., G. Ganguli, Y. C. Lee, and P. Palmadesso, Simulation of electrostatic ion instabilities in the presence of parallel currents and transverse electric fields, *Physics of Space Plasma (1988), SPI Conference Proceedings and Reprints Series, Number 8*, T. Chang, G. B. Crew, and J. R. Jasperse, eds., Scientific Publishers, Inc., Cambridge, MA, 1989b, p. 405; also in *Proceedings of an International School and Workshop on Plasma Astrophysics, ESA SP-285 (Vol. 1)* p. 385, ESA Paris, France, 1989c.

- Nishikawa, K.-I., G. Ganguli, Y. C. Lee, and P. J. Palmadesso, Simulation of electrostatic turbulence due to sheared flow parallel and transverse to the magnetic field, *J. Geophys. Res.*, 95, 1029, 1990a.
- Nishikawa, K.-I., J. L. Roeder, and H. C. Koons, Simulation of electron cyclotron harmonic waves by AMPTE lithium release, to be resubmitted to *J. Geophys. Res.* 1990b.
- Nishikawa, K.-I., I. H. Cairns, and D. A. Gurnett, Simulation of the nonlinear evolution of electron plasma waves, to be submitted to *J. Geophys. Res.* 1990c.
- Roeder, J. L., H. C. Koons, R. H. Holzworth, R. R. Anderson, O. H. Bauer, D. A. Gurnett, G. Haerendel, B. Häusler, and R. Treumann, Electron cyclotron harmonic waves observed by the AMPTE-IRM plasma wave experiment following a lithium release in the solar wind, *J. Geophys. Res.*, 92, 5768, 1987.
- Tataronis, J. A. and F. W. Crawford, Cyclotron harmonic wave propagation and instability: I. Perpendicular propagation, *J. Plasma Phys.*, 4, 231, 1970.