

AD-A229 119

DTIC FILE COPY

①

CONTRACT NO.: 87PP7852

TITLE: The 3D Structures of Some Diarrheal Causing Bacterial
Toxins

PRINCIPAL INVESTIGATOR: Martin Sax

PI ADDRESS: Veterans Administration Medical Center,
Pittsburgh, Pennsylvania 15240

REPORT DATE: July 1, 1990

TYPE OF REPORT: Annual

PREPARED FOR: U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND
FORT DETRICK
FREDERICK, MARYLAND 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

20030206211

DTIC
ELECTE
OCT 15 1990
S B D

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Veterans Administration Medical Center		6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) Pittsburgh, PA 15240			7b. ADDRESS (City, State, and ZIP Code)		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION U.S. Army Medical Research & Development Command		8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER 87PP7852		
8c. ADDRESS (City, State, and ZIP Code) Fort Detrick Frederick, MD 21702-5012			10. SOURCE OF FUNDING NUMBERS		
PROGRAM ELEMENT NO. 62787A	PROJECT NO. 3M1- 62787A871	TASK NO. AA	WORK UNIT ACCESSION NO. 387		
11. TITLE (Include Security Classification) (U) The 3D Structures of Some Diarrheal Causing Bacterial Toxins					
12. PERSONAL AUTHOR(S)					
13a. TYPE OF REPORT Annual		13b. TIME COVERED FROM 7/1/89 TO 6/31/90		14. DATE OF REPORT (Year, Month, Day) 1990 July 1	
15. PAGE COUNT 7					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	RA 1; X-ray Diffraction; Crystallography; Protein Toxin; Enterotoxin; PO; BD		
20	02				
06	01				
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The overall is to determine the 3D structures of the <i>S. aureus</i> enterotoxins by X-ray diffraction. A 3.5A maps of SEB has been prepared, and the tracing of the main chain is in progress.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT ☐ UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT ☐ DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Mary Frances Bostian			22b. TELEPHONE (Include Area Code) 301-663-7325		22c. OFFICE SYMBOL SGRD-RMT-S

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

Where copyrighted material is quoted, permission has been obtained to use such material.

Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).


For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institute of Health.

Martin Sax 7/3/90
PI Signature Date

INTRODUCTION:

Progress in understanding the biological activities of the diarrheal causing staphylococcal enterotoxins (SEs) at the molecular level now has reached the point where detailed knowledge of their 3D structures is required. Much biochemical data has been accumulated about the SEs, including the amino sequences of several of them (1). A hotly pursued goal in this field presently is to explain the immunogenic activity of the SEs, in particular, their powerful stimulation of T cell proliferation (2). The first step in this process involves the formation of a binary complex between these SE and a MHC class II molecule. The latter is situated on the surface of the accessory cell, whose function is to present the binary complex to a V-beta receptor on the T cell surface, where a ternary complex consisting of the binary complex and the V-beta molecule, then forms. T-cell proliferation ensues. The process does not occur without the prior formation of the binary complex. Relevant questions about this process remain to be answered at the molecular level. Which groups on the enterotoxin molecule make binding contacts with the MHC-II molecule? Which groups interact with the V-beta receptor? What molecule structural features of the enterotoxin molecule are crucial for T cell stimulatory activity? A combination of mitagenetic and crystallographic studies relating activity on 3D structure is the most obvious approach to solving this problem. In addition the 3D structure is vitally important for interpreting other biochemical data at the molecular level (1).



Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

BODY:

At first we tried the method of single isomorphous replacement (SIR) to the derivation of structure factor phases for use in the preparation of a 3D map of the SEB molecule. Combining the SIR and anomalous data with solvent leveling yielded a 4A map which showed the molecular boundary, but efforts to trace the main chain were fruitless even after the phases were extended to 3.5A. An MIR (multiple isomorphous replacement) map next was constructed with data from seven heavy atom derivatives. Anomalous data from two of the derivatives also were included. The resolution of the map was 4A and the figure of merit was 0.66. Application of the solvent leveling algorithm (3) extended the resolution to 3.7A and yielded the figure of merit of 0.83. It was possible to trace approximately 75 percent of the main chain. A polyalanine model was fitted to the main chain trace on a Silicon Graphics unit. The phases were extended stepwise to 3A by repeated application of the same method. The phase extended map brought out important new features of the protein molecule. The helical regions were defined better, and the main chain certainly was altered in at least one place. Next a combined phase difference map (4) was prepared in an attempt to bring out new and improved structural information.

Our plan is to complete the tracing of the main chain and to fit the side chains. This will be accomplished by utilizing 2Fo-Fc until a completed model is obtained. The derived model will be refined by restrained least squares.

CONCLUSIONS:

Any conclusions about the 3D structure of SEB at this point must be regarded as being highly tentative since the structure determination is only partially completed. What we see presently is that there are three distinct domains of varying size in the SEB molecule, that the percentage of alpha helical regions is small, that the beta strands are considerably more prevalent than the helices, and that a large fraction of the molecules consist of a periodic secondary structural feature.

REFERENCES:

1. Ono et al. In Bacterial Toxins, Vol. 4, Ed. C. Hardegree and A. Tu, Macel Dekker, Inc., New York, NY (1988).
2. Marrack, P. & Kappler. J. Science 248, 705-711 (1990).
3. Wang, B.C. Methods of Enzymology 115, 90-112 (1985).
4. Read, R.J. Acta Cryst. A42, 140-149 (1986).