

**DTIC FILE COPY**

© 1989 Elsevier Science Publishers B.V. (Biomedical Division)  
FRTL-5 today. F.S. Ambesi-Impiombato, H. Perrild, editors

**S** **D** **D**  
**ELECTE**  
**MAR 29 1991**  
**G** **D** 191

**MONENSIN INCREASES CYTOSOLIC CALCIUM IN FRTL-5 CELLS**

**AD-A233 409**

CLARA AMBROZ, HENRY FEIN, AND ROBERT SMALLRIDGE

Department of Clinical Physiology, Walter Reed Army Institute of Research, Washington, D.C. 20307-5100 (USA)

**INTRODUCTION**

Calcium is thought to play an important role in stimulus-secretion coupling in secretory cells. Monensin, a Na<sup>+</sup>-specific ionophore which raises intracellular sodium content, also increases the cytosolic free calcium concentration [Ca<sup>2+</sup>]<sub>i</sub> in several cells (1,2). The objectives of the present study were (a) to examine the effect of increasing [Na<sup>+</sup>]<sub>i</sub> on cytosolic Ca<sup>++</sup> in FRTL-5 cells, and (b) to determine the relative contributions of extra- versus intracellular calcium in producing any such changes.

**MATERIALS AND METHODS**

Materials

Cells. FRTL-5 cells were maintained in 95% air - 5% CO<sub>2</sub> at 37°C. The medium used was Coon's modified Ham's F-12 supplemented with 5% calf serum, 1 mM non-essential amino acids, TSH (1 mU/ml), insulin (10 µg/ml), transferrin (5 µg/ml), glycyl-L-histidyl-L-lysine acetate (2 ng/ml), somatostatin (10 ng/ml), and cortisol (1 nmol/l). Cells were plated in petri dishes at 2x10<sup>5</sup> cells/ml, fed twice weekly, and passaged every 7-8 days.

Methods

Loading of Cells. Cells were washed, resuspended at 5x10<sup>7</sup> cells/ml in HBSS, and incubated with Indo-1 AM (5 µM) for one hour at 37°C in a shaking water bath. Cells were resuspended at 2.5x10<sup>6</sup> cells/ml in HBSS with 0.02% BSA for study.

Fluorescence Measurements. Two ml aliquots were placed in quartz cuvettes and fluorescence recordings obtained on an SLM 8000 spectrofluorometer. The excitation wavelength was 350 nm. Emission was recorded as the rate of intensities at 405 and 480 nm. Calculations of [Ca<sup>2+</sup>]<sub>i</sub> were determined as previously reported (3).

**RESULTS**

Extracellular Ca<sup>++</sup> Present

Baseline Calcium. Cytosolic calcium content under basal conditions averaged 220 ± 6 nM (n=50).

Accession For	
NTIS - CRUI	00
DTIC TAB	00
Unannounced	
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
	A1201

QUANTITY  
180-00280  
3

This document has been approved for public release and sale; its distribution is unlimited.

**91 3 08 010**

Monensin Effect. The addition of monensin produced a dose dependent increase in  $[Ca^{2+}]_i$  (Table).

Calcium Blockers. Three classes of  $Ca^{++}$  channel blockers (nifedipine, verapamil, diltiazem) had no effect on inhibiting the monensin stimulation of  $[Ca^{2+}]_i$ . Similar results were obtained when TMB-8 and ryanodine were used to block the release of intracellular  $Ca^{++}$  stores.

$Na^+/K^+$  ATPase. Ouabain ( $10^{-3}M$ ) had no effect on the monensin stimulation of  $[Ca^{2+}]_i$ .

Extracellular Cations Absent

Monensin Effect. In the absence of extracellular  $Ca^{2+}$ , monensin produced a negligible change in  $[Ca^{2+}]_i$  (Table). Subsequent addition of ionomycin produced a release of  $Ca^{2+}$  from intracellular stores. In  $Na^+$  free buffer, the  $[Ca^{2+}]_i$  response to monensin was inhibited by 80% (Table).

TABLE: MONENSIN EFFECT ON CYTOSOLIC CALCIUM (% Change)

Monensin Dose (M)	$Ca^{2+}$	+	-	+
	$Na^+$	+	+	-
$10^{-4}$		$420 \pm 54\%$	$6 \pm 3\%$	$79 \pm 10\%$
$10^{-6}$		$81 \pm 11\%$	N.D.	$15 \pm 15\%$
$10^{-8}$		$32 \pm 6\%$	N.D.	N.D.

$Ca^{2+} = 1.27 \text{ mM}; Na^+ = 140 \text{ mM}; \text{N.D.} = \text{not tested}$

DISCUSSION

This study demonstrates that changes in intracellular sodium dramatically increase the cytosolic free calcium concentration in FRTL-5 cells. The source of calcium is virtually all extracellular. The calcium influx is independent of calcium channels or  $Na^+/K^+$  ATPase activity. While the response is highly dependent upon extracellular  $Na^+$ , a small effect of monensin was independent of  $Na^+$ .

REFERENCES

1. Grinstein S, Goetz JD (1985) *Biochim Biophys Acta* 819:267
  2. Schackmann RW, Chock PB (1986) *J Biol Chem* 261:8719
  3. Grynkiewicz G, Poenie M, Tsien RY (1985) *J Biol Chem* 260:3440
- The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or reflecting the opinions of the Department of the Army or the Department of Defense.