


AD-A234 813 REPORT DOCUMENTATION PAGE

UNCL		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution is unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NMRI 91-14		7a. NAME OF MONITORING ORGANIZATION Naval Medical Command	
6a. NAME OF PERFORMING ORGANIZATION Naval Medical Research Institute	6b. OFFICE SYMBOL (if applicable)	7b. ADDRESS (City, State, and ZIP Code) Department of the Navy Washington, DC 20372-5120	
6c. ADDRESS (City, State, and ZIP Code) 8901 Wisconsin Avenue Bethesda, MD 20889-5055		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Naval Medical Research & Development Command	8b. OFFICE SYMBOL (if applicable)	10. SOURCE OF FUNDING NUMBERS	
8c. ADDRESS (City, State, and ZIP Code) 8901 Wisconsin Avenue Bethesda, MD 20889-5044		PROGRAM ELEMENT NO. 61102A	PROJECT NO. 3M161102BS13
		TASK NO. AK111	WORK UNIT ACCESSION NO. DA313955
11. TITLE (Include Security Classification) Taxonomy			
12. PERSONAL AUTHOR(S) Weiss E			
13a. TYPE OF REPORT Book chapter	13b. TIME COVERED FROM _____ TO _____	14. DATE OF REPORT (Year, Month, Day) 1991	15. PAGE COUNT 3
16. SUPPLEMENTARY NOTATION In: Manual of Clinical Microbiology 5th edition, editor in chief, Albert Balows; editors, William J. Hausler, Jr ...[et al.]. Washington, D.C.: American Society for Microbiology, 1991. pp. 1033-1035			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	human pathogens; spotted fever group; genotypes; Rickettsiae; Ehrlichia; Proteobacteria; Chlamydiae; serologic relationships; bacteriology; taxon; phylogeny; phylogenetic analysis; phenotypes; morphology	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)			
			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Physician Librarian		22b. TELEPHONE (Include Area Code) (301) 295-2188	22c. OFFICE SYMBOL MRL/NMRI

Section VIII. Rickettsiae and Chlamydiae

Chapter 103

Taxonomy

EMILIO WEISS

Section 9 of *Bergey's Manual of Systematic Bacteriology* (6) describes in detail the orders *Rickettsiales* and *Chlamydiales*. The *Rickettsiales*, even though separated from the endosymbionts in this edition of *Bergey's Manual*, represent a complex taxon, which includes the families *Rickettsiaceae*, *Bartonellaceae*, and *Anaplasmataceae*. The *Chlamydiales*, on the other hand, include a single family and a single genus, possibly underclassified. Numerous observations, isolations of new species, and phylogenetic analyses made during the past decade and for the most part after the publication of *Bergey's Manual* suggest that we should take another look at the classification of the family *Rickettsiaceae* and the genus *Chlamydia*.

Table 1 lists the species in the family *Rickettsiaceae* and in the genus *Chlamydia*, including species that have recently been added, and summarizes the information thus far obtained through comparisons of 16S rRNA sequences (20).

Among the changes that have occurred during the past decade, human infection in Europe with the spotted fever group rickettsia *Rickettsia conorii* was found to occur more frequently and to produce a more severe disease than in the past (10, 11). To the large number of rickettsial species in the spotted fever group, some established pathogens and others most likely commensals in the invertebrate host, two have been added, the pathogen *R. japonica* (14) and the commensal *R. helvetica* (8). The commensal *R. bellii* was isolated and described as a rickettsia that is possibly not a member of the spotted fever group (9).

The most extensive additions to our information are possibly those that involve the genus *Ehrlichia*. Ristic et al. (12) demonstrated that there is a small but highly significant degree of serologic relationship between the dog pathogen *Ehrlichia canis* and the human pathogen *Rickettsia sennetsu*. This finding was quickly followed by the observations that the two microorganisms strongly resemble each other in morphology and mode of development (3). As a result the sennetsu agent was reclassified in *Bergey's Manual* as *Ehrlichia sennetsu*. A few years later the etiologic agent of Potomac horse fever (equine monocytic ehrlichiosis) was isolated and found to be distantly related to *E. canis* and more closely related to *E. sennetsu* (4, 5). The evolutionary history of *E. sennetsu* and *E. risticii*, two microorganisms that are biologically similar but that differ widely in geographic distribution and pathogenesis for vertebrate hosts, well deserves to be investigated. A possible link between the two microorganisms is suggested by the recent discovery (discussed in chapter 106) that *E. canis*, or a closely related microorganism, is also a human pathogen.

Also worth mentioning is the similarity between an-

other dog pathogen and a potential human pathogen. *Neorickettsia helminthoeca* is a dog pathogen that occurs in the U.S. West Coast states. Dogs acquire the disease by eating raw salmon infested with the metacercariae of a trematode, which in turn is infected with *N. helminthoeca* (6). Fukuda et al. (1) presented evidence that a disease involving a similar cycle may occur in Japan among people who eat improperly cooked fish. The fish, the grey mullet, is also often infested with a trematode that carries a bacterium similar to *N. helminthoeca*.

To the genus *Chlamydia*, the important human pathogen *C. pneumoniae* has been added (2).

The study of the phylogeny of the *Rickettsiaceae* and of chlamydiae conducted thus far has produced some results that confirm observations made on phenotypic similarity among certain species, while others contrast sharply with our present taxonomic concepts. For example, as expected, *Rickettsia prowazekii*, *R. typhi*, and *R. rickettsii* were shown to have very similar rRNA sequences (differences of <2%, smaller than the typical difference between the genera *Escherichia* and *Salmonella*) (15). These three rickettsial species belong to the alpha subdivision of the purple bacteria (20), recently named *Proteobacteria* (13). In this subdivision, the rickettsial species are specifically, although distantly, related to *Ehrlichia risticii* (15). This came as a surprise, since rickettsiae and the monocytic ehrlichiae differ in morphology and intracellular location, cytoplasm in the case of rickettsiae and phagosome in ehrlichiae (6). On the other hand, it has recently been shown that ehrlichiae and rickettsiae resemble each other in in vitro substrate utilization and gain in ATP content from this activity (18, 19). The rickettsiae are not specifically related to *Rochalimaea quintana*. This microorganism is also a member of the alpha subdivision of *Proteobacteria*, but belongs to subgroup 2 and is specifically related to the plant-associated agrobacteria and rhizobacteria (17). This is also unexpected since *R. prowazekii* and *R. quintana* have historically been transmitted by the same arthropod vector, the body louse, and the two DNAs hybridize with each other to the extent of 25 to 33% (7).

Not surprising, because of marked differences in phenotypic characteristics, was the finding that *Coxiella burnetii* is unrelated to the three species of rickettsia. It belongs in the gamma subdivision of the *Proteobacteria* and in this group it shows a rather distant but specific relationship to the genus *Legionella*. The tick commensal, *Wolbachia persica*, was also shown to belong in the gamma subdivision and to be phylogenetically related to the *Coxiella-Legionella* cluster (15).

Most astonishing was the finding that chlamydiae have no close relatives among the eubacteria (16). They represent a hitherto unrecognized major eubacterial group, peripherally related only to the planctomyces. These

TABLE 1. List of species in the family *Rickettsiaceae* and genus *Chlamydia* and summary of present knowledge of their phylogeny based on 16S rRNA sequences*

Genus	Group	Species	Proteobacterial subdivision ^b	Related genus ^b	
<i>Rickettsia</i>	Typhus	<i>R. prowazekii</i> , ^c <i>R. typhi</i> ^c	Alpha	<i>Ehrlichia</i>	
		<i>R. canada</i>	Unknown	Unknown	
	Spotted fever	<i>R. rickettsii</i> ^c	Alpha	<i>Ehrlichia</i>	
		<i>R. conorii</i> , ^c <i>R. sibirica</i> , ^c <i>R. australis</i> ^c	As above	As above	
		<i>R. akari</i> , ^c <i>R. japonica</i> ^c <i>R. montana</i> , <i>R. parkeri</i> , <i>R. rhipicephali</i>	As above Unknown	As above Unknown	
	Uncertain	<i>R. helvetica</i>	Unknown	Unknown	
		<i>R. bellii</i>	Unknown	Unknown	
		<i>R. tsutsugamushi</i> ^c	Unknown	Unknown	
	<i>Rochalimaea</i>	Scrub typhus	<i>R. quintana</i> ^c	Alpha, subgroup 2	Agrobacteria, rhizobacteria
			<i>R. vinsonii</i>	Unknown	Unknown
<i>Ehrlichia</i>	Monocytic	<i>E. risticii</i> ^d	Alpha	<i>Rickettsia</i>	
		<i>E. canis</i> , ^{c,d} <i>E. sennetsu</i> ^c	As above	As above	
	Granulocytic	<i>E. equi</i> , ^d <i>E. phagocytophila</i> ^d	Unknown	Unknown	
<i>Cowdria</i>		<i>C. ruminantium</i> ^d	Unknown	Unknown	
<i>Neorickettsia</i>		<i>N. helminthoeca</i> ^d	Unknown	Unknown	
<i>Coxiella</i>		<i>C. burnetii</i> ^{c,d}	Gamma	<i>Legionella</i>	
<i>Wolbachia</i>		<i>W. persica</i>	Gamma	<i>Coxiella</i> , <i>Legionella</i>	
		<i>W. pipientis</i> , <i>W. melophagi</i>	Unknown	Unknown	
<i>Rickettsiella</i> ^c		<i>R. popilliae</i> , <i>R. grylli</i> , <i>R. chironomi</i>	Unknown	Unknown	
<i>Chlamydia</i>		<i>C. trachomatis</i> , ^c <i>C. psittaci</i> ^{c,d}	None (not a proteobacterium)	<i>Planctomyces</i>	
		<i>C. pneumoniae</i> ^c	As above	As above	

* The *Rickettsiales* families *Bartonellaceae* and *Anaplasmataceae* are not included. Not all of the species listed are discussed in subsequent chapters.

^b "As above" below a proteobacterial subdivision or genus designation means that it is reasonable to assume that the designation above is applicable. Such an inference cannot be made for some of the other species of the same genus.

^c Established human pathogen.

^d Animal pathogen.

^e Pathogenic for invertebrate host.

two groups of bacteria have very little in common phenotypically, except that the cell wall in both cases contains no peptidoglycan.

Much remains to be done in the field of phylogeny of the rickettsiae. If this work is extended to other pathogens of the spotted fever group, such as *R. conorii* or *R. sibirica*, few surprises are expected, because of the solid base of phenotypic information, which suggests that these rickettsiae are closely related. However, *R. tsutsugamushi* is sufficiently different from the other rickettsiae to deserve separate phylogenetic analysis. Despite the large body of information that we have on this species, in the absence of such a study we cannot venture to say that it is a proteobacterium.

The place of taxonomy in a manual of clinical microbiology is not clear. When phylogenetic information reinforces what we have learned by the study of phenotypic characteristics, it encourages us to proceed as we have. When a conflict develops between genotypic and phenotypic information, for the immediate purpose of isolation and identification of pathogens, it is preferable to pay greater attention to the phenotypic characteristics. However, genotypic information cannot be ignored. A conflict may suggest that important biologic or ecologic characteristics of the microorganisms involved are yet to be uncovered.

I am indebted to Gregory A. Dasch for his helpful discussions.

The preparation of this chapter was in part supported by the Naval Medical Research and Development Command, Department of the Navy, work unit no. 3M161102.BS13.AK.111.

ADDENDUM IN PROOF

Taxonomy and the study of the bacterial genome have recently acquired new significance as the polymerase chain reaction (PCR) is being developed for the rapid identification of infectious agents (K. E. Hechemy, D. Paretsky, D. H. Walker, and L. P. Mallavia, ed., *Ann. N.Y. Acad. Sci.* 590:1-586, 1990). For example, although the scrub typhus group of rickettsiae is still represented by a single species, *R. tsutsugamushi*, there is considerable variation in antigenic properties among the various strains. This variation is not necessarily accompanied by marked differences in virulence for humans. Heterogeneity has been attributed to variability of certain surface components that occur in a background of strong homology (E. V. Oaks, R. M. Rice, D. J. Kelly, and C. K. Stover, *Infect. Immun.* 57: 3116-3122, 1989). It is therefore important, for the purpose of the PCR test, to select a region on the genome that is relatively stable and recognizes most, if not all, strains. In the case of the typhus and spotted fever group rickettsiae, there is sufficient similarity in their genomes for the selection of a region that detects several species, such as *R. prowazekii*, *R. typhi*, *R. rickettsii*, and *R. conorii*. If the test is positive, the identity of the infectious agent can, in most cases, be inferred on the basis

of geography and other epidemiological considerations. Although more specific PCR tests can be applied, even if the species is not recognized in the initial test, treatment need not be delayed, since all of the above rickettsial species are susceptible to tetracycline or doxycycline. It has been recognized, however, that rickettsiae share antigens with several other bacterial species, including in some cases *Legionella* species. A genome that recognizes *Legionella* species should not be included in any test for rickettsiae, since the antibiotic of choice for *Legionella* species is erythromycin, which is not effective against rickettsiae (Hechemy et al., Ann. N.Y. Acad. Sci. 590: 1-586, 1990).

LITERATURE CITED

1. Fukuda, T., T. Sasahara, S. Yamamoto, N. Kawabata, and Y. Minamishima. 1985. Rickettsiae and rickettsioses in southern Kyushu, Japan, p. 357-363. In J. Kazar (ed.), Proceedings of the 3rd International Symposium on Rickettsiae and Rickettsial Diseases. Slovak Academy of Sciences, Bratislava, Czechoslovakia.
2. Grayston, J. T., C.-C. Kuo, L. A. Campbell, and S.-P. Wang. 1989. *Chlamydia pneumoniae* sp. nov. for *Chlamydia* sp. strain TWAR. Int. J. Syst. Bacteriol. 39:88-90.
3. Høllten, C. A., M. Ristic, D. L. Huxsoll, and G. Rapmund. 1982. *Rickettsia sennetsu* in human blood monocyte cultures: similarities to the growth cycle of *Ehrlichia canis*. Infect. Immun. 35:314-319.
4. Holland, C. J., M. Ristic, A. I. Cole, P. Johnson, G. Baker, and T. Goetz. 1985. Isolation, experimental transmission, and characterization of causative agent of Potomac horse fever. Science 227:522-524.
5. Holland, C. J., E. Weiss, W. Burgdorfer, A. I. Cole, and I. Kakoma. 1985. *Ehrlichia risticii* sp. nov.: etiologic agent of equine monocytic ehrlichiosis (synonym, Potomac horse fever). Int. J. Syst. Bacteriol. 35:524-526.
6. Krieg, N. R., and J. G. Holt (ed.). 1984. Bergey's manual of systematic bacteriology, vol. 1, p. 687-739. The Williams & Wilkins Co., Baltimore.
7. Myers, W. F., and C. L. Wisseman, Jr. 1980. Genetic relatedness among the typhus group of rickettsiae. Int. J. Syst. Bacteriol. 30:142-150.
8. Peter, O., J. C. Williams, and W. Burgdorfer. 1985. *Rickettsia helvetica*, a new spotted fever group rickettsia: immunochemical analysis of the antigens of 5 spotted fever group rickettsiae, p. 99-108. In J. Kazar (ed.), Proceedings of the 3rd International Symposium on Rickettsiae and Rickettsial Diseases. Slovak Academy of Sciences, Bratislava, Czechoslovakia.
9. Phillip, R. N., E. A. Casper, R. L. Anacker, J. Cory, S. F. Hayes, W. Burgdorfer, and C. E. Yunker. 1983. *Rickettsia bellii* sp. nov.: a tick-borne rickettsia, widely distributed in the United States, that is distinct from the spotted fever and typhus biogroups. Int. J. Syst. Bacteriol. 33:94-106.
10. Raoult, D., P. J. Weiller, A. Chagnon, H. Chaudet, H. Gallais, and P. Casanova. 1986. Mediterranean spotted fever: clinical, laboratory and epidemiological features of 199 cases. Am. J. Trop. Med. Hyg. 35:845-850.
11. Raoult, D., P. Zuchelli, P. J. Weiller, C. Charrel, J. L. SanMarco, H. Gallais, and P. Casanova. 1986. Incidence, clinical observations and risk factors in the severe form of Mediterranean spotted fever among patients admitted to hospital in Marseilles 1983-1984. J. Infect. 12:111-116.
12. Ristic, M., D. L. Huxsoll, N. Tachibana, and G. Rapmund. 1981. Evidence of a serologic relationship between *Ehrlichia canis* and *Rickettsia sennetsu*. Am. J. Trop. Med. Hyg. 30:1324-1328.
13. Stackebrandt, E., R. G. E. Murray, and H. G. Trüper. 1988. *Proteobacteria* classis nov., a name for the phylogenetic taxon that includes the "purple bacteria and their relatives." Int. J. Syst. Bacteriol. 38:321-325.
14. Uchida, T., X. Yu, T. Uchiyama, and D. H. Walker. 1989. Identification of a unique spotted fever group rickettsia from humans in Japan. J. Infect. Dis. 159:1122-1126.
15. Weisburg, W. G., M. E. Dobson, J. E. Samuel, G. A. Dasch, L. P. Mallavia, O. Baca, L. Maldelco, J. E. Sechrest, E. Weiss, and C. R. Woese. 1989. Phylogenetic diversity of the rickettsiae. J. Bacteriol. 17:4302-4306.
16. Weisburg, W. G., T. P. Hatch, and C. R. Woese. 1986. Eubacterial origin of chlamydiae. J. Bacteriol. 167:570-574.
17. Weisburg, W. G., C. R. Woese, M. E. Dobson, and E. Weiss. 1985. A common origin of rickettsiae and certain plant pathogens. Science 230:556-558.
18. Weiss, E., G. A. Dasch, Y.-H. Kang, and H. N. Westfall. 1988. Substrate utilization by *Ehrlichia sennetsu* and *Ehrlichia risticii* separated from host constituents by Renografin gradient centrifugation. J. Bacteriol. 170:5012-5017.
19. Weiss, E., J. C. Williams, G. A. Dasch, and Y.-H. Kang. 1989. Energy metabolism of monocytic *Ehrlichia*. Proc. Natl. Acad. Sci. USA 86:1674-1678.
20. Woese, C. R. 1987. Bacterial evolution. Microbiol. Rev. 51:221-271.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	20