

AD-A236 222



①

SMM OBSERVATIONS OF INTERSTELLAR ^{26}Al : A STATUS REPORT

W. R. Purcell and M. P. Ulmer
Department of Physics and Astronomy
Northwestern University > N00173-85-C-2501
Evanston, Illinois 60208

and

G. H. Share and R. L. Kinzer
E. O. Hulburt Center for Space Research
Naval Research Laboratory
Washington, D.C. 20375

1988
ABSTRACT

The discovery of radioactive ^{26}Al in the interstellar medium has provided direct evidence for the recent nucleosynthesis of intermediate-mass nuclei in the Galaxy. While modern nucleosynthesis theories suggest that ^{26}Al production may occur in a variety of astrophysical objects, they have not been able to identify the dominant production site of the observed ^{26}Al . Observationally, the greatest hope for identifying its source is through a comparison of the angular distribution of the 1.809 MeV radiation, emitted during the decay of ^{26}Al , with the angular distributions of the suggested production sites. Limits on the distribution of the 1.809 MeV radiation have been obtained from data collected by the *Solar Maximum Mission* Gamma-Ray Spectrometer by using the Earth as an occulting disk in its large field-of-view. These limits are: 1) the data are inconsistent with a point source origin of 1.809 MeV radiation located at $l = 0^\circ$, $b = 0^\circ$ at the 4.7σ confidence level; 2) the angular diameter for a uniform face-on disk distribution centered at $l = 0^\circ$, $b = 0^\circ$ is $> 10^\circ$ at the 4.1σ confidence level; and 3) the data are consistent with each of the diffuse distribution models studied representing ^{26}Al production by many individual events occurring throughout the Galaxy.

DISTRIBUTION STATEMENT A
Approved for public release
Distribution Unlimited

91-01449

INTRODUCTION

The discovery of 1.809 MeV radiation from the decay of radioactive ^{26}Al in the interstellar medium (ISM) by *HEAO-3* (Mahoney *et al.* 1984), and the subsequent confirmation by *SMM* (Share *et al.* 1985a), represents the first direct observation of extra-solar radioactivity. These observations suggest the presence of $\sim 2-3 M_\odot$ of ^{26}Al in the ISM which, because ^{26}Al decays with a mean lifetime of 1.04×10^6 years, provides direct evidence that significant quantities of ^{26}Al have been synthesized in the Galaxy in the last 1 - 2 million years. An identification of the production site, or sites, of the observed ^{26}Al will provide a unique opportunity for directly comparing observation with theories of nucleosynthesis.

While nucleosynthesis was originally considered to occur primarily in supernovae (Burbidge *et al.* 1957), modern theories suggest that significant nucleosynthesis occurs in a variety

01 6 6 064

The GRS axis has remained oriented toward the Sun throughout most of its mission. Since the Sun passes near the Galactic Center in late December, the Galactic Center region is visible for ~ 3 months each year as it passes through the large GRS field-of-view. It was the presence of an annual modulation of the 1.809 MeV line flux which led to the original *SMM* detection of ^{26}Al ; however, significant constraints could not be placed on its angular distribution. It is the combined use of Earth occultation, the large GRS field-of-view, and the over eight years of nearly continuous data which makes a detailed study of the distribution of the 1.809 MeV radiation possible.

^{26}Al DISTRIBUTION MODELS

Since the GRS is not an imaging instrument, it can not directly measure the angular distribution of the 1.809 MeV emission. Rather, an angular distribution model must be assumed and then compared with the data, so the angular distributions representative of the suggested production sites must be identified. For this work, two scenarios for the production of the observed interstellar ^{26}Al have been explored: 1) the ^{26}Al is assumed to be the result of nucleosynthesis by many individual events occurring throughout the Galaxy, and 2) the ^{26}Al is assumed to be the result of nucleosynthesis in a single event taking place in the direction of the Galactic Center.

If the observed ^{26}Al was produced by many individual events occurring throughout the Galaxy, its distribution would then be expected to follow the Galactic distribution of its production site. The suggested production sites can be separated into two Galactic populations: 1) Population I objects, including type II supernovae, Of and Wolf-Rayet stars, and massive ($M \gtrsim 3 M_{\odot}$) red giants; and 2) Population II objects, of which novae are members. Figure 1 shows two longitude distribution models for each of these Galactic populations. These models have been normalized to the 1.809 MeV flux of $0.033 \text{ photons sec}^{-1}$ reported by Share *et al.* (1985b). The distribution models for Population I objects are based on observations of interstellar carbon monoxide (CO) and are labeled CO Model #1 and CO Model #2. CO Model #1, from Leising and Clayton (1985), is based on a model for the radial distribution of molecular clouds in the Galaxy while CO Model #2, from Dame *et al.* (1987), represents the observed CO flux integrated along the line of sight and in Galactic latitude within $\pm 1.25^{\circ}$. The distribution models for Population II objects are based on observations of novae in M31 and are labeled Nova Model #1 and Nova Model #2. Nova Model #1, also from Leising and Clayton (1985), is based on a model for the radial distribution of nova in M31 while Nova Model #2, from Mahoney *et al.* (1985), is based on the assumption that the distribution of Galactic novae follows the visual luminosity.

If the observed ^{26}Al was produced by a single event, its distribution would be expected to follow roughly that of a face-on disk centered on the production site. The angular size of the disk would depend on the distance and age of the event. Two models were selected to examine this case; the Point Source Model and the Disk Model. The Point Source Model represents a point source of 1.809 MeV radiation while the Disk Model represents a uniform, face-on disk distribution having an angular diameter of $\sim 10^{\circ}$. From von Ballmoos *et al.* (1987), the centroid of the 1.809 MeV emission was found to be consistent with the Galactic Center, so the centroids of these models were located at $l = 0^{\circ}$, $b = 0^{\circ}$. These models were also normalized to the 1.809 MeV flux reported by Share *et al.* (1985b).




Figure 1: The ^{26}Al Galactic longitude models used in this work, normalized to the 1.809 MeV flux reported by Share *et al.* (1985b).

ANALYSIS TECHNIQUE

Using the Earth as an occulting disk in the intrinsically large GRS field-of-view, a technique of data selection has been developed which significantly improves the GRS angular resolution. This method, which can be applied to nearly any celestial position, also significantly reduces the instrumental background features produced by isotopes having half-lives $\gtrsim 10^5$ of minutes. The data selection method consists of identifying spectra as either source or background measurements based on whether a selected celestial position was occulted by the Earth during the spectral accumulation. If the celestial position was occulted, the associated spectrum is considered to be a background measurement, otherwise the spectrum is considered a source measurement. Using this technique, data were selected to perform observations of celestial positions at five degree intervals along the Galactic plane within $\pm 70^\circ$ of both the Galactic Center and the Galactic Anti-center; the Galactic Anti-center positions were used to monitor any residual systematic effects. In order to optimize the signal-to-noise, the following data selection criteria were applied. First, the data were required to have been collected $> 10,000$ seconds after the last significant passage of the *SMM* satellite through the South Atlantic Anomaly. Second, all data collected during known transient events were excluded. Third, only data collected when the GRS field-of-view was within 40° of the Galactic Plane were used. Finally, for the observation of each celestial position, the source spectra were defined as those collected when the celestial position was between 1° and 13° above the horizon, while the background spectra were defined as those collected when the celestial position was between 1° and 13° below the horizon. Only occultation transitions for which there were both source and background measurements available were used. In order to fur-




Figure 2: The normalized effective field-of-view for the Galactic Center observation. The contour levels are at 10, 25, 50, 75, and 90% of the peak response.

ther reduce the systematic effects, the source minus background differences were performed for each occultation transition separately, and the resulting background subtracted spectra accumulated for all of the valid occultation transitions. The same procedure was followed for each celestial position.

Once the spectra had been selected and the differences performed for the observation of each celestial position, the effective GRS field-of-view was generated for each observation by estimating the sky exposure for each of the individual spectra used. The effective field-of-view for the observation of the Galactic Center is shown in Figure 2. The asymmetry seen in this figure, which is found to vary with celestial position, is attributed to the orbital geometry and to periods during which the GRS instrument was placed into an in-flight calibration mode during which no data are available.

RESULTS AND CONCLUSIONS

The final difference spectrum for the observation of each celestial position was fitted over the range 1.5 – 2.1 MeV by a model consisting of three gaussian lines, one at 1.809 MeV and two background lines, and a continuum term. The lines were fixed in energy and in width at the instrument resolution (95 keV at 1.8 MeV). Analysis of the resulting 1.809 MeV fluxes indicated the presence of a residual systematic effect at the $\lesssim 1 \times 10^{-4}$ counts $\text{cm}^{-2} \text{sec}^{-1}$ level. This systematic was attributed to a prompt ($\lesssim 1$ second) background line at 1.809 MeV produced by neutron interactions on ^{27}Al . It was found that this residual

Table 1:
Results of Statistical Analysis

²⁶ Al Distribution Model	Reduced χ^2	
	Galactic Center	Galactic Anti-center
Point Source	2.71 ($2 \times 10^{-4}\%$)	0.80 (77%)
Disk Source	2.40 ($4 \times 10^{-3}\%$)	0.84 (71%)
CO Model #1	1.24 (17%)	1.06 (38%)
CO Model #2	1.14 (28%)	0.90 (62%)
Nova Model #1	1.38 (8%)	1.01 (45%)
Nova Model #2	1.14 (28%)	1.13 (29%)

systematic effect could be reduced below the statistical level by using the magnetic L-shell parameter as a measure of the cosmic-ray environment of the spacecraft during the spectral accumulations. The results of the spectral fits for each of the celestial positions, with the systematic effect removed, are shown in Figure 3. This figure also shows the expected signal for each of the ²⁶Al distribution models discussed above. As can be seen from the data in the Galactic Anti-center direction, most of the residual systematic effects have been eliminated. Since the final difference spectra for any two celestial positions may have used some of the same individual spectra, they are not statistically independent. A Monte Carlo simulation was used to estimate the covariance between adjacent positions; the errors shown in Figure 3 have been corrected for this covariance term.

A χ^2 test was applied between the data and each of the distribution models. The resulting reduced χ^2 , for 29 degrees-of-freedom, and the corresponding probability for each distribution model is given in Table 1. Included in Table 1 are the results of the analysis of the Galactic Anti-center observations to demonstrate the validity of the analysis method. The results of the statistical analysis are that: 1) the data are inconsistent, at the 4.7σ confidence level, with a point source origin of 1.809 MeV radiation located at the Galactic Center, 2) the angular diameter of diffuse emission (uniform face-on disk) centered on the Galactic Center is $> 10^\circ$ at the 4.1σ confidence level, and 3) the data are consistent with each of the CO and nova angular distribution models studied.





Figure 3: The background corrected counting rates for observations of positions in the direction of the Galactic Center (*upper*) and the Galactic Anti-center (*lower*). Note that this does *not* represent the *distribution* of the 1.809 MeV radiation. Also shown are the expected responses for each of the ^{26}Al distribution models.

The GRS observation of the Galactic Center using this technique, for which the effective field-of-view was $\sim 22^\circ$ FWHM (see Figure 2), is found to be $(1.1 \pm 1.0) \times 10^{-4}$ photons $\text{cm}^{-2} \text{sec}^{-1}$. This value is consistent with the fluxes of $(1.4 \pm 0.9) \times 10^{-4}$ and $(1.3 \pm 0.9) \times 10^{-4}$ photons $\text{cm}^{-2} \text{sec}^{-1}$ reported by Mahoney *et al.* (1986) and MacCallum *et al.* (1987), respectively, and is $\sim 1.9 \sigma$ below the flux of $(6.4 \pm 2.6) \times 10^{-4}$ photons $\text{cm}^{-2} \text{sec}^{-1}$ reported by von Ballmoos *et al.* (1987). The $\sim 2.3 \sigma$ excess seen in Figure 3 for the observation of the -20° longitude position would seem to suggest the presence of a concentrated source of 1.809 MeV emission in this direction. Due to the statistics available, however, and since the data are well fitted by the diffuse models, no significant conclusions about this excess can be made at this time. Work is in progress to study positions out of the Galactic plane in an attempt to place constraints on the latitude extent of the 1.809 MeV emission.

ACKNOWLEDGEMENTS

We wish to thank Steve Matz and Mark Leising for their valuable comments and suggestions throughout this analysis, and Dan Messina for his direction of the production analysis and careful screening of the data. We would also like to thank Tom Dame for supplying the data from his CO observations. This work was supported by NASA contracts S-10987-C at NU and S-14513-D and NRL.

REFERENCES

- Arnett, W. D. 1969, *Ap. J.*, **157**, 1369.
- Arnould, M., Nørgaard, H., Thielemann, F.-K. and Hillebrandt, W. 1980, *Ap. J.*, **237**, 931.
- Burbidge, E. M., Burbidge, G. R., Fowler, W. A., and Hoyle, F. 1957, *Rev. Mod. Phys.*, **29**, 547.
- Clayton, D. D. and Leising, M. D. 1987, *Phys. Rev.*, **144**, 1.
- Dame, T. M., *et al.* 1987, *Ap. J.*, **322**, 706.
- Dearborn D. S. P. and Blake, J. B. 1984, *Ap. J.*, **277**, 783.
- Forrest, D. L., *et al.* 1980, *Solar Physics*, **65**, 15.
- Hillebrandt, W., Thielemann, F.-K. and Langer, N. 1987, *Ap. J.*, **321**, 761.
- Leising, M. D. and Clayton, D. D. 1985, *Ap. J.*, **294**, 591.
- MacCallum C. J., Hutters, A. F., Stang, P. D. and Leventhal, M. 1987, *Ap. J.*, **317**, 877.
- Mahoney, W. A., Higdon, J. C., Ling, J. C., Wheaton, W. A. and Jacobson, A. S. 1985, *Proc. of the 19th International Cosmic Ray Conf.*, **1**, 357.
- . 1986, *BAAS*, **18**, 979.
- Mahoney, W. A., Ling, J. C., Wheaton, Wm. A. and Jacobson, A. S. 1984, *Ap. J.*, **286**, 578.
- Nørgaard, H. 1980, *Ap. J.*, **236**, 895.
- Purcell, W. R. 1988, Ph. D. thesis, Northwestern University, Evanston.
- Share, G. S., Kinzer, R. L., Kurfess, J. D., Forrest, D. J., Chupp, E. L. and Rieger, E. 1985a, *Ap. J. (Letters)*, **292**, L61.
- Share, G. H., Kinzer, R. L., Chupp, E. L., Forrest, D. J., and Rieger, E. 1985b, in *Proc. of the 19th International Cosmic Ray Conf.*, **1**, 353.
- von Ballmoos, P., Diehl, R., and Schonfelder, V. 1987, *Ap. J.*, **318**, 654.